

ROCKWOOD CHEMICAL CO.

P. O. Box 34

ROCKWOOD BRAND

Brawley, California

CAUTION

Harmful if swallowed, inhaled or absorbed through skin! DO NOT BREATHE DUST. DO NOT GET IN EYES OR ON CLOTHING. In case of contact remove contaminated clothing, flush skin or eyes with water. For eyes, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. During commercial or prolonged exposure in dust mixing and loading operations wear clean synthetic rubber gloves and a mask or respirator of a type passed by U. S. Bureau of Mines for Thiodan and organic phosphate protection. Prolonged exposure will result in cholinesterase depression. Do not apply to or allow dust to drift into areas occupied by unprotected humans or animals. DO NOT CONTAMINATE FEED OR FOOD. Apply this product only as specified on the label.

For further information on the use of this product consult your State Agricultural Extension Service or Experiment Station.

DO NOT APPLY or allow dust to drift onto varieties of sorghum which are sensitive to phosphates.

NOTICE TO PHYSICIAN: Atropine sulfate is antidotal.

WASTE DISPOSAL: Dispose of wastes by burying in non crop land areas, or by burning.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER, must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, wildlife and birds. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STATE REG. NO. 10226-50048-AA

EPA EST. NO. 10226-CA-1

EPA REG. NO. 10226-41

Thi-Lox 3-3 Dust

ACTIVE INGREDIENTS:

Endosulfan* (hexachloro-hexahydro-methano-
2, 4, 3-benzodioxathiepin oxide 3.0%
Dimethyl (2, 2, 2-trichloro-1-hydroxy-ethyl
phosphonate)** 3.0%

INERT INGREDIENTS: 94.0%

TOTAL 100.00%

*ENDOSULFAN (THIODAN) IS T.M. REG. BY CANADIAN
HOECHST LTD. U.S. PAT NO. 2,799,685.

**DYLOX IS T.M. REG. BY FARBENFABRIKEN BAYER A.G.
CHEMAGRO CORPORATION LICENSEE. U.S. PAT NO. 529,
379.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER PRECAUTIONS ON THE
BACKSIDE PANEL.

DO NOT REUSE EMPTY CONTAINER.

DESTROY IT BY BURYING IT WITH WASTE.

C/R Printers P4903RV 72575

DIRECTIONS FOR USE

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THE LABEL. A thorough application should be made, covering the undersides of leaves as well as the top. This product (Thi-Lox 3-3 Dust) is prepared for use on the following crops against the listed insects. Apply when infestation first appears and repeat as necessary. Thi-Lox 3-3 Dust may be applied by ground or airplane equipment. For brevity days prior to harvest for prohibited use of this product will be in parenthesis ().

ALFALFA - (Grown for Seed) For control of alfalfa caterpillar, alfalfa webworm, Beet Armyworm, Lygus bug, Meadow spittlebug, Stinkbug, Tarnished plantbug, Variegated cutworm, Western yellow-striped armyworm apply 33 lbs. of product per acre. (14) DO NOT FEED treated forage to dairy animals being finished for slaughter. DO NOT USE in forage to be used commercially or to be sold interstate.

CABBAGE AND CAULIFLOWER - For control of Western yellow-striped armyworm, Imported cabbageworm, Variegated cutworm, Diamondback moth, Cabbage aphid, Cabbage looper, Cross-striped cabbageworm, Flea beetle, Harlequin Bug and Stinkbug use 33-50 lbs. of product per acre on cabbage. Do not apply after edible parts start to form. (21)

COTTON - For control of Cotton fleahopper, Cotton leafworm, Cotton bollweevil, Bollworm, Cotton leaf perforator, Cabbage looper, Armyworm (Western yellow-striped and beet), Lygus bug, Stink bug, Darkling ground beetle, Southern garden leafhopper, Black fleahopper complex and Saltmarsh caterpillar apply 25-50 lbs. of product per acre. DO NOT APPLY after bolls open. DO NOT graze meat or dairy animals in treated fields.

SAFFLOWER (SEED) - For control of Green Peach aphid, Dipterous leaf miner, Imported cabbageworm, Stink bug, apply 20-33 lbs. of product per acre. DO NOT APPLY AFTER flowering stage.

SUGAR BEETS - For control of Beet webworm, Green Peach Aphid, Variegated cutworm, Beet armyworm, Dipterous leaf miners, Meadow Spittle bug and Saltmarsh caterpillar apply 20-33 lbs. of product per acre. (14) DO NOT FEED treated tops which have been treated within 30 days of harvest to livestock.

ACCEPTED
AUG 25 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 10226-41

NET CONTENTS