

ROCKWOOD CHEMICAL CO.

P.O. Box 34

Brawley, California

ROCKWOOD BRAND

Zineb-Cryolite Dust No. 6-50

DIRECTIONS FOR USE

Rockwood Brand Zineb-Cryolite Dust No. 6-50 should be applied to dry foliage at times when the air is still and the temperature does not exceed 90 degrees F. Thorough and uniform coverage is absolutely essential for disease control. Begin applications when disease is first present and continue as long as disease is prevalent. To be used only by those fully trained or otherwise familiar with the handling of hazardous insecticides. For brevity, days indicate interval between final treatment and harvest.

RECOMMENDED USES

CANTALOUPEs, CUCUMBERS, WATERMELONS, SQUASH
For the control of Downy Mildew, Alternaria blight, Anthracnose, and Gummy stem blight and Cabbage looper - apply 30 to 45 pounds per acre as long as necessary. Seven days. Repeat at 7 - 10 day intervals.

NOTE: This product may leave a poisonous residue on plants to which it is applied. The U. S. Food and Drug Administration has established a minimum amount of such poisonous residue that may remain on raw agricultural products at harvest and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of this minimal amount.

ACTIVE INGREDIENTS

Zinc ethylene bisdithiocarbamate
(zineb)* 6.0%

Sodium Fluoaluminate (Cryolite) 50.0%

INERT INGREDIENTS 44.0%
Total 100.0%

*Registered trade name of Rohm and Haas Company.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EPA Reg. No. 10226-44

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation of nose, throat and skin. Do not breathe dust or spray mist. Wear clean protective clothing and bathe after each exposure. AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

USE ONLY ON SULFUR RESISTANT CANTALOUPEs. Burn empty bag, never re-use. When burning, work to windward to avoid danger from fumes.

NON-WARRANTY: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

C/R Printers - P4302RV-4225

Made in U.S.A.

APR 7 1964
10226-44

ROCKWOOD CO.

P.O. Box 34

ROCKWOOD

Zineb-Cryolite No. 6

DIRECTIONS FOR USE

Rockwood Brand Zineb-Cryolite Dust No. 6-50 should be applied to dry foliage at time when the air is still and the temperature does not exceed 90 degrees F. Thorough and uniform coverage is absolutely essential for disease control. Begin applications when disease is first present and continue as long as disease conditions prevail. To be used only by thoroughly trained operators familiar with the handling of hazardous insecticides. For brevity, days indicate intervals between final treatment and harvest.

RECOMMENDED USES

CANTALOUPEs, CUCUMBERS, WATERMELONS, SQUASH
For the control of Downy Mildew, Alternaria blight, Anthracnose, and Gummy stem blight and Cabbage looper - apply 10 to 45 pounds per acre as long as necessary. Seven days. Repeat at 7-10 day intervals.

NOTE: This product may leave a poisonous residue on plants to which it is applied. The U. S. Food and Drug Administration has established a maximum amount of such poisonous residue that may remain on raw agricultural products at harvest and it is the user's responsibility to see that there is no residue on such crops at harvest time in excess of the maximum limit.

ACTIVE INGREDIENTS

Zinc ethylene bisdithiocarbamate (zineb)*

Sodium Fluoaluminumate (Cryolite)

INERT INGREDIENTS

Total

*Registered trade name of Rockwood

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product in accordance with the label.

EPA Reg. No.