

ROCKWOOD CHEMICAL CO.

P. O. Box 34

Brawley, California

ACCEPTED

AUG 25 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10226-48

ROCKWOOD BRAND

MALATHION THIO

SULFUR 5-4-20

RECOMMENDED USES

BROCCOLI, CABBAGE, CAULIFLOWER: For control of cabbage looper, flea beetle and stink bug. Apply 15 to 20 lbs. on cauliflower before the edible parts start to form; 15 to 25 lbs. on broccoli and cabbage. 14 days.

COTTON: For control of aphids, bollworm, cabbage looper, cotton leaf perforator, cotton stink bug, lygus bug. Apply 10 to 30 lbs. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

HEAD LETTUCE: For control of cabbage looper, aphids. Apply 20 to 25 lbs. product per acre. Do not apply within 14 days of harvest. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock. Do not apply more than 3 times after thinning.

MELONS: For control of aphids, cucumber beetle, striped cabbage flea beetle. Apply 20 to 35 lbs. per acre. 1 day.

NOTICE: Recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of ROCKWOOD CHEMICAL COMPANY, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practice. THE BUYER, must assume all responsibility including injury and/or damage, resulting from its misuse as such or in combination with other products.

STATE REG. NO. 10226-50033AA

C-R Printers-P5417 Rev.-7-18-75

ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithio-phosphate of diethyl mercaptosuccinate . .	5%
Endosulfan (Hexachloro-hexahydro-methano 2, 4, 3-benzodioxathiepin oxide . .	4%
Sulfur	20%
INERT INGREDIENT	71%
Total	100%

*Cythion, the premium grade Malathion, is T.M. Reg. U.S. Pat. Off. American Cyanamid Co.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NET CONTENTS _____

EPA EST. NO. 10226-CA-1

EPA REG. NO. 10226-48

DIRECTIONS FOR USE

ROCKWOOD BRAND MALATION THIO SULFUR 5-4-20 should be applied to dry foliage at time when the air is still, using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops. To be used only by thoroughly trained operators familiar with the handling of hazardous insecticides. For brevity, days indicate interval between final treatment and harvest.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking, wear clean clothing. Avoid contamination of feed and foodstuffs. Do not allow drift to areas occupied by unprotected humans or animals. To protect fish and wildlife, do not contaminate streams, lakes or ponds. Workers entering treated areas the day of application should wear protective clothing.