


10272-1A

FRONT PANEL



NOTICE
 KEEP FROM FREEZING. If this product is stored in cold or freezing weather it may separate or thicken. This will not harm the performance. Before using, warm to room temperature and stir thoroughly.

MANTEK
MAN-WEED X-40

Low Volatile Selective

Kills Broadleaf Weeds and
 Woody Plants

ACTIVE INGREDIENTS:
 Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid 24.5%
 Isooctyl Ester of 2,4,5-Trichlorophenoxyacetic Acid 11.7%

INERT INGREDIENTS: 63.8%

CAUTION: (18 pt. type) KEEP OUT OF REACH OF CHILDREN (12 pt tp)
 (See left panel for additional precautions.)

Contains 1 1/3 lbs. of 2,4-D acid equivalent plus 2/3 lb. 2,4,5,-T acid equivalent per gallon minimum. Contains equivalent of 16.3% 2-4-Dichlorophenoxyacetic acid and 8.1% Trichlorophenoxyacetic acid.

MANTEK CORPORATION
 12 2 Loop 12
 Irving, Texas 75060

LEFT PANEL

CAUTION:
 Avoid contact with skin, eyes or clothing. Harmful if swallowed. In case of contact flush eyes with plenty of water; wash skin with soap and water. Avoid breathing spray mist. NOT allow solution or spray drift to contact desirable plants. Among crops highly susceptible to 2,4-D and 2,4,5-T injury are cotton, tobacco, blackeyed peas, beans, tomatoes, melons, other vegetables, grapes fruit trees and some ornamentals. (Coarse sprays are less likely to drift than are fine mist sprays.) It is difficult to completely clean 2,4-D and 2,4,5-T from sprayers or other equipment, therefore do not use such sprayers and other equipment for purposes where even trace amounts of this chemical may cause injury. Do not reuse shipping containers; destroy by crushing and burying. Avoid contamination of water intended for irrigation and domestic use. Do not transport or store near seeds, fertilizers, insecticides or fungicides. Vapors from this product may injure susceptible plants in the vicinity. Do not use or store near heat or open flame. Consult agriculture experiment station or extension service weed specialists for specific recommendations as to dosage and time of application and for control of other weeds and woody plants. Note: Keep Spray tank agitated.

LEFT PANEL - Continued

Apply only to areas that will not be grazed after treatment by milking animals or animals being finished for slaughter.

DO NOT STORE WITH FOODSTUFFS

RIGHT PANEL -

Controls -

Buttercups	Ground Ivy	Sweetgum	Elm
Chervil	Henbit	Willows	Many Others
Chickweeds	Knotweed	Shepherds Purse	
Chicory	Oxalis	Sow Thistle	
Cinquefoil	Pennyworts	Speedwell	
Clovers	Plantains	Spotted Spurge	
Dandelions	Self-Heal	Wild Carrot	
Poison Ivy	Heal-All	Wild Onion	
Poison Oak	Sheepsorrel	Yellow Rocket	
Brambles	Oaks	Locust	
Honeysuckle	Chokecherry	Plums	
Cottonwood	Sassafras	Sumac	
Aspens	Wild Roses	Maples	

DIRECTIONS:

Lawn and Turf - - To control dandelions, chickweed, plaintain, Japenese Clover, oxalis, wild onions and certain other weeds, dilute 2 quarts of MANTEK MAN-WEED X-40 with 100 gallons of water to treat one acre. Wet foliage thoroughly. Water thoroughly before application. Do not apply during heat of day. If grass begins to yellow, water thoroughly and fertilize. Do not apply to grass only 3 or 4 months old. Temporary burning or yellowing may occur to Bent and Fescue grasses. Do not use on St. Augustine, clover, Dichondra or Lippia lawns. Oxalis - - Repeated treatment may be necessary. Wild onions - Treat while young and before bulbils are formed, usually in late fall or early spring.

For Docks, Thistles, Morning Glory, in fence rows and rights-of-way - - Mix 1 gallon to 100 gallons of water to treat 1 acre. Wet plants thoroughly and repeat when necessary.

For woody plants such as Poison Oak, Brambles, Poison Ivy, Sumac, Honeysuckle - - Mix 8 ounces per gallon of water and drench thoroughly while plants are in full leaf. Repeat application in four weeks if necessary.

To Kill Stumps - - Mix 1 pint in one quart of Kerosene or Diesel Oil. Spray all of the stump area thoroughly.

USDA REG. NO. _____