

GENERAL USE PRECAUTIONS

Do not plant any root crop not listed on this label in rotation with PCNB treated soils. This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

RECOMMENDATIONS AND DIRECTIONS FOR USE

BEANS (Root and Stem Rot - Rhizoctonia solani): FURROW - Application rates are to 14,500 linear feet of row for bush type beans and to 8,430 linear feet of row for pole beans. Apply 2.3 qts. in 8 to 10 gallons of water. Spray planting furrow and covering soil at planting time. Avoid applying directly to seed; otherwise delayed emergence may occur. IN CALIFORNIA AND ARIZONA apply 3.4 pints in 8 to 10 gals. of water. Apply in-furrow and mix with covering soil at planting. Use lower rate for lighter soils. Avoid application to bare seed.

SNAP AND DRY BEANS (White Mold-Sclerotinia sclerotiorum): Use 1 gal. mixed in 8 to 10 gals. of water and apply to 14,500 linear feet of row for bush type beans or to 8,430 linear feet of row for pole beans. Spray in an 8" wide band, centered on the row, immediately after or at time of seeding. Avoid applying directly to seed; otherwise delayed emergence may occur. (Where dinitro type pre-emergence herbicides are used, add 1 gallon to the final herbicide spray mix and apply simultaneously). If disease is severe, application may be repeated at 2 to 3 week intervals, using 1 or 2 nozzles per row and directing spray at base of plants. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

COTTON (Damping Off - Rhizoctonia solani): Use 1/2 to 1 gallon per acre in at least 15 gallons of water (per 13,000 linear feet of row based on 40" spacings). Spray on seed and surrounding soil at planting time. Do not allow the feeding or grazing of cotton foliage by livestock.

GARLIC (White Rot - Sclerotium cepivorum): Use 10 gals. mixed in 100 gals. of water and apply to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter, using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the seed as it is dropped. Direct the rear nozzle to spray the covering soil.

(Continued on right panel)

LOT No. _____

Net Contents _____

Rinse empty container thoroughly with water and discard it.

Cameraman's Note

Poor Copy

PROFESSIONAL TERRACLOR* 2 E

Emulsifiable Liquid-Soil Fungicide

CONTROLS CERTAIN SOIL-BORNE DISEASES OF BEANS-COTTON-GARLIC-POTATOES-LAWNS -ORNAMENTALS

ACTIVE INGREDIENT:
Pentachloronitrobenzene

23.9%
76.1%
100.0%

INERT INGREDIENTS:

*Terraclor is a registered trade mark of Olin Corporation.

CAUTION

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED.

AVOID INHALING SPRAY MIST.

THIS MATERIAL MAY CAUSE SKIN IRRITATION.

WASH HANDS AND FACE THOROUGHLY AFTER USING.

DO NOT CONTAMINATE FEED OR FOODSTUFFS.

COMBUSTIBLE—Do not use or store near heat or open flame.

PROTECT FROM TEMPERATURES BELOW 32°F.

*Product crystallizes at lower temperatures. Do not store at higher temperatures and mix thoroughly to solve crystals and assure uniformity.

PROFESSIONAL CHEMICAL CO., INC.

4517 YALE ST.

HOUSTON, TEXAS 77018

EPA Reg. No. 10290-21

692-7175

RECOMMENDATIONS
POTATOES (Rhizoctonia) with 100 to 150 gal. and cross disc to the last discing. of the soil. It is de
ROW (in-furrow) AP to total approximately all the soil that sure to the planter work acre); one cone net of spray per acre; Spray pattern should
ORNAMENTALS
AZALEA AND CAM under plants at a r osening buds and reg
LAWNS (Brown Patch Bermuda and similar Mix 1 qt. in sufficient spray or any other c area should be light area is subjected to the area should be s ation of the grass occ above, using 1 1/2 qts. areas. Do not
NOTICE: Profe or implied, inc any particular contained on the Observe all products use

PROFESSIONAL TERRACLOR* 2 E

Emulsifiable Liquid-Soil Fungicide
CONTROLS CERTAIN SOIL-BORNE DISEASES OF
BEANS-COTTON-GARLIC-POTATOES-LAWNS
-ORNAMENTALS

ACTIVE INGREDIENT:
Pentachloronitrobenzene

23.9%
76.1%
100.0%

INERT INGREDIENTS:

*Terraclor is a registered trade mark of Olin Corporation.

CAUTION KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED.
AVOID INHALING SPRAY MIST.
THIS MATERIAL MAY CAUSE SKIN IRRITATION.
WASH HANDS AND FACE THOROUGHLY AFTER USING.
DO NOT CONTAMINATE FEED OR FOODSTUFFS.
COMBUSTIBLE—Do not use near fire, heat, or open flame.

Under the Federal Insecticide, Fungicide, and Rodenticide Act,
this product is registered for the pesticide use indicated on the label.
*Product crystallizes at lower temperatures. Store at higher temperatures and mix thoroughly to break down crystals and ensure uniformity of application.

PROFESSIONAL CHEMICAL CO., INC.
4517 YALE ST. HOUSTON, TEXAS 77018
692-7175
EPA Reg. No. 10290-21

RECOMMENDATIONS (Continued)

POTATOES (Rhizoctonia solani) - **BROADCAST APPLICATION:** - Apply 9 to 12½ gallons mixed with 100 to 150 gallons of water to one acre. Apply to the soil surface, prior to planting and disc and cross disc to a depth of 4 to 6 inches. A drag harrow is usually attached behind the discs on the last discing. It is very important to thoroughly mix the Terraclor into the tuber-forming zone of the soil. It is desirable to make treatment when the soil is slightly moist.

ROW (in-furrow) APPLICATION: Use 5 gals. per 12,400 linear feet of row mixed in sufficient water to total approximately 20 gallons of finished spray. Spray several points in the planting row to treat all the soil that surrounds and covers the seed piece and potato forming zone. Three nozzles attached to the planter work best; one fan nozzle ahead of opening discs (approximately 5 gals. of spray per acre); one cone nozzle placed above and slightly ahead of closing discs (approximately 10 gals. of spray per acre); one fan nozzle to rear of closing discs (approximately 5 gals. of spray per acre). Spray pattern should treat a 12 to 14 inch band centered on the row.

ORNAMENTALS

AZALEA AND CAMELLIA (Azalea Petal blight and Camellia flower blight): Apply as a drench under plants at a rate of 1½ qts. in 30 gallons of water per 150 sq. ft. Begin application prior to opening buds and repeat every 3 to 4 weeks through the bloom period.

LAWNS (Brown Patch - Rhizoctonia solani): For Southern or warm season lawns such as St. Augustine Bermuda and similar grasses: Make one application in the fall or spring at first indication of infection. Mix 1 qt. in sufficient water (10 - 15 gals.) to cover 1,000 sq. ft. Apply by means of a sprinkler car or any other convenient method of obtaining even coverage. Following application, the treated area should be lightly watered to insure movement through the grass to the soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be retreated 3-4 weeks later. Under certain growing conditions, a temporary discoloration of the grass occasionally may occur, and will disappear in a short time. For Dichondra: Treat as above, using 1½ qts. mixed in 40 gals. of water per 1,000 sq. ft. Do not graze treated areas. Do not feed clippings to livestock.

NOTICE: Professional Chemical Company, Inc. makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on this label.

Observe all cautions and limitations on labeling of all products used in mixtures.

Cameraman's Note

Poor Copy