

DOUBLE ACTION INSECT KILLER - This is a fast-acting formula that can be used as a contact spray against flying and crawling insects and as a residual insecticide for providing up to 4 weeks of control of roaches.

DIRECTIONS FOR USE

Shake well before each use. Use protective cap. Aim spray away from person. Push button to spray.

HOUSEFLIES, MOSQUITOES AND GNATS
 Indoor - Close windows and doors. Point nozzle upward. Direct spray next to a light source, especially windows and other light sources which attract these insects. In the room, wait until they leave the treated area. Keep the room closed for a least 15 minutes. Vent air to outside when you return.

TO REDUCE ANNOYANCE FROM FLYING INSECTS
 Outdoor - Before you go into a garden, spray the legs and flycatcher traps. Then spray the resting places. For lawns, use a spray nozzle. Hold container 12 to 18 inches from shrubs, lawns and grassy areas. Spray these areas with a slow sweeping motion at the rate of 1 to 2 seconds per 100 square feet of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dry before occupying the treated area. Repeat as necessary.

ROACHES, EARWIGS, GRAIN MITES, SOYBUGS, SPIDERS AND CENTIPEDES

Contact Spray - Spray thoroughly into hiding places such as under areas, cracks, crevices, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly into nests where possible. Repeat as necessary.

ROACHES, EARWIGS, GRAIN MITES, SOYBUGS, SPIDERS AND CENTIPEDES

Residual Spray - Spray baseboards, cracks and crevices around door and window frames and other hiding places such as closets in back of and under appliances, sinks and cupboards in the kitchen and bathroom. In basements, spray hiding places in cracks and crevices, lower walls and dark places. Spray at the rate of 1 to 2 seconds per linear foot at a distance of 12 to 14 inches from the surface. Repeat as necessary.

WASPS - Applications should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

ANTS - Apply to ant trails and around door sills, window frames and other places where ants may enter.

ORNAMENTAL PLANTS PESTS: Spray aphids exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs, Japanese beetles, minute weevils, larvae and oakworm larvae. Direct spray on insects by holding the container 12 to 18 inches from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides, covering foliage until slightly moist. Avoid heavy wetting. Repeat applications as necessary.

USE WITH CAUTION on the foliage of Flowering Quince, Juniper, Cypress, Mulberry, Ash, Rose, Camellia, Azalea, as certain varieties have shown some leaf browning.

When infested, spray smaller evergreens, deciduous trees and shrubs as rhododendrons, pines, spruces, hemlock, yews, holly, arbovitae, junipers, boxwood, laurel, lilac, spirea, dogwood, palm trees, mimosa, toyon, viburnum, oaks, maples, locust, forsythia, althea and roses for control of aphids and other pests listed above. Also spray infested flowering plants as peonies, iris, zinnia, snapdragons, ageratum, chrysanthemum, nasturtium, daisy, Japanese geraniums, marigolds, ivy, sago, broadleaf acacia for the same listed pests.

SHAKE WELL BEFORE USING

NET WT. 1 lb.

1014

Venuscide

"Residual" Insect Killer

FOR USE IN HOMES AND NON-FOOD AREAS OF MOTELS AND INSTITUTIONS

ACTIVE INGREDIENTS:

*Resmethrin
 5-(Phenylmethyl)-1-furanyl methyl
 2,2-dimethyl-3-(2-methyl-1-propenyl)
 cyclopropanecarboxylate 0.350% w/w
 Related compounds 0.040%

INERT INGREDIENTS 99.610%
 100.000%

*Cis/trans isomers ratio: max 30% cis and min. 70% trans.

*U.S. Patent No. 3,465,007

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops. Do not use in eating products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. **WARNING:** Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator.

E.P.A. Registration No. 10292-26

E.P.A. Est. No. 10792-IL-01

Manufactured by

VENUS LABORATORIES, INC.

855 Lively Boulevard, Wood Dale, IL 60191

INSTITUTIONAL SIZE

ACCEPTED
 JUN 11 1979
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide registered under EPA Reg. No. 10292-26

CAUTION: KEEP OUT OF REACH OF CHILDREN
 (Pesticide label text)

**100% NON FLAMMABLE
PROPELLANT**

Venuscide
Flying Insect Killer

ACTIVE INGREDIENTS:

* Resmethrin	
5-(phenylmethyl)-3-furanyl methyl 2,2-di-	
methyl-3-(2-methyl-1-propenyl)	cyclopro-
panecarboxylate	0.200% w/w
Related compounds	0.028%
**d-trans Allethrin (allyl homolog of	
Cinerin I)	0.400%
Related compounds	0.030%
Aromatic petroleum hydrocarbons	0.272%
Petroleum distillate	6.500%
INERT INGREDIENTS	92.570%
	100.000%

*Cis/trans isomers ratio: max 30% (±)
cis and min. 70% (±) trans.

*Resmethrin, SBP-1382

**Bioallethrin

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove birds, pets, and cover aquariums before spraying. Do not use on edible crops. Apply this product only as specified on this label.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the plant is not in operation. **WARNING:** Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator.

E.P.A. Registration No. 10292-27

E.P.A. Est. No. 10292-IL-01

Manufactured by

VENUS LABORATORIES, INC.

855 Lively Boulevard, Wood Dale, IL 60191

INSTITUTIONAL SIZE

DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray. Apply this product only as specified on this label.

FOR USE IN HOUSEHOLDS AND INDUSTRIAL PLANTS

KILLS FLIES, MOSQUITOES AND GNATS: Close all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room especially windows and other light sources which attract these insects. Fill the room with mist then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. Sweep up and destroy fallen insects. Repeat treatment as necessary.

KILLS WASPS AND HORNETS: Space spray - before applying, close all doors, windows and ventilator openings. In unfinished attics and crawl spaces where direct sprays cannot reach pests, point nozzle upward and fill the area with mist at the rate of 35 seconds per 1,000 cu. ft. of space. Leave treated area immediately after application is complete, closing off area for at least 1 hour. Ventilate area when treatment is complete. Repeat as necessary.

FOR CRAWLING INSECTS: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes. Spray thoroughly into cracks, moist areas, crevices, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible as a contact spray. Repeat application as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

OUTDOORS - REDUCE ANNOYANCE FROM FLYING INSECTS AROUND PICNIC AND RECREATIONAL AREAS, NGMES AND INDUSTRIAL PLANTS.

Kills mosquitoes, gnats (midges), house flies, horse flies, deer flies and flying moths (millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

NUISANCE PESTS - ON OR ENTERING BUILDINGS:

WASPS AND HORNETS: Direct spray, application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary. Spray directly on stray wasps that enter buildings.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

SPIDERS: Spray directly on spiders and repeat as necessary.

EARWIGS: Spray directly on earwigs in crevices and other hiding places of outdoor porches, patios, flower beds and doorway entrances. Repeat as necessary.

IMPORTANT: Sweep up all dead insects and deposit these sweepings into a closed garbage container within two hours after each treatment.

ACCEPTED
JUN 11 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10292-27
SHAKE WELL BEFORE USING

NET WT. 1 LB.

#1013

CAUTION: Keep out of reach of children.
Contents under pressure.
See other caution on Back Panel.