

**DIRECTIONS**

Use one ounce per 100 gallons of water. Use in only noncorrodible watering utensils (Founts).

For Automatic proportioners: Add 1¼ ounces of MAQUAT DLC 1490 - 45% per gallon of water to prepare a stock solution. Use stock solution through chemical proportioner at the rate of one ounce per gallon of water. Use continuously except when water vaccines are in use. See danger statement.

MAQUAT DLC 1490 - 45% is a highly concentrated blend of chemical agents formulated for use as an aid in the control of potability of poultry drinking water.

Net Contents

KEEP FROM FREEZING

**MAQUAT  
DLC1490-45%**

E.P.A. Reg. No. 10324 - 11

**Concentrated - Poultry Drinking Water Sanitizer**

**ACTIVE INGREDIENTS:**

- n-Alkyl (90% C<sub>14</sub>, 5% C<sub>12</sub> & 5% C<sub>16</sub>)  
dimethyl dichlorobenzyl ammonium chloride .. 42.5%
- n-Alkyl (90% C<sub>14</sub>, 5% C<sub>12</sub> & 5% C<sub>16</sub>)  
dimethyl ethyl ammonium bromide ..... 2.5%

**INERT INGREDIENTS:** ..... 55.0%  
100.0%

**DANGER**

**KEEP OUT OF REACH OF CHILDREN**

**See antidote statement and other precautions on the side panel.**

**Rinse empty container thoroughly with water and discard it.**



**Mason Chemical Company**

A Division of Schrade Batterson Co.

5253 WEST BELMONT AVENUE  
CHICAGO, ILLINOIS 60641  
312/ 282 0200

**DANGER**

**Keep out of reach of children.**

Not for use or storage in or around the home. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear Goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

Do not use this product when water vaccines or water fed medications are in use. Discontinue use in poultry drinking water when birds are receiving any medication such as vaccines or any drug in their drinking water, for example, Sulfonamides, etc. If in doubt, consult your veterinarian.


**FIRST AID**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:**

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.



# MAQUAT DLC1490-45%

E.P.A. Reg. No. 10324 - 11

## Concentrated - Poultry Drinking Water Sanitizer

### ACTIVE INGREDIENTS:

n-Alkyl (90% C <sub>14</sub> , 5% C <sub>12</sub> & 5% C <sub>16</sub> )	
dimethyl dichlorobenzyl ammonium chloride . . .	42.5%
n-Alkyl (90% C <sub>14</sub> , 5% C <sub>12</sub> & 5% C <sub>16</sub> )	
dimethyl ethyl ammonium bromide . . . . .	2.5%

INERT INGREDIENTS: . . . . .	55.0%
	<u>100.0%</u>

### DANGER

**KEEP OUT OF REACH OF CHILDREN**

See antidote statement and other precautions on the side panel.

Rinse empty container thoroughly with water and discard it.



## Mason Chemical Company

A Division of Schrade-Batterson Co.

5253 WEST BELMONT AVENUE  
CHICAGO, ILLINOIS 60641

312-282-0200

### DIRECTIONS

Use one ounce per 100 gallons of water. Use in only noncorrodible watering utensils (Founts).

For Automatic proportioners: Add 1 1/4 ounces of MAQUAT DLC 1490 - 45% per gallon of water to prepare a stock solution. Use stock solution through chemical proportioner at the rate of one ounce per gallon of water. Use continuously except when water vaccines are in use. See danger statement.

MAQUAT DLC 1490 45% is a highly concentrated blend of chemical agents formulated for use as an aid in the control of potability of poultry drinking water.

Net Contents

KEEP FROM FREEZING

### DANGER

**Keep out of reach of children.**

Not for use or storage in or around the home. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear Goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

Do not use this product when water vaccines or water fed medications are in use. Discontinue use in poultry drinking water when birds are receiving any medication such as vaccines or any drug in their drinking water, for example, Sulfonamides, etc. If in doubt, consult your veterinarian.

### FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

### NOTE TO PHYSICIAN:

Probable mucosal damage may be indicated by the presence of proteinuria. Do not use if proteinuria is present.