

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Ms. Elizabeth Tannehill
Regulatory Manager
Mason Chemical Company
721 W. Algonquin Road
Arlington Heights, IL 60005

NOV 29 2004

SUBJECT: Maquat 5.5-M
EPA Registration Number: 10324-80
Applications Dated: May 25, 2004
Receipt Date: June 2, 2004

Dear Ms. Tannehill:

The following amendment, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable with the conditions listed below:

- update label language
- add public health pests

Efficacy Data

Acceptable Studies

Data Requirement	Means of Support	Status
Hantavirus (Using Prospect Hill virus as surrogate)	Submitted study: MRID 457837-01	Acceptable @ 1:64 in 400 ppm hard water, 5% soil for 10 minutes

As stated previously, additional directions for use specific to treatment of surfaces contaminated with Hantavirus are required. Add the following text to your label on page six immediately below the direction for the use of this product against HBV, HCV, and HIV for the treatment of areas which may be contaminated with Hantavirus:

CONCURRENCES							
SYMBOL	7510C						
SURNAME	J. Jones						
DATE	11/29/04						

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Special Instructions for Cleaning and Disinfecting areas which may be infested with Hantavirus

- 1. Infection with Hantavirus occurs by inhalation of infectious materials. CDC recommends that persons involved with cleanup wear coveralls, (disposable, if possible), rubber boots or disposable shoe covers, rubber or plastic gloves, protective goggles, and a half mask air purifying, negative pressure respirator with a high efficiency particulate air (HEPA) filter or a powered air-purifying respirator (PAPR) with HEPA filter.
2. All potential infective waste material (including respirator filter) from cleanup operations that cannot be burned or deep buried on site must be double bagged in appropriate plastic bags. The bagged material must then be labeled as infectious (if it is to be transported) and disposed of in accordance with local requirements for infectious waste.
3. Rodent droppings and visible dust may be reservoirs for Hantavirus. If you are cleaning out a building that has been closed up, such as a cabin, shed or garage:
a. Air out the building for at least 30 minutes by opening windows and doors.
b. Leave the building while it is airing out.
c. Do not vacuum, sweep or dust. This may spread the virus through the air.
d. Thoroughly wet the contaminated areas with the product and allow to stand undisturbed for 10 minutes.
e. Carefully remove contaminated material and dispose by burial or burning. Contact your local and state health department for additional disposal methods.
f. Treat the surface again following the label directions and allow to stand undisturbed for 10 minutes.

Additional Label Comments

Make the following changes to your labeling:

A. Make the following revision to page one:

Revise the "If Swallowed" statement to include: "...treatment advice..."

B. Make the following revisions to page three:

- 1. Correct the statement (left column, tenth item) to read "...grease and grime."
2. Correct the statement (right column, twentieth item) to read "...bowling alleys."

C. Make the following revisions to page four:

1. Change "moonwalk" to read "moonwalks" and "obstacle coarse play" to read "obstacle course play."

Table with 8 columns and 3 rows. Header: CONCURRENCES. Rows: SYMBOL, SURNAME, DATE.

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

3/15

2. Delete the entire section from the Directions for Use which begins "Solution used to disinfect medical devices..." and ends: "...Must have clear instructions for mixing and use." The Agency currently does not classify disinfectants as either "low or medium level" activity.

D. Make the following revisions to page five:

1. Delete the following statement as this product is used at a different rate as a virucide. Delete "according to the virucidal qualification on hard inanimate surfaces" from the second statement on this page beginning: "This product is bactericidal according to the AOAC Use dilution Test..."

2. Delete the word "Virucidal" from the heading for the "Disinfection/*Virucidal Directions" as well as from the general disinfection use dilution chart. This product is used at two different rates for effectiveness against viruses, which is not in agreement with the disinfection directions.

E. Make the following revisions to page six:

1. Revise the directions for "Kills HBV, HCV and HIV on precleaned Environmental Surfaces..." by adding the abbreviation, HCV and complete name; Hepatitis C Virus to the end of this statement.

2. Change "moonwalk" to read "*moonwalks*" and "obstacle coarse play" to read "*obstacle course play*."

3. Change "For Odors Caused by Dogs..." to read as follows: "...mats, *litter* boxes..."

4. Revise the directions for "Disinfection of hard, Non-Porous Surfaces in Footbaths" by changing the order of the statements as follows: "...Brush or swab...to stand for 10 minutes. *For spray applications, use a coarse spray device. Do not breathe spray.* After the unit has been thoroughly disinfected..."

F. Make the following revision to page seven:

Add the following statement to the directions "To Disinfect Food Service Establishments Food Contact Surfaces": "*After use, all surfaces in the area must be thoroughly rinsed with potable water.*"

G. Make the following revision to page eight:

Revise the "Farm Premise Use" directions to read: "...scrub all treated feed racks, *mangers*..."

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

H. Make the following revisions to page nine:

1. Since the section on "Poultry and Swine Premise Sanitization" refers to disinfection, and the organisms of interest at these sites are killed at the disinfection rate, move the sections beginning "Site Preparation" and "Application and Use dilution..." into the disinfection section of the label. Also revise the "Poultry and Swine Premise Sanitation" to read as follows: change "wails" to read "walls," and change "...feeding end watering appliance" to read "feeding and watering appliances..."

2. Revise the "Footbath Sanitizer Directions" such that the rate is in agreement with the non food contact sanitizing data by indicating a contact time of 5 minutes. Also revise by changing the order of the statements as follows: "...Brush or swab...to stand for 5 minutes. *For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly sanitized...*"

I. Make the following revision to page ten:


Revise the heading "Carpets, Ccarpet cushions..." to read "Carpets, *Carpet* cushions..."

General Comments

As stated in the Agency letter of 4/28/04, the cited study MRID 455392-01 does not support a claim for *Penicillium variable*, ATCC 32333. This study has been deleted from your matrix. Do the same to the matrix in your files.

A stamped copy of the accepted labeling is enclosed. Submit one (1) copy of your final printed labeling before distributing or selling the product bearing the revised labeling. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3 (c)(5) and section 4 (a) when the Agency requires all registrations of similar products to submit such data.

If the above conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6)e. Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. Should you have any questions concerning this letter, please contact Tracy Lantz at (703) 308-6415.

Sincerely,

Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510C)

Enclosures: Stamped label, DER D302117
7510C:T.Lantz:11/15/04:10324-80h

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER, Keep Out of Reach of Children. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if absorbed through the skin. Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist. Do not get into eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash clothing before reuse.

(If container is 5 gallons or larger the following statement must appear on the label.)

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents.

(If the container is greater than one gallon use the following storage and disposal statements.)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL (Larger than 1 gal.) - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(If container is one gallon or less use the following storage and disposal statements)

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children and animals. Do not store on side. Avoid creasing or impacting of side walls. Do not reuse empty container. Wrap and discard in trash (or recycle).

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 10324-80

MAQUAT® 5.5-M

Disinfectant, Cleaner, Mildewstat, Fungicide, Sanitizer (nonfood contact surfaces) Virucide*, Deodorizer for Hospitals Nursing Home, Whirlpool, Home, Institutional, Industrial, School, Dairy, Equine, Poultry/Turkey, Farm, Veterinary, Restaurant, Food Handling and Process Areas, Federally Inspected Meat and Poultry Plants, Bar and Institutional Kitchen Use and for Harvesting & Handling Equipment. Effective in hard water up to 400 ppm hardness (Calculated as CaCO₃) in the presence of 5% serum contamination.

Formulated for effective Poultry Sanitation.

Formulated for effective Swine Premise Sanitation.

Formulated for effective Mushroom Farm Sanitation

ACTIVE INGREDIENTS:

- Octyl Decyl Dimethyl Ammonium Chloride 1.650%
- Didecyl Dimethyl Ammonium Chloride 0.825%
- Diocetyl Dimethyl Ammonium Chloride 0.825%
- Alkyl (C₁₄, 50%; C₁₂, 40%; C₁₆, 10%)
dimethyl benzyl ammonium chloride 2.200%

INERT INGREDIENTS: 94.500%

TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See left panel for additional precautionary statements

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advise. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

MASON CHEMICAL COMPANY

"THE QUATERNARY SPECIALISTS"

721 W. Algonquin Road
Arlington Heights, IL 60005
847-290-1621

Toll Free:
1-800-362-1855

E.P.A. Reg. No. 10324-80
E.P.A. Est. No. 10324-IL-1



NET CONTENTS

BATCH NO

5/5

This product contains no phosphorous.

This product is a phosphate free formulation designed to provide effective cleaning, deodorizing and disinfection in areas where housekeeping is of prime importance in controlling the hazard of cross contamination on treated surfaces.

This product maximizes (improves) labor results by effectively controlling odors.

This product is for use in kitchens, bathrooms, and other household areas.

When used as directed, this product will deodorize surfaces in restroom and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

Cross-contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but in schools, institutions and industry.

This product delivers non-acid disinfection performance in an economical concentrate.

This product is a concentrated Hospital Use disinfectant that is effective against a broad spectrum of bacteria, is virucidal*, (and) fungicidal, and eliminates odor causing bacteria when used as directed.

This product is an economical concentrate that can be used with a mop and bucket, trigger sprayers, sponge or by soaking.

This product will not leave a grit or soap scum.

Will not cause swelling of transducer membrane or harm compressor plates.

This product is a versatile Disinfectant/Sanitizer for Veterinarian, Veterinary Practice, Animal Care, Animal Laboratory, and Agricultural and Farm Premise applications.

This product is a complete, chemically balanced disinfectant/(sanitizer) that provides clear use solutions even in the presence of hard water.

(This product inhibits bacterial growth on moist surfaces and deodorizes by killing microorganisms that cause offensive odors.)

This product is recommended for use in household and commercial humidifiers. Use of this product will control unpleasant (malodors) odors.

This product is recommended for use as a disinfectant on hard, non-porous surfaces (at 850 ppm active). A potable water rinse is required after application on food contact surfaces.

This product is a broad spectrum disinfectant formulated for use in Ultrasonic Baths (Ultrasonic cleaning units).

This product is a versatile cleaner and broad-spectrum disinfectant formulated for use on bath and therapy equipment (Whirlpools).

Use this product to disinfect/sanitize non-porous salon/barber tools and instruments: (such as) combs, clippers, brushes, trimmers, razors, scissors, blades, tweezers, and manicure/pedicure instruments and footbath surfaces.

This product is a one-step disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1 & HBV) and inhibits the growth of mold and mildew and their odors when used as directed.

This product is a no rinse disinfectant cleaner that disinfects, cleans and deodorizes in one labor saving step.

This product neutralizes musty odors and tough odors from smoke, pet accidents, and spills on contact.

This product is specially formulated to effectively eliminate offensive odors cause(d) by mold and mildew.

This product controls (reduces) (eliminates) (neutralizes) (destroys) odors to make your home (kitchen) (bathroom) sanitary.

This product is effective against household germs and odors (caused) by animal waste, septic tank or sewage backup, smoke and bathroom and kitchen odors.

Use this product on the multi-touch surfaces responsible for cross-contamination.

This product is effective at controlling mold and mildew on shower curtains.

This product may be used in work areas such as tool rooms and garages for odor control and light duty cleaning.

This product provides long lasting freshness against tough (pet) odors such as odors from litter boxes and pet accidents

This product cleans, shines, deodorizes and disinfects all hard non-porous surfaces listed on the label. It inhibits the growth of mold and mildew, leaving bathrooms and kitchens clean and fresh smelling.

This product is recommended for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

This product is a multi-surface cleaner, deodorizer and disinfectant. Use on windows, mirrors, and other non-food contact glass surfaces.

Use this product to clean, sanitize and disinfectant non-porous personal protective safety equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors and ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Caution: Cleaning at the recommended 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

This product is an effective antimicrobial cleaner, designed for use by wholesale and retail florists, shippers and green houses.

When used as directed, this product will disinfect hard, non-porous surfaces, such as flower buckets, floors and walls of coolers, design and packing benches, and countertops.

This product may be relied on to deodorize coolers, buckets, garbage pails and other areas where obnoxious odors may develop.

Use this product to clean, disinfect and deodorize flower buckets, walls, floors of coolers, shippers, greenhouse packing areas, garbage pails and other areas where obnoxious odors may develop.

This product may be used as a general purpose antimicrobial detergent in florist shops, wholesale florist, shippers, green house packing areas and other commercial floriculture places for efficient cleaning and antimicrobial action against certain bacteria which cause:

1. Plugging of stems with slime, which reduces uptake of water for various flowers including roses, chrysanthemums, gladioli and tulips.
2. Production of ethylene gas, which may injure blooms of the various sensitive flowers including carnations, snapdragons, some orchids, baby's breath, sweet peas, freesia and alstroemeria.

4/15

This product is an effective one-step sanitizer-cleaner for use on non-food contact surfaces.

Floor cleaner.

One-step cleaner.

Cleans everyday kitchen messes.

Cleans non-food contact kitchen surfaces and food preparation areas.

Cleans and shines.

Great for use (on) (in) the (kitchen), (bathroom), (floors) and other household areas. Removes (eliminates) odors. Deodorizes. Safe for most surfaces.

Safe for most surfaces. For a cleaner, fresher household. Removes stains. Removes dirt. Non-staining. Clear formula.

Eliminates odors caused by bacteria (and) (mildew) (and) (non-fresh foods). Kills odor causing bacteria in the kitchen (bathroom).

Cleans everyday kitchen messes like dirt, grease and food stains. Cuts through tough grease and grime.

Escherichia coli (*E. coli*), *Salmonella choleraesuis* (*Salmonella*), and *Staphylococcus aureus* (*Staph*) are common germs found where food is prepared and stored. This product kills these (harmful) germs and helps prevent the spread of food borne contamination on treated kitchen surfaces listed on this label. A potable water rinse is required after application on food contact surfaces. This product is not for use on dishes, glassware or eating utensils.

Disinfects (and Sanitizes) kitchen surfaces (bathroom surfaces and floors). Sanitizes kitchen surfaces (bathroom surfaces and floors). Kills germs. Kills household bacteria. A potable water rinse is required after application on food contact surfaces.

Antibacterial. Kills Athlete's Foot fungus on bathroom surfaces.

Hospital use Disinfectant. Institutional Disinfectant (Sanitizer).

Kills common kitchen (bathroom) germs (and viruses*).

Effective sanitizer in the presence of 5% serum contamination. (This product is an effective disinfectant/sanitizer in the presence of 5% serum contamination.)

Effective sanitizer in the presence of soils. (This product is an effective sanitizer in the presence of soils.)

This product is effective against Citrus Canker and Bacterial Spot of Tomatoes through the treatment of the handling, harvesting, storage and transportation equipment.

This product is recommended for Poultry Premise Sanitation (Hatcheries)

Egg Receiving Area	Tray Dumping Area	Chick Processing Area
Egg Holding Area	Chick Holding Area	Chick Loading Area
Setter Room	Hatchery Room	Poultry Buildings

This product is recommended for Swine Premise Sanitation:

Farrowing Barns and Areas	Dressing Plants	Blocks
Waterers and Feeders	Loading Equipment	Creep Area
Hauling Equipment	Nursery	Chutes

Maquat® 5.5-M for use in:

- Hospitals, nursing homes, medical and dental offices and clinics, physician offices, operating rooms, isolation wards, and medical research facilities.

- Patient care rooms & facilities, recovery, anesthesia, Emergency Rooms, X-ray cat labs, newborn nurseries, orthopedics, whirlpools, footbath surfaces, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms.
- EMS & fire facilities, emergency vehicles, ambulance(s), police cars.
- Day care centers and nurseries, sick rooms.
- Acute care institutions, alternate care institutions, home healthcare institutions.
- Life care retirement communities.
- Restaurants, restaurants and bars, bars, cafeterias, institutional kitchens, fast food operations and food storage areas.
- Supermarkets, convenience stores, retail and wholesale establishments, department stores, shopping malls, gift shops, video stores, bookstores, dressing rooms and laundries, photocopy centers, bicycle shops, auto repair centers.
- Computer manufacturing sites, toy factories.
- Food establishments, coffee shops, donut shops, bagel stores, pizza parlors, liquor stores.
- Crime scenes and funeral homes, mortuaries, burial vaults, mausoleums, autopsy rooms.
- Police stations, courthouses, correctional facilities, jails, prisons, municipal government buildings, bus stations, train stations.
- Institutional facilities, laboratories, factories, business and office buildings, restrooms, hotels and motels, and transportation terminals.
- Public restrooms, public facilities, waysides, travel rest areas, shower rooms, shower stalls, bathrooms.
- Hotel, motels, dormitories.
- Kitchens, bathrooms and other household areas.
- Homes (households).
- Institutions, schools and colleges, churches, classrooms, community colleges, universities, athletic facilities and locker rooms, exercise rooms, exercise facilities, gyms, gymnasiums.
- Cosmetic manufacturing facilities, medical device manufacturing facilities, biotechnology firms, pharmaceutical manufacturing facilities.
- Heath clubs, spas, tanning spas/beds, footbath surfaces, massage/facial salons, hair/nail/pedicure salons, barber/beauty shops, salons, tattoo parlors.
- Museums, art galleries, post offices, performance/theater centers, banks, libraries, movie houses, bowling alleys.
- Recycling centers.
- Humidifier water tanks.
- Campgrounds, playgrounds, recreational facilities, picnic facilities, sports arenas, sports complexes.
- Food processing plants, USDA inspected food-processing facilities, dairy farms, hog farms, equine farms, poultry and turkey farms and egg processing plants.

meat/poultry processing plants, meat/poultry producing establishments, mushroom farms, rendering plants.

- Processing facilities for Fish, Milk, Citrus, Fruit, Vegetable, Ice Cream, and Potatoes and beverage plants.
- Tobacco plant premises.
- Veterinary clinics, animal life science laboratories, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops and other animal care facilities.
- Household and automotive garages, boats, ships, barges, campers, trailers, mobile homes, cars, trucks, buses, trains, taxis and airplanes.
- Cruise lines, airline terminals, shipping terminals, public transportation.
- Commercial florist and flower shops.
- Basements, cellars, bedrooms, attics, living rooms and porches.

This product may be used on washable hard non-porous surfaces such as:

- Food preparation and storage areas.
- Counters (countertops), stoves (stovetops), sinks (bathroom, kitchen), tub surfaces, non-food contact surfaces of Slurppy[®] machines, and exterior surfaces of appliances, refrigerators, and ice machines.
- Glass, metal, stainless steel, glazed porcelain, glazed ceramic, granite, marble, plastic, (such as polystyrene or polypropylene), sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome and vinyl.
- Enameled surfaces, painted woodwork (finished), Formica[®], vinyl and plastic upholstery.
- Foundations, steps, plumbing fixtures, finished baseboards and window sills.
- Tables, chairs, desks, bed frames, lifts, washable walls, cabinets, doorknobs and garbage cans, cuspidors and spittoons.
- Exhaust fans, refrigerated storage and display equipment, coils and drain pans of air conditioning and refrigeration equipment and heat pumps.
- Large, inflatable, non-porous plastic and rubber structures: animals, promotional items, moonwalk, slides, obstacle course play and exercise equipment.
- Picnic tables and outdoor furniture.
- Telephones and Telephone booths.
- Highchairs.
- Drinking fountains.
- Shower stalls, shower doors and curtains, bathtubs and glazed tiles, chrome plated intakes, toilets, toilet bowls, toilet bowl surfaces, urinals, empty diaper pails, portable and chemical toilets and latrine buckets, porcelain tile and restroom fixtures.
- Ultrasonic baths, whirlpools, whirlpool bathtubs.
- Kennels, kennel runs, kennel/cages floors, conductive flooring.

This product meets AOAC Use – Dilution Test Standards for hospital disinfectants at 400 ppm water hardness.

This product meets AOAC efficacy standards for hard surface non-food contact sanitizers.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing (*Duck*) Hepatitis B Virus.

This product has passed the Virucidal Effectiveness Test using Bovine viral diarrhea virus (BVDV) (Surrogate for human Hepatitis C virus).

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Human Coronavirus.

For larger areas such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfection.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Solution used to disinfect medical devices, instruments and equipment. This solution may be used as a cleaning agent, and once device is properly cleaned, solution will provide low or medium level of disinfection. Solution must be able to be used in normal ventilation areas, and non-toxic for breathing and will not cause harm to equipment or patients with normal contact after being disinfected. Must have clear instructions for mixing and use.

(Note for reviewer: For labels that list medical premises and metal and/or stainless steel surfaces, one of the following statements must be used:)

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

Or

This product is not for use on Medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is required.

Apply this product with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray (mist).

NOTE: With spray applications cover or remove all food products. Prepare a fresh solution at least daily or when visibly soiled or diluted. For heavily soiled areas, a preliminary cleaning is required.

DISINFECTION

PREPARATION OF USE SOLUTION: For water hardness up to 400 ppm add 2 ounces of this product per gallon (16 milliliters per liter) of water (1:64) (850 ppm active quat) to disinfect hard non-porous surfaces. Treated surfaces must remain wet for 10 minutes. OR mix a 2 ounce packet with 1 gallons of water.

This product is a Hospital Use Disinfectant at 2 ounces per gallon (16 milliliters per liter) (1:64 dilution) modified in the presence of 400 ppm hard water.

This product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces, according to the virucidal qualification on hard inanimate surfaces, modified in the presence of 5% organic serum against:

- | | |
|---|--|
| <i>Campylobacter jejuni</i> | <i>Proteus vulgaris</i> |
| <i>Corynebacterium ammoniagenes</i> | <i>Pseudomonas aeruginosa</i> |
| <i>Enterobacter aerogenes</i> | <i>Pseudomonas aeruginosa</i> (Antibiotic resistant) |
| <i>Enterobacter cloacae</i> | <i>Salmonella choleraesuis</i> |
| <i>Enterococcus faecalis</i> | <i>Salmonella typhi</i> |
| <i>Enterococcus faecium</i> (Vancomycin resistant) | <i>Serratia marcescens</i> |
| <i>Escherichia coli</i> | <i>Shigella dysenteriae</i> |
| <i>Escherichia coli</i> (Antibiotic resistant) | <i>Shigella flexneri</i> |
| <i>Escherichia coli</i> O157:H7 | <i>Shigella sonnei</i> |
| <i>Legionella pneumophila</i> | <i>Staphylococcus aureus</i> |
| <i>Klebsiella pneumoniae</i> | <i>Staphylococcus aureus</i> (Methicillin resistant) |
| <i>Klebsiella pneumoniae</i> (Antibiotic resistant) | <i>Staphylococcus epidermidis</i> (antibiotic resistant) |
| <i>Listeria monocytogenes</i> | <i>Streptococcus pyogenes</i> |
| <i>Proteus mirabilis</i> | <i>Xanthomonas axonopodis</i> pv. <i>Citri</i> |
| | <i>Xanthomonas campestris</i> pv. <i>Vesicatoria</i> |

Fungicidal against:

- | | |
|------------------------------------|-------------------------|
| <i>Trichophyton mentagrophytes</i> | <i>Candida albicans</i> |
|------------------------------------|-------------------------|

Hospital Disinfection Use Dilution Chart

Ounces of Product	Amount of Water
½ ounce	1 quart
1 ounce	½ Gallon
2 ounces	1 gallon
5 ounces	2½ gallons
10 ounces	5 gallons
20 ounces	10 gallons

***Virucidal Performance:** At 1¼ ounces per gallon of water use level, (540 ppm active quat) treated surfaces must remain wet for ten minutes. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution at least daily or more often if solution becomes diluted or soiled.

This product was evaluated and found to be effective in the presence of 5% blood serum against the following viruses on hard non-porous surfaces:

- | | |
|---|--|
| Avian influenza A/Turkey/Wisconsin | Newcastle disease virus |
| Canine Distemper | Porcine Respiratory & Reproductive Virus (PRRSV) |
| Herpes Simplex Type1 | Porcine Rotavirus |
| Herpes Simplex Type2 | Pseudorabies virus |
| HIV-1 (AIDS virus) | Transmissible Gastroenteritis (TGE) |
| Influenza A/Brazil Virus | Vaccinia virus |
| Infectious Bovine Rhinotracheitis virus (IBR) | |

At 2 ounces per gallon of water use level (850 ppm active quat), with treated surfaces remaining wet for 10 minutes, this product was evaluated and found to be effective in against the following viruses on hard non-porous surfaces:

- | | |
|-------------------------|--------------------------|
| Canine Coronavirus | Heptatitis C Virus (HCV) |
| Human Coronavirus | Hantavirus |
| Hepatitis B Virus (HBV) | |

Disinfection/Virucidal Directions: Apply use solution to hard inanimate, non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For

sprayer applica. is use a coarse spray device. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist).

Before using this product, food products and packaging materials must be removed from the room or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

General Disinfection and Virucidal Use Dilution Chart

Ounces of Product	Amount of Water
0.3125 ounces	1 quart
0.625 ounces	½ gallon
1¼ ounces	1 gallon
3.125 ounces	2½ gallons
6¼ ounces	5 gallons
12½ ounces	10 gallons

At 1¼ ounces per gallon of water in the presence of 5% blood serum and 400 ppm of hardness for a 2 minute contact time, this product was found to be effective against HIV-1 (AIDS Virus).

OR

At 1¼ ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water (CaCO₃) and 5% blood serum, this product was proven to be effective against HIV-1 (AIDS Virus). Treated surfaces must remain wet for 2 minutes. Prepare a fresh solution at least daily or when use dilution becomes diluted or soiled.

At 2 ounces per gallon of water in the presence of 100% serum and 400 ppm of hardness for a 10 minute contact time, this product was found to be effective against (Duck) HBV.

OR

At 2 ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water (CaCO₃) and 100% serum, this product was proven to be effective against (Duck) Hepatitis B Virus with a contact time of 10 minutes.

At 2 ounces per gallon of water in the presence of 400 ppm of hardness for a 10 minute contact time, this product was found to be effective against Hepatitis C virus (HCV).

OR

At 2 ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water (CaCO₃) and 5% organic load, this product was proven to be effective against Hepatitis C Virus (HCV) with a contact time of 10 minutes.

At 2 ounces per gallon of water in the presence of 400 ppm of hardness for a 10 minutes contact time, this product was found to be effective against Hantavirus.

OR

At 2 ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water (CaCO₃) and 5% organic load, this product was proven to be effective against Hantavirus with a contact time of 10 minutes.

For Schools, Industry and Non-Medical Institutional Uses: At 1¼ ounces per gallon of water, this product delivers *cleaning and germicidal effectiveness. Treated surfaces must remain wet for 10 minutes, then allow to air dry or wipe. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution at least daily or more often if solution becomes diluted or soiled. It is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Escherichia coli* and *Serratia marcescens*. The same AOAC tests used to confirm performance for hospitals were used.

9/15

***KILLS HBV, HCV and HIV ON PRECLE. D ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus (HBV) and human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

"SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HBV, HCV AND HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS."

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, or eye coverings. (Clean up should always be done wearing protective latex gloves, gowns, masks and eye protection.)

Cleaning Procedure: Blood and other body fluids (containing HIV-1, HBV & HCV) must be thoroughly cleaned from surfaces and objects before application of this product.

Disposal of Infectious Materials: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

Contact Time: Leave surface wet for 2 minutes and 10 minutes for HIV-1 and HBV, respectively. This 2 minute contact time will not control other common type of viruses and bacteria listed on this label. For HCV, leave surface wet for 10 minutes.

OR

For HIV, leave surfaces wet for two minutes with 1¼ ounces per gallon (10 milliliters per liter) (or equivalent use dilution) use solution in the presence of 400 ppm hard water (CaCO₃ and 5% blood serum). This contact time will not control other common types of viruses and bacteria. For HBV, leave surfaces wet for 10 minutes with 2 ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water (CaCO₃ and 100% serum). For HCV, leave surfaces wet for 10 minutes with 2 ounces per gallon (or equivalent use dilution) in the presence of 400 ppm hard water and 5% organic load.

Efficacy tests have demonstrated that this product is an effective bactericide, virucide and fungicide in the presence of organic soil (5% blood serum).

General Deodorization: To deodorize, add a 1¼ ounces of this product to 1 gallon of water (or equivalent use dilution) (540 ppm active quat). Excess material should be wiped up or allowed to air dry.

For Deodorizing Garbage Cans, Garbage Trucks, Industrial Waste Receptacles and Garbage Handling Equipment: It is especially important to preclean for the product to perform properly. Use 1¼ ounces per gallon of this product or other suitable detergent system for pre cleaning step. Then, apply a wetting concentration of 6 ounces of this product per gallon of water (or equivalent use dilution) (2,500 ppm quat active).

For Odors Caused by Dogs, Cats and Other Domestic Animals: Use on rugs, floors, walls, tile, cages, crates, mats, litter boxes, floor coverings, or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area. Then follow directions for "General Deodorization".

Fungicidal Activity: At 2 ounces per gallon use-level, (or equivalent use dilution) (850 ppm active quat) this product is effective in locker rooms, dressing rooms, shower and bath areas and exercise facilities against *Trichophyton mentagrophytes* (athlete's foot fungus) (a cause of Ringworm), *Candida albicans* on inanimate hard non-porous

surfaces in the presence of 5% blood serum and 400 ppm water hardness as CaCO₃. Treated surface must remain wet for 10 minutes. Allow to air dry.

Mold and Mildew Control Directions: Add 1¼ ounces per gallon (10 milliliters per liter) of water to control the growth of mold and mildew (*Aspergillus niger*) and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappear.

Cleansing of Body Surfaces and Body Orifices of Human Remains: To cleanse away skin secretions and accompanying malodor, to insure the removal of all soil and bloodstains, and to remove and reduce surface contamination, apply 2 ounces of this product to 1 gallon of water (850 ppm active) to the surfaces and body openings, natural or artificial. Allow a 10 minute contact time for optimal results. Bathe the entire body using sponge or washcloth. A soft brush may be employed on surfaces other than the face. Prepare a fresh solution for application of each remains.

To control the growth of mold and mildew on large inflatable non-porous plastic and rubber structures (animals, promotional items, moonwalk, slides, obstacle course play and exercise equipment): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 1¼ ounces per gallon (540 ppm active quat) of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not house employ equipment until treatment has been absorbed, set or dried.

Disinfection/Fungicide/Virucide for Barber/Salon Tools Directions: Immerse pre-cleaned barber/salon tools (such as combs, brushes, razors manicure/pedicure tools, clippers, tweezers, clipper and trimmer blades, and scissors) in 2 ounces per gallon solution (or equivalent use dilution) (850 ppm active quat) of the product. Completely immerse instruments and tools for at least 10 minutes. Rinse thoroughly and dry before use. Prepare a fresh solution at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 5 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Disinfection of Hard, Non-Porous Surfaces in Footbaths: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 2 ozs per gallon of water (or equivalent use dilution) (850 ppm active quat) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 10 minutes. After the unit has been thoroughly disinfected, rinse all cleaned surfaces with fresh water. For spray applications, use a coarse spray device. Do not breathe spray. Do not use equipment until treatment has been absorbed, set or dried.

NON-ACID TOILET BOWL (AND URINAL) DISINFECTION/CLEANER DIRECTIONS

Remove gross filth prior to disinfection.

From Concentrate: Add 1½ ounces to the toilet bowl (0.75 gallons) and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop, or sponge. Allow to stand for 10 minutes and flush.

From Use solution: Empty toilet bowl or urinal and apply 2 ounces per gallon (16 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow to stand for 10 minutes and flush. For sprayer application, use a coarse spray device.

For Heavy Duty Cleaning: Empty toilet bowl or urinal and apply 8 ounces per gallon (64 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow to stand for 10 minutes and flush. For sprayer application, use a coarse spray device.

Cleaning/Deodorizing Directions: Add 1¼ ounces per gallon of water to clean and deodorize windows, mirrors and non-food contact glass surfaces. Apply solution with a mop, cloth, sponge or mechanical sprayer so as to wet all surfaces thoroughly. Allow to air dry or wipe up excess liquid.

Ultrasonic Bath Disinfectant Directions: Use this product to disinfect hard (inanimate) non-porous non-critical objects compatible with Ultrasonic cleaning units. Pour (a) fresh solution of 2 ounces of this product per gallon of water (1:64) directly into bath chamber. Pre-clean heavily soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to the manufactures' (use) directions. Remove objects and wash (rinse) with sterile water (sterile water for injection) or allow to air dry. Prepare fresh solution for each use.

Disinfection of Hard Non-Porous Surfaces in Whirlpool Units: After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 2 ounces of this product for each gallon of fresh water added. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

For Disinfecting Hard, Non-porous (Fiberglass) Bath and Therapy Equipment: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the whirlpool unit, drain the water and refill with fresh water to just cover the intake valve. Add 10 ounces of this product for each 5 gallons of water (2 ounces per one gallon) (850 ppm active quat) in the unit at this point. Briefly start the pump to circulate the solution. Turn off pump. Wash down the unit sides, seat of the chair lift, and any/all related equipment with a clean swab, brush or sponge. Product to surface contact time must be at least 10 minutes for proper disinfection. After the unit has been thoroughly disinfected, drain solution from the unit and rinse any/all cleaned surfaces with fresh water. The unit is ready for reuse.

To Disinfect Food-Processing and Tobacco Premises: (For) floors, walls and storage areas, add 2 ounces of this product to gallon of water (or equivalent use dilution) (850 ppm active quat). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications use a coarse spray device. Allow to remain wet for 10 minutes, then remove excess liquid. For sprayer applications, use a coarse (mist or) spray. Before using this product, food products and packaging materials must be removed from area or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

To Disinfect Food Service Establishment Food Contact Surfaces: (For) countertops, appliances, and tables. Before using this product food products and packaging materials must be removed from area or carefully protected. Add 2 ounces of this product to 1 gallon of water (or equivalent use dilution) (850 ppm active quat). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes. Then remove excess liquid and rinse the surface with potable water. For sprayer applications, use a coarse spray (device). This product is not for use on dishes, glassware or eating utensils.

For use on non-food contact surfaces as a general disinfectant in the brewery industry: Use 2 ounces of this product per gallon of water (or equivalent use dilution) (850 ppm active quat). Follow the general disinfectant directions above.

Directions for Fogging: (For use in) dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using one quart per 1000 cubic feet of room area with a product solution containing 3 ounces product to 1 gallon of water (or equivalent use dilution) (1,200 ppm). Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed prior to reuse with potable water.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

Disinfection of Poultry/Turkey Equipment, Swine Quarters, Animal Quarters and Kennels: Prior to disinfection, remove all poultry, other animals and their feed from premises, trucks, coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 1¼ ounces of this product per gallon of water (or equivalent use dilution) (540 ppm active quat). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

Disinfection/Fungicide/Virucide of Poultry/Turkey Equipment, Swine Quarters, Animal Quarters and Kennels Directions: Remove all animals and feeds from premises, vehicles and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 2 ounces of this product per gallon of water (or equivalent use dilution) (850 ppm active quat). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After application, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set or dried.

11/5

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent, then rinsed with potable water before reuse.

HATCHERIES: Use 2 ounces of this product per gallon of water to treat the following hard non-porous surfaces: hatching, setters, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more. Rinse with potable water before reuse. Then allow to air dry.

VEHICLES: Clean all vehicles including mats, crates, cabs, and wheels with high-pressure water and this product. Use 2 ounces per gallon to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more and allow to air dry. For sprayer applications, use a coarse spray device.

DRESSING AND RENDERING PLANT USE (RENDERING PLANT/DRESSING PLANT DISINFECTANT DIRECTIONS): Disinfect equipment, walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the recommended use solution, 1½ ounces of this product per gallon of water (540 ppm active quat). Scrub to loosen all soils. Allow to soak for 10 minutes and thoroughly rinse all wetted and cleaned surfaces with potable water.

RENDERING PLANT/DRESSING PLANT DISINFECTANT/FUNGICIDE/VIRUCIDE DIRECTIONS: Cover or remove all food packaging material before disinfection. Remove gross soils. Apply to equipment, walls and floors in poultry and animal dressing plants with a solution of 2 ounces of this product per gallon of water (or equivalent use dilution) (850 ppm active quat). Equipment and utensils must remain wet for 10 minutes then thoroughly rinsed with potable water before operations are resumed. Disinfect all rooms, exterior walls and loading platforms of dressing plants. Thoroughly rinse all wetted and cleaned surfaces with potable water.

FARM PREMISE USE: Do not use in milking stalls, milking parlors or milk houses. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or transversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with a use solution of 2 ounces of this product per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, managers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

OR

FARM PREMISE DISINFECTION DIRECTIONS

For use in Equine, Dairy and Hog Farms:

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, chutes and other facilities and fixtures occupied or transversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.

5. Saturate all surfaces with the recommended disinfecting/Virucidal or fungicidal solution for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, managers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

Veterinary Practice (Clinics)/Animal Care Animal Laboratory/Animal Life Science Laboratory/Zoos/Pet Shop/Kennels/Breeding and Grooming Establishment/Tack Shops Disinfection Directions: For cleaning and disinfecting the following hard non-porous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals (and) feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces occupied or transversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 2 ounces of this product per gallon of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed set or dried.

COMMERCIAL FLORIST USE DIRECTIONS: To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare Use solution by adding (mixing) 2 ounces per gallon of water (1:64). For heavy-duty cleaning, add (mix) 8 ounces per gallon of water (1:16).

Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply Use solution to hard (inanimate) non-porous surfaces, thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge or sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface, (and) rub with brush, sponge or cloth. Do not breathe spray (mist).

Treated surfaces must remain wet for 10 minutes. Allow to air dry.

Prepare a fresh solution at least daily or when use solution becomes visibly dirty.

Citrus Canker Control and Control of Bacteria Spot of Tomatoes (Not Applicable in California): For prevention of Citrus Canker and Bacterial Spot of Tomatoes Diseases through treatment of equipment. Effective against *Xanthomonas axonopodis* pv. citri (Citrus Canker) and *Xanthomonas campestris* pv. vesicatoria (bacteria spot of tomatoes) at 2060 ppm active quat. Treat all trucks, vehicles, and equipment thoroughly at a dilution ratio of 1:27 (2060 ppm active quat). Treatment can be applied to trucks, attached trailers, field harvesting equipment; including cargo area, wheels, tires, under carriage, hood, roof, fenders and any other part of transportation equipment that can be taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using maximum rate 2060 ppm quat. For sprayer applications use a coarse spray. After use, all surfaces which come in contact with food or crop must be rinsed with potable water. Clothing should be either thoroughly rinsed or laundered before reuse. Footwear should be rinsed before reuse.

12/5

SANITIZING NON-FOOD CONTACT SURFACE DIRECTIONS

Preparation of Sanitizer Use Solution: Add 1 ounce per 2¾ gallons of water to sanitize hard non-porous environmental surfaces. Treated surfaces must remain wet for 5 minutes. Allow to air dry.

Staphylococcus aureus

Klebsiella pneumoniae

To Sanitize (and Deodorize): Apply use solution to hard inanimate, non-porous surfaces, thoroughly wetting surfaces as recommended and required with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist). Let stand for 5 minutes. Then wipe. Allow to air dry.

POULTRY (AND SWINE) PREMISE SANITATION

Site Preparation: The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated thoroughly clean all surfaces with soap or detergent and rinse with water.

Application and Use Dilution Poultry (and Swine) Premise: Remove all animals and feeds from premises, trucks, cars, coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 2 ounces of this product per gallon of water. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers and other equipment which may contact food or water with soap or detergent, and rinse with potable water before reuse. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

SHOE BATH SANITIZER (DIRECTIONS): (To prevent tracking harmful organisms into animal areas, and the packaging and storage areas of food plants.) Shoe baths containing one inch of freshly made solution should be placed at all entrances to buildings, hatcheries and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and place in 2 ounces of this product per gallon (16 milliliters per liter) of water solution for 5 minutes prior to entering area. Change the sanitizing solution in the bath at least daily or sooner if solution appears dirty.

SANITIZING HATCHERY ROOMS USING FOGGING DEVICES: Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Close room off so fog is confined to room to be treated. Mix 2 gallons this product to 2½ gallons water. Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with soap or detergent and rinse with water.

SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES: Mix 24 ounces of this product to 122 ounces of water. Fog 3-8 ounces of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. It is acceptable to fog setters and hatchers with 2 ounces per gallon solution of this product on an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

FOOD PROCESSING PLANTS USING FOGGING DEVICES

Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using 1 quart per 1000 cubic feet of room area with a solution containing 3 ounces of product to 1 gallon of water (1,200 ppm). Vacate the area of all personnel for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed prior to reuse with potable water.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

SALON/BARBER INSTRUMENTS AND TOOLS SANITIZER DIRECTIONS: Pre-clean soiled instruments and tools. Completely immerse water safe nonporous instruments and tools in a solution of 1 ounce per 2¾ gallons (or equivalent use dilution) (150 ppm active) of water for at least 5 minutes. Rinse thoroughly and dry before use. Prepare a fresh solution at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 5 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Footbath Sanitizer Directions: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 1 oz per 2¾ gallon of water (or equivalent use dilution) (150 ppm active quat) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 60 seconds. After the unit has been thoroughly sanitized, rinse all cleaned surfaces with fresh water. For spray applications, use a coarse spray device. Do not breathe spray. Do not use equipment until treatment has been absorbed, set or dried.

Humidifier Sanitizer Directions: Thoroughly clean water tank and filters. Add 2 ounces of this product per gallon of refill water. Not for use in heat or atomizing type humidifiers.

WATER/SMOKE DAMAGE RESTORATION (Not Applicable in California)

Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use. This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim, and frame lumber, tackless strip and paneling. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

Sewer backup & river flooding: During mitigation procedures, dilute 2 to 4 ounces of this product per gallon of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affective areas with a sprayer using a coarse spray tip, before and after cleaning and extraction.

Carpets, Carpet cushions and other porous materials such as sub floors, drywall, trim and frame lumber, tackless strip and paneling: For water damage from a clean water source, extract excess water. Test hidden area for colorfastness. Dilute 2 to 4 ounces of the product per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a sprayer using a coarse spray tip, to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for 10 minutes. Follow with a through extraction. Dry rapidly and thoroughly.

Special Instructions for Cleaning Carpet Against Odor Causing Bacteria: This product may be used in industrial and institutional areas such as homes, motels & hotel chains, nursing homes, schools and hospital. For use on wet cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for color fastness.

For portable extraction units: Mix 2 ounces of this product per gallon of water.

For truck mounted extraction machines: Mix 24 ounces of the product per gallon of water and meter at 4 gallons per hour.

For rotary floor machines: Mix 4 ounces of this product per gallon of water and apply at the rate of 300-500 sq. ft. per gallon.

Do not mix this product with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using this product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the carpet manufacturer's directions.

CARPET SANITIZER AGAINST ODOR-CAUSING BACTERIA, FOR HOME, INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE (Not Applicable in California)

This product sanitizes the carpet by controlling/reducing the growth of odor-causing bacteria. It can be used in industrial and institutional areas such as homes, motels, hotels chains, nursing homes and hospitals.

Vacuum carpet thoroughly prior to application. Mix 2 ounces of product per gallon of water. Follow the Injection and/or Extraction procedures as specified for any conventional steam cleaning equipment you are using. For rotary floor machines, mix 16 ounces per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon.

For use on washable synthetic fibers. Do not use on wool. Test color fastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to cloth, a water-based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

Note: This product should not be mixed with other cleaning products.

SMOKE DAMAGE RESTORATION (Not Applicable in California): Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use: This product is particularly suitable for use in smoke damage restoration situations against

odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim, and frame lumber, tackless strip and paneling. Follow directions as outlined in the Water Damage Restoration section. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

MUSHROOM FARM INDUSTRY USE DIRECTIONS

Site Preparation: The first step in any on going sanitation program should be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

Disinfection: Use 2 ounces of this product per gallon of water (or equivalent dilution). Wet all surfaces thoroughly. Treated surfaces should be allowed to remain wet for 10 minutes. Let air-dry. For heavily soiled areas, preclean first. Prepare a fresh solution for each use.

For Heavy Duty Cleaning: When greater cleaning is desired, use 4 ounces of this product per gallon of water (or equivalent dilution). Heavily soiled areas may require repeated cleaning before treatment.

DO NOT APPLY TO THE MUSHROOM CROP, COMPOST OR CASING. Rinse treated surfaces with potable water before they contact the crop, compost or casing.

**TWO OUNCE PACKET TO BE USED WITH MASTER
CONTAINER LABEL**

MIX EACH PACKET WITH 1 GALLON OF WATER

MAQUAT 5.5-M

E.P.A. Reg. No. 10324-80 E.P.A. Est. No. 10324-IL-1

NET CONTENTS: 2.0 FLUID OZ.

.....
ACTIVE INGREDIENTS:

Octyl Decyl Dimethyl Ammonium Chloride	1.650%
Didecyl Dimethyl Ammonium Chloride.....	0.825%
Dioctyl Dimethyl Ammonium Chloride.....	0.825%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%).....	
dimethyl benzyl ammonium chloride	2.200%
INERT INGREDIENTS:	94.500%
TOTAL	100.000%

.....

KEEP OUT OF REACH OF CHILDREN

DANGER

**SEE OUTER CONTAINER FOR PRECAUTIONARY STATEMENTS AND
USE DIRECTIONS**

DISPOSAL: Do not reuse container. Wrap and put in trash.

**MASON CHEMICAL COMPANY
"THE QUATERNARY SPECIALISTS"**

721 W. Algonquin Road
Arlington Heights, IL 60005

847-290-1621

Toll Free:
1-800-362-1855



5/5