

10324-99

10/16/2007

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

OCT 16 2007

Ms. Elizabeth Tannehill
Regulatory Manager
Mason Chemical Company
721 W. Algonquin
Arlington Heights, IL 60005

Subject: Maquat 10-PD
EPA Registration Number 10324-99
Application Dated June 6, 2007
Received Date June 18, 2007

Dear Ms. Tannehill:

The following label revisions submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable with the following comments.

Proposed Amendment:

- Label amendment and adding new organism

Labeling Comments:

1. Revise the claim in the left column of page five which begins; "Disinfects (and Sanitizes) kitchen surfaces (bathroom surfaces and floors). Sanitizes *non-food contact* kitchen surfaces..."
2. Revise the statement "Saturate surfaces with recommended disinfecting solution for a period of 10 minutes," located in the list of directions below in the "Animal Premise Use Directions" subsection on pages nine and ten. Delete "recommended" such that the statement is in agreement with PR Notice 2000-5, Mandatory Language. The following sets of directions require this revision:

Disinfection/Fungicide/Virucide of Poultry/Turkey Equipment, Swine Quarters, Equine Quarters Rendering Plants/Dressing Plants For use in Equine, Dairy, and Hog

Treatment of Animal Housing Facilities
Terrarium and Small Animal Cage Furniture Disinfection
Special Instructions for Inactivating Avian Influenza A

General Comments:

A stamped copy of the acceptable labeling is enclosed. Submit one (1) copy of your final printed labeling before distributing or selling the product bearing the revised labeling.

Should you have any questions or comments concerning this letter, please contact Drusilla Copeland at (703) 308-6224 or Velma Noble at (703) 308-6233

Sincerely,



Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510P)

Enclosure



MASON CHEMICAL COMPANY
 "The Quaternary Specialists"

721 W. Algonquin Road | Arlington Heights, IL 60005 | 847-290-1621 or 800-362-1855

E.P.A. Reg. No. 10324-99

E.P.A. Est. No. 10324-IL-1

MAQUAT® 10-PD

Net Contents:

Batch No:

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS
 AND DOMESTIC ANIMALS**

DANGER, Keep Out of Reach of Children. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get into eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

(If container is 5 gallons or larger the following statement must appear on the label.)

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents.

(If the container is greater than one gallon use the following storage and disposal statements.)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL (Larger than 1 gal.): Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. In addition, plastic containers may be disposed of by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(If container is one gallon or less use the following storage and disposal statements.)

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to small children. Do not store on side. Avoid creasing or impacting of side walls. Do not reuse empty container. Wrap and discard in trash (or recycle).

PELIGRO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

Disinfectant, Sanitizer (Non-Food Contact), Deodorizer, Fungicide, Mildewstat, Virucide* With Organic Soil Tolerance For Hospitals, Nursing Home(s), Whirlpool, Home, Hotels & Motels, Waterbed Conditions, Institutional, Industrial, School, Dairy, Equine, Poultry/Turkey Farm, Veterinary, Federally Inspected Meat and Poultry Plants, Bar and Institutional Kitchen.

Non-Acid Bathroom Cleaner.

Athletic Surface Disinfectant.

Small Fly Ovicidal Treatment.

Formulated for effective Poultry Sanitation, Federal Insecticide, Fungicide, and Rodenticide Act as

Formulated for effective Swine Pen Sanitation, Pesticide,

Formulated for effective Mushroom Farm Sanitation, registered under EPA Reg. No.

ACTIVE INGREDIENTS:

Alkyl (C₁₄ 60%, C₁₆ 30%, C₁₂ 5%, C₁₈ 5%)
 dimethyl benzyl ammonium chloride 5.0%

Alkyl (C₁₂ 68%, C₁₄ 32%)
 dimethyl ethylbenzyl ammonium chloride 5.0%

INERT INGREDIENTS: 90.0%

TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
 DANGER PELIGRO**

See left (back) (side) (right) panel (of label) (below) for additional precautionary and first aid statements.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

(Note to Reviewer: This information has been verified per DOT regulations. It is NOT required to be on the label but is being requested by a customer.)

Transportation Information

DOT Hazard Class: 8 Corrosive

DOT Proper Shipping Name: Disinfectant Liquid Corrosive (Quaternary Ammonium Compound), 8, UN1903, PGIII

ACCEPTED
 with COMMENTS
 in EPA Letter Dated:

OCT 16 2007

Sanitation, Federal Insecticide, Fungicide, and Rodenticide Act as Pesticide, registered under EPA Reg. No. 10324-99

3/14

(Note to reviewer: The Table of Contents will not be on any label. This is for our customer's reference only. The page numbers will change as the label changes.)

Table of Contents

Description	Page No.
Organism Guide	2
Marketing Claims	3
Disinfection	
General	7
Hospital/Healthcare	8
Deodorization/Mold & Mildew	9
Barber/Salon	10
Restroom/Bathroom	10
Food Processing	11
Animal Premises	11
Other	13
Non-Food Contact Sanitization	
Laundry	14
Food Processing	14
Animal Housing	15
Other	15
Food Contact Sanitization	
General	15
Public Eating	16
Food Processing	17

ORGANISM LIST

(Note: For reference use only. This is a list of all organisms covered by this label. Please see appropriate Directions for Use for proper use dilution.)

Disinfection Performance in Hospitals, Dental Offices, Nursing Homes, and Other Health Care Institutions: This product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum against:

<i>Botrytis cinerea</i> (ATCC 12481)	<i>Pseudomonas aeruginosa</i> (ATCC 15442)
<i>Burkholderia cepacia</i> (ATCC 25416)	<i>Salmonella enterica</i> (ATCC 10708)
<i>Campylobacter jejuni</i> (ATCC 29428)	<i>Salmonella typhi</i> (ATCC 6539)
<i>Corynebacterium ammoniagenes</i> (ATCC 6871)	<i>Shigella sonnei</i> (ATCC 9290)
<i>Escherichia coli</i> 0157:H7 (ATCC 35150)	<i>Staphylococcus aureus</i> (ATCC 6538)
<i>Enterococcus faecium</i> (Vancomycin Resistant) (VRE)	<i>Staphylococcus aureus</i> (Methicillin resistant) (ATCC 33591)
<i>Klebsiella pneumoniae</i> (ATCC 13883)	<i>Yersinia enterocolitica</i> (ATCC 23715)
<i>Listeria monocytogenes</i> (ATCC 984)	
<i>Proteus mirabilis</i>	

General Disinfection Performance Non-Medical (schools, restaurants, food services, beverage and food processing plants): This product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum against:

<i>Campylobacter jejuni</i> (ATCC 29428)	<i>Staphylococcus aureus</i> (ATCC 6538)
<i>Escherichia coli</i> 0157:H7 (ATCC 31250)	<i>Staphylococcus aureus</i> (Methicillin resistant) (ATCC 33591)
<i>Listeria monocytogenes</i> (ATCC 984)	<i>Yersinia enterocolitica</i> (ATCC 23715)
<i>Proteus mirabilis</i>	
<i>Salmonella enterica</i> (ATCC 10708)	

Virucidal Performance: This product was evaluated in the presence of 5% serum and found to be effective against the following viruses on hard, non-porous environmental surfaces:

Hepatitis B Virus	Human Coronavirus (ATCC VR-740)
-------------------	---------------------------------

Hepatitis C Virus
 Herpes Simplex Type1 (ATCC VR-260)
 Herpes Simplex Type 2 (ATCC VR-734)
 HIV-1

Infectious Bronchitis Virus
 Influenza A2/Japan Virus (ATCC VR-100)
 Vaccinia virus

Animal Premise Virucidal Performance: This product was evaluated in the presence of 5% serum and found to be effective against the following viruses on hard, non-porous environmental surfaces:

Avian influenza A H9N2/Turkey/Wisconsin virus (ATCC VR-798)	Newcastle disease virus
Avian Reovirus (ATCC VR-2449)	Porcine Respiratory & Reproductive Virus (PRRSV)
Bovine Viral Diarrhea Virus (ATCC VR-534)	Porcine Rotavirus (ATCC VR-893)
Canine Coronavirus (ATCC VR-809)	Pseudorabies virus (ATCC VR-135)
Canine Distemper Virus	Transmissible Gastroenteritis (TGE) Virus
Equine Arteritis virus (ATCC VR-796)	
Infectious Bovine Rhinotracheitis virus (IBR)	
Infectious Laryngotracheitis Virus	

Non-Food Contact Sanitizing Performance: This product is an effective one-step sanitizer against the following:

<i>Staphylococcus aureus</i> (ATCC 6538)	<i>Enterococcus aerogenes</i> (ATCC 13048)
--	--

Fungicidal Activity: This product is effective against *Trichophyton mentagrophyti* (athlete's foot fungus) (a cause of Ringworm) on inanimate surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities.

MARKETING CLAIMS

Maquat® 10-PD for use in:

- Hospitals, nursing homes, medical and dental offices and clinics, physician offices, operating rooms, isolation wards, and medical research facilities.
- Patient care rooms & facilities, recovery rooms, anesthesia, Emergency Rooms, X-ray cat labs, newborn nurseries, orthopedics, whirlpools, footbath surfaces, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms.
- EMS & fire facilities, emergency vehicles, ambulance(s), ambulance equipment/surfaces, police cars.
- Day care centers and nurseries, sick rooms.
- Acute care institutions, alternate care institutions, home healthcare institutions.
- Life care retirement communities.
- Restaurants, restaurants and bars, bars, cafeterias, institutional kitchens, fast food operations and food storage areas.
- Supermarkets, convenience stores, retail and wholesale establishments, department stores, shopping malls, gift shops, video stores, bookstores, dressing rooms and laundries, photo copy centers, bicycle shops, auto repair centers.
- Computer manufacturing sites, toy factories.
- Food establishments, coffee shops, donut shops, bagel stores, pizza parlors, liquor stores.
- Crime scenes and funeral homes, mortuaries, burial vaults, mausoleums, autopsy rooms.

- Police stations, courthouses, correctional facilities, jails, prisons, municipal government buildings, penitentiaries, correctional institutions, bus stations, train stations.
- Institutional facilities, laboratories, factories, business and office buildings, restrooms, hotels and motels, and transportation terminals.
- Public restrooms, public facilities, waysides, travel rest areas, shower rooms, shower stalls, bathrooms.
- Hotel, motels, dormitories.
- Kitchens, bathrooms and other household areas.
- Homes (households).
- Institutions, schools and colleges, churches, classrooms, community colleges, universities, athletic facilities and locker rooms, exercise rooms, exercise facilities, gyms, gymnasiums.
- Cosmetic manufacturing facilities, medical device manufacturing facilities, biotechnology firms, pharmaceutical manufacturing facilities.
- Heath clubs, spas, tanning spas/beds, footbath surfaces massage/facial salons, hair/nail/pedicure salons, barber/beauty shops, salons, tattoo parlors.
- Museums, art galleries, post offices, performance/theater centers, banks, libraries, movie houses, bowling alleys.
- Recycling centers.
- Humidifier water tanks. *(Not for use in California)*
- Campgrounds, playgrounds, recreational facilities, picnic facilities, sports arenas, sports complexes.
- Dairy farms, hog farms, equine farms, poultry and turkey farms and egg processing plants, meat/poultry processing plants, meat/poultry producing establishments, mushroom farms, rendering plants.
- Tobacco plant premise.
- Veterinary clinics, animal life science laboratories, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops and other animal care facilities.
- Household and automotive garages, boats, ships, barges, campers, trailers, mobile homes, cars, trucks, buses, trains, taxis and airplanes.
- Cruise lines, airline terminals, airports, shipping terminals, public transportation.
- Commercial florist and flower shops.
- Basements, cellars, bedrooms, attics, living rooms and porches.

This product may be used on washable hard non-porous surfaces such as:

- Counters (countertops), stoves (stovetops), sinks (bathroom, kitchen), tub surfaces, exterior surfaces of appliances, refrigerators, and ice machines.
- Glass, metal, stainless steel, glazed porcelain, glazed ceramic, granite, marble, plastic (such as polystyrene or polypropylene), sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome and vinyl.
- Enameled surfaces, painted woodwork (finished), Formica®, vinyl and plastic upholstery.

- Foundations, steps, plumbing fixtures, finished baseboards and windowsills.
- Tables, chairs, desks, bed frames, lifts, washable walls, cabinets, doorknobs and garbage cans, cuspidors and spittoons.
- Exhaust fans, refrigerated storage and display equipment, coils and drain pans of air conditioning and refrigeration equipment and heat pumps.
- Large inflatable non-porous plastic and rubber structures: animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment.
- Hard, non-porous surfaces of picnic tables and outdoor furniture or Non-wooden Picnic tables and outdoor furniture except cushions and wood frames. (Note: Only one statement will be used on label.)
- Telephones and telephone booths.
- Highchairs, baby cribs, diaper changing stations, infant bassinets/cribs/warmers/incubators/care equipment, folding tables.
- Bed railings, bedpans, cervical collars, CPR training mannequins, curing lights, neck braces, oxygen hoods, slit lamps, spine backboards, stretchers and unit stools.
- External Lenses Vision correction including eyeglasses (not for use on contact lenses), protective eyewear, goggles, light lens covers, optical instruments/implements.
- Drinking fountains.
- Shower stalls, shower doors and curtains, bathtubs and glazed tiles, chrome plated intakes, toilets, toilet bowls, toilet bowl surfaces, urinals, empty diaper pails, portable and chemical toilets and latrine buckets, glazed porcelain tile and restroom fixtures.
- Ultrasonic baths, whirlpools, whirlpool bathtubs.
- Kennels, kennel runs, cages, kennel/cage floors, conductive flooring.
- Wrestling and gymnastic mats, athletic training tables, physical therapy tables.

This product contains no phosphorous.

This product is a phosphate-free formulation designed to provide effective cleaning, deodorizing and disinfection in areas where housekeeping is of prime importance in controlling the hazard of cross contamination on treated surfaces.

This product maximizes (improves) labor results by effectively controlling odors.

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

This product can be used to disinfect, clean and deodorize terrarium and small animal cages, hot rock, substrate and cage furniture (plastic terrarium ornaments, driftwood, heat caves and water dishes). *(Not for use in California.)*

This product is for use in kitchens, bathrooms, and other household areas.

This product is a multi-surface cleaner, deodorizer and disinfectant. Use on windows, mirrors, and other non-food contact glass surfaces.

When used as directed, this product will deodorize surfaces in rest room and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

516

Cross contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but also in schools, institutions and industry.

This product delivers non-acid disinfection performance in an economical concentrate.

This product is a concentrated Hospital Use disinfectant that is effective against a broad spectrum of bacteria, is virucidal*, (and) fungicidal, and eliminates odor causing bacteria when used as directed.

This product is an economical concentrate that can be used with a mop and bucket, trigger sprayers, sponge or by soaking.

This product will not leave a grit or soap scum.

Will not cause swelling of transducer membrane or harm compressor plates.

This product is a versatile Disinfectant & Sanitizer for Veterinarian, Veterinary Practice, Animal Care, Animal Laboratory, and Agricultural and Farm Premise applications.

This product is a complete, chemically balanced disinfectant/sanitizer that provides clear use solutions even in the presence of hard water.

This product inhibits bacterial growth on moist surfaces and deodorizes by killing microorganisms that cause offensive odors. *(Not for use in California.)*

This product is for use as a disinfectant on hard, non-porous surfaces (at 625 ppm active).

This product is for use in household and commercial humidifiers. Use of this product will control unpleasant (malodors) odors. *(Not for use in California.)*

This product is a versatile broad-spectrum disinfectant formulated for use in Ultrasonic Baths (Ultrasonic cleaning units).

This product is a versatile cleaner and broad-spectrum disinfectant formulated for use on bath and therapy equipment (Whirlpools).

Use this product to disinfect non-porous salon/barber tools and instruments: (such as) combs, brushes, scissors, clippers, trimmers, razors, blades, tweezers, and manicure instruments and footbath surfaces.

Use this product to sanitize and disinfect non-porous salon/barber tools and instruments: (such as) combs, brushes, scissors, clippers, trimmers, razors, blades, tweezers and manicure instruments.

This product is a one-step disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, HCV & HBV) and inhibits the growth of mold and mildew and their odors when used as directed.

This product neutralizes musty odors and tough odors from smoke, pet accidents, and spills on contact.

This product is specially formulated to effectively eliminate offensive odors caused by mold and mildew.

This product controls (reduces) (eliminates) (neutralizes) (destroys) odors to make your home (kitchen) (bathroom) sanitary.

This product is effective against household germs and odors caused by animal waste, septic tank or sewage backup, smoke and bathroom and kitchen odors.

Use this product on the multi-touch surfaces responsible for cross-contamination.

This product is effective at controlling mold and mildew on shower curtains.

This product is for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

This product is a no-rinse disinfectant cleaner that disinfects, cleans and deodorizes in one labor saving step.

This product may be used in work areas such as tool rooms and garages for odor control and light duty cleaning.

This product provides long lasting freshness against tough (pet) odors, such as odors from litter boxes and pet accidents.

This product provides effective cleaning strength that will not dull most metal-interlock floor finishes, and does not require a rinse prior to recoat.

This product cleans, shines, deodorizes and disinfects all hard non-porous surfaces listed on the label. It inhibits the growth of mold and mildew, leaving bathrooms and kitchens clean and fresh smelling.

This product is an effective antimicrobial cleaner, designed for use by wholesale and retail florists, shippers and green houses.

When used as directed, this product will disinfect hard, non-porous surfaces, such as flower buckets, floors and walls of coolers, design and packing benches, and countertops.

This product may be relied on to deodorize coolers, buckets, garbage pails and other areas where obnoxious odors may develop.

Use this product to clean, disinfect and deodorize flower buckets, walls, floors of coolers, shippers, greenhouse packing areas, garbage pails and other areas where obnoxious odors may develop.

This product may be used as a general purpose antimicrobial detergent in florist shops, wholesale florist, shippers, greenhouse packing areas and other commercial floriculture places for efficient cleaning and antimicrobial action against certain bacteria which cause: *(Not for use in California.)*

1. Plugging of stems with slime, which reduces uptake of water for various flowers including roses, chrysanthemums, gladioli and tulips.
2. Production of ethylene gas, which may injure blooms of the various sensitive flowers including carnations, snapdragons, some orchids, baby's breath, sweet peas, freesia and alstroemeria.

Floor cleaner.

One-step cleaner.

Cleans everyday kitchen messes.

Cleans kitchen surfaces and food preparation areas.

Cleans and shines.

Great for use (on) (in) the (kitchen), (bathroom), (floors) and other household areas. Removes (eliminates) odors. Deodorizes. Will not harm most surfaces.

Will not harm most surfaces. For a cleaner, fresher household. Removes stains. Removes dirt. Non-staining. Clear formula.

Eliminates odors caused by bacteria (and) (mildew) (and) (non-fresh foods). Kills odor causing bacteria in the kitchen (bathroom).

Cleans everyday kitchen messes like dirt, grease and food stains. Cuts through tough grease and grime.

216

Disinfects (and Sanitizes) kitchen surfaces (bathroom surfaces and floors). Sanitizes kitchen surfaces (bathroom surfaces and floors). Kills germs. Kills household bacteria.

Antibacterial.

Kills Athlete's Foot fungus on bathroom surfaces.

Hospital Use Disinfectant. Institutional Disinfectant (Sanitizer).

Kills common kitchen (bathroom) germs.

Use this product to clean, sanitize and disinfectant non-porous personal protective safety equipment, protective headgear, athletic helmets, wrestling/boxing headgear, athletic shoe soles, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors and ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

Use this product to clean, sanitize and disinfectant non-porous ambulance equipment and surfaces by rinsing all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at 120°F temperature will avoid overheating and distortion of the ambulance equipment and surfaces that would necessitate replacement.)

This product is for use in Poultry Premise Sanitation (Hatcheries) Egg Receiving Area, Egg Holding Area, Setter Room, Tray Dumping Area, Trays, Buggies, Racks, Egg Flats, Chick Holding Room, Hatchery Room, Chick Processing Area, Chick Loading Area, Poultry Buildings, Ceilings, Side walls and Floors, Drinkers, and all other hard non-porous Poultry House related equipment, and surfaces in the Hatchery environment.

This product is for use in Swine Premise Sanitation, Waterers and Feeders, Hauling Equipment, Dressing Plants, Loading Equipment, Farrowing Barns and Areas, Nursery, Blocks, Creep Area, Chutes.

This product meets AOAC Use – Dilution Test Standards for hospital disinfectants.

This product meets AOAC efficacy standards for hard, non -porous surface sanitizers.

This product is effective against the control of small flies on non-food contact surfaces such as floors, walls, drains, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing (Duck) Hepatitis B Virus.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Hepatitis C Virus. (Not for use in California.)

OR

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus).

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Human and Canine Coronavirus.

This product is an effective disinfectant/sanitizer in the presence of 5% serum contamination.

This product is an effective sanitizer in the presence of soils on non-food contact surfaces.

This product meets AOAC efficacy standards for hard surface non-food contact sanitizers.

For larger areas, such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfection.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

(Note for reviewer: For labels that list medical premises and metal and/or stainless steel surfaces, one of the following statements must be used.)

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

Or

This product is not for use on Medical device surfaces

Apply this product with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray (mist). (Use of "mist" is not allowed in California.)

NOTE: With spray applications, cover or remove all food products. Prepare a fresh solution at least daily or when use solution becomes visibly dirty. For heavily soiled areas, a preliminary cleaning is required.

DISINFECTION

General Disinfectant-Non-medical Dilution Chart

Ounces of Product	Amount of Water
0.15 ounces	1 quart
0.3 ounces	½ Gallon
0.6 ounces	1 gallon
1.5 ounces	2 ½ gallons
3.0 ounces	5 gallons
6.0 ounces	10 gallons

Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions Disinfection and Virucidal Dilution Chart

Ounces of Product	Amount of Water
0.2 ounces	1 quart
0.4 ounces	½ Gallon
0.8 ounces	1 gallon
2.0 ounces	2 ½ gallons
4.0 ounces	5 gallons
8.0 ounces	10 gallons

Dilution Chart:

For Hospital or Medical Environment claims	4oz/5 gal. water
For General or Broad Spectrum claims	3oz/5 gal. water
For Virucidal claims	4oz/5 gal. water
For Animal Virucidal claims	4oz/5 gal. water
For Non-Food Contact Sanitizing claims	1oz/4 gal. water
For Fungicide claims	4oz/5 gal. water

Disinfection Performance in Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions: Add 4 ounces of this product to 5 gallons of water (or equivalent use dilution) (625 ppm active quat) OR mix a 4-ounce packet with 5 gallons of water. Treated surfaces must remain wet for 10 minutes. At this use level, this product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum against:

<i>Botrytis cinerea</i> (ATCC 12481)	<i>Pseudomonas aeruginosa</i> (ATCC 15442)
<i>Burkholderia cepacia</i> (ATCC 25416)	<i>Salmonella enterica</i> (ATCC 10708)
<i>Campylobacter jejuni</i> (ATCC 29428)	<i>Salmonella typhi</i> (ATCC 6539)
<i>Corynebacterium ammoniagenes</i> (ATCC,6871)	<i>Shigella sonnei</i> (ATCC 9290)
<i>Escherichia coli</i> 0157:H7 (ATCC 35150)	<i>Staphylococcus aureus</i> (ATCC 6538)
<i>Enterococcus faecium</i> (Vancomycin Resistant) (VRE)	<i>Staphylococcus aureus</i> (Methicillin resistant) (ATCC 33591)
<i>Klebsiella pneumoniae</i> (ATCC 13883)	<i>Yersinia enterocolitica</i> (ATCC 23715)
<i>Listeria monocytogenes</i> (ATCC 984)	
<i>Proteus mirabilis</i>	

General Disinfectant Non-medical (schools, restaurants, food services, dairies, farm, beverage and food processing plants): Add 3 ounces of this product to 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Treated surfaces must remain wet for 10 minutes. At this use level, this product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum with a 10 minute contact time against:

<i>Campylobacter jejuni</i> (ATCC 29428)	<i>Staphylococcus aureus</i> (ATCC 6538)
<i>Escherichia coli</i> 0157:H7 (ATCC 31250)	<i>Staphylococcus aureus</i> (Methicillin resistant) (ATCC 33591)
<i>Listeria monocytogenes</i> (ATCC 984)	<i>Yersinia enterocolitica</i> (ATCC 23715)
<i>Proteus mirabilis</i>	
<i>Salmonella enterica</i> (ATCC 10708)	

Virucidal Performance: At 4 ounces per 5 gallons use level (or equivalent use dilution) (625 ppm active quat), this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard non-porous environmental surfaces.

Hepatitis B Virus	Human Coronavirus (ATCC VR-740)
Hepatitis C Virus	Infectious Bronchitis Virus
Herpes Simplex Type1 (ATCC VR-260)	Influenza A2/Japan Virus (ATCC VR-100)
Herpes Simplex Type 2 (ATCC VR-734)	Vaccinia virus

At 4.5 ounces per 5 gallons (703 ppm quat active) (or equivalent use dilution) use level, this product was evaluated in the presence of 5% serum with a 2-minute contact time and found to be effective against the following virus on hard, non-porous environmental surface:

HIV-1

Fungicidal Activity: At 4 ounces per 5 gallons use-level (or equivalent use dilution) (625 ppm active quat) this product is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) on inanimate surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities.

For Disinfecting Hard, Non-Porous Surfaces Against Avian Influenza: For hospital/medical/health facilities, the directions for use for hospital disinfection must be followed.

HOSPITAL/HEALTH CARE

Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions: For disinfecting floors, walls, countertops, bathing areas, lavatories, bed frames, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 4 ounces of this product to 5 gallons of water (625 ppm active quat) OR mix a 4-ounce packet with 5 gallons of water. Treated surfaces must remain wet for 10 minutes.

At 4 ounces per 5 gallons (or equivalent use dilution) (625 ppm active quat) in the presence of 200 ppm hard water (CaCO₃) and 100% serum this product was proven to be effective against (*Duck*) Hepatitis B Virus with a contact time of 10 minutes.

At 4 ounces per 5 gallons (or equivalent use dilution) (625 ppm active quat) was proven to be effective in the presence of 5% organic load against Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus) with a contact time of 10 minutes.

At 4.5 ounces per 5 gallons (or equivalent use dilution) in the presence of 5% serum for a two-minute contact time this product was found to be effective against HIV-1 (AID Virus).

***KILLS HIV, HCV AND HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type I (HIV-1) (associated with AIDS), Hepatitis C virus (HCV) and Hepatitis B virus (HBV).

"SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS."

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, or eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

Disposal of Infectious Materials: Blood and other body fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

Contact Time: For HIV, leave surfaces wet for two minutes with 4.5 ounces per 5 gallons (or equivalent use dilution) (703 ppm active quat) use solution. This contact time will not control other common types of viruses and bacteria. For HBV and HCV, leave surfaces wet for 10 minutes with 4 ounces per 5 gallons (or equivalent use dilution) (625 ppm active quat).

Efficacy tests have demonstrated that this product is an effective bactericide, virucide and fungicide in the presence of organic soil (5% blood serum).

Surgical Instrument Presoak : Add 4 ounces of this product to 5 gallons of water (or equivalent use dilution) (625 ppm active quat). Place pre-cleaned instruments in

2118

solution to presoak surgical instruments for a minimum of 10 minutes, then proceed with normal sterilization procedure.

NOTE: Plastics may remain immersed until ready to use. Instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Ultrasonic Bath Disinfectant Directions: Use this product to disinfect hard, (inanimate) non-porous, non-critical objects compatible with Ultrasonic cleaning units. Pour (a) fresh solution of 0.8 ounces per gallon of water (or equivalent use dilution) (625 ppm active quat) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visibly dirty or discolored.

For Disinfecting Hard, Non-porous Bath (Fiberglass) and Therapy Equipment: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the unit, drain the water and refill with fresh water to just cover the intake valve. Add 4 ounces of this product for each 5 gallons of water (0.80 ounces per one gallon) (or equivalent use dilution) (625 ppm active quat) in the unit at this point. Briefly start the pump to circulate the solution. Turn off pump. Wash down the unit sides, seat of the chair lift, and any/all related equipment with a clean swab, brush or sponge. Product to surface contact time must be at least 10 minutes for proper disinfection. After the unit has been thoroughly disinfected, drain solutions from the unit and rinse any/all cleaned surfaces with fresh water. The unit is ready for reuse.

Disinfection of Hard, Non-Porous Surfaces in Whirlpool Units: After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 4 ounces of this product for each 5 gallons of (0.80 ounces to one gallon) (or equivalent use dilution) (625 ppm active quat) water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

DEODORIZATION / MOLD & MILDEW

Fungicidal Activity: At 4 ounces per 5 gallons use-level (or equivalent use dilution) (625 ppm active quat) this product is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) on inanimate surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities.

Mold and Mildew Control: At 4 ounces per 5 gallons (or equivalent use dilution) (625 ppm active quat) this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat treatment every seven days or when new growth appears.

General Deodorization: To deodorize, add 3 ounces of this product to 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Excess material must be wiped up or allowed to air dry.

Cleaning/Deodorizing Directions: Add 2¼ to 9 ounces per gallon (or equivalent dilution) of water to clean and deodorize windows, mirrors and glass surfaces. Use a

coarse spray device. Spray 6-8 inches from surface. Rub with sponge or cloth. Do not breathe spray (mist). (Use of "mist" is not allowed in California.)

For Deodorizing Garbage Cans, Garbage Trucks, Industrial Waste Receptacles and Garbage Handling Equipment: It is especially important to pre-clean for the product to perform properly. Then, apply a wetting concentration of 3.25 ounces of this product per gallon of water (or equivalent use dilution) (2,500 ppm quat active).

For Deodorizing Septic Storage Tanks: When tanks are empty, pour 3 oz. per 5 gallons of water (469 ppm) (or equivalent use dilution) into septic storage tank on recreational vehicles, campers and boats to control the bacteria that can cause odors. As tank fills with sewage and the vehicle is moving, this disinfectant solution will mix and provide an adequate deodorizing solution for this use. This product is to be used on gray and black water tanks only. Check with local, state, or federal authorities before disposing of sewage in accordance to federal, state and local regulations for waste disposal.

Air Freshener/Automotive Uses: 3 ounces per 5 gallons of water will effectively neutralize damp musty odors caused by mildew in storage areas, basements, closets, bathrooms and A/C filters. Effective on smoking and cooking odors (garlic, fish, onions, etc.), automobile odors from tobacco, musty carpet smell, and beverage smells will be eliminated. Spray or apply onto seats, carpets, headliner and ashtray or into trunk and all vents to eliminate odors from tobacco, beverage spills and musty carpet. For sprayer applications, use a coarse spray device. Allow to air dry.

For Odors Caused by Dogs, Cats and Other Domestic Animals: Use on rugs, floors, walls, tile, cages, crates, litter boxes, mats, floor coverings, or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area. Then follow directions for "General Deodorization".

To Control the Growth of Mold and Mildew on Large, Inflatable, Non-Porous Plastic and Rubber Structures (animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 3 ounces per 5 gallons (469 ppm active quat) of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

To Control the Growth of Mold and Mildew on Non-Porous Athletic Equipment (wrestling and gymnastic mats, athletic training tables, physical therapy tables, athletic helmets, wrestling/boxing headgear, athletic shoe soles): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 3 ounces per 5 gallons (469 ppm active quat) of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

BARBER/SALON

Disinfection of Barber/Salon Tools Directions: Pre-cleaned barber/salon tools, such as combs, brushes, razors, clipper and trimmer blades, tweezers manicure/pedicure tools and scissors, can be disinfected by immersing in 0.6 ounces per gallon solution of this product (or equivalent use dilution) (469 ppm active quat). Completely immerse instruments and tools for at least 10 minutes. Rinse thoroughly and dry before use. Prepare fresh solution at least daily or more often if solution becomes cloudy or soiled.

Disinfection/Fungicide/Virucide for Barber/Salon Tools Directions: Immerse pre-cleaned barber/salon tools, such as combs, brushes, razors, clipper and trimmer blades, tweezers manicure/pedicure tools and scissors, in a 0.8 ounce per gallon

PH

solution of the product (or equivalent use dilution) (625 ppm active quat). Completely immerse instruments and tools for at least 10 minutes. Rinse thoroughly and dry before use. Prepare a fresh solution at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Disinfecting/Virucide Grooming Clippers : Turn the clipper off occasionally during use and spray between the teeth of blades a solution of 4 ounces per 5 gallons (or equivalent use dilution) (625 ppm active quat). Allow surface to remain wet for at least 10 minutes. Remove hair, dandruff and dust particles prior disinfecting the blades.

Cleaning Hair Clippers and Electric Shears : While clipper/shear is running, hold in the downward position and spray 3 ounces per 5 gallons of water solution (or equivalent use dilution) (469 ppm active quat) of this product directly onto the blades two or three times to thoroughly wet the blades. Do not spray on the clipper case or drip into clipper housing. Turn off clipper/shear. Allow surfaces to remain wet for at least 10 minutes. Then wipe dry with a clean soft cloth. Lubricate as per clipper/shear manufacturer's instructions.

Cleaning Barber/Salon Shears and Other Implements : Spray shear/implement until thoroughly wet. Wipe away visible debris using a soft bristle brush or cloth. Immerse pre-cleaned shear/implement into a container of 3 ounces per 5 gallons of water solution (or equivalent use dilution) (469 ppm active quat) of this product for at least 10 minutes. Remove shear/implement and wipe dry. No rinsing is necessary. Use fresh solution at least daily or when immersion tank (bowl) (sink) solution becomes diluted or visibly soiled.

Disinfection of Hard, Non-Porous Surfaces in Footbaths: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 3 ounces per 5 gallons of water (or equivalent use dilution) (469 ppm active quat) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 10 minutes. For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly disinfected, rinse all cleaned surfaces with fresh water. Do not use equipment until treatment has been absorbed, set or dried.

Disinfection/Fungicide/Virucide of Hard, Non-Porous Surfaces in Footbaths: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 0.8 ounce per gallon of water (or equivalent use dilution) (625 ppm active quat) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 10 minutes. For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly disinfected, rinse all cleaned surfaces with fresh water. Do not use equipment until treatment has been absorbed, set or dried.

RESTROOM/BATHROOM

NON-ACID TOILET BOWL (AND URINAL) DISINFECTANT DIRECTIONS

Remove gross filth prior to disinfection.

From Concentrate: Add ¾ oz. to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop, or sponge. Repeat and allow to stand for 10 minutes and flush.

From Use-solution: Empty toilet bowl or urinal and apply a 0.6 oz per gallon use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer; brush or swab thoroughly and allow to stand for 10 minutes and flush. For spray application use a coarse spray device.

For Heavy Duty Cleaning: Empty toilet bowl or urinal and apply 18 ounces per 5 gallons (or equivalent dilution) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly and allow to stand for 10 minutes and flush.

For Use To Clean and Disinfect Shower Rooms, Locker Rooms and Other Large Open Areas with Floor Drains:

1. Pre-clean heavily soiled areas.
2. Apply Use Solution of 4 ounces per 5 gallons of water (625 ppm quat active) to floors, walls and ceilings making sure not to over spray. To disinfect, all surfaces must remain wet for 10 minutes.
3. Scrub using deck brush or other coarse material as necessary.
4. Rinse surfaces thoroughly and let air dry.

RV Holding Tanks/Recreational Vehicles : For toilet waste and holding tanks, cover bottom of holding tank with water and 2.25 to 9 ounces per 5 gallons of water to deodorize. If odors return before time to empty, add 2.25 to 9 ounces to the tank. For kitchen waste, add 2.25 to 9 ounces to gray water tank as needed to control malodors created by dirty dishwasher.

FOOD PROCESSING PREMISES

To Disinfect Food-Processing and Tobacco Premises: Before using this product, food products and packaging materials must be removed from area or carefully protected. For use on floors, walls and storage areas, add 3 ounces of this product to 5 gallons of water (or equivalent use dilution) (469 ppm active quat). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications use a coarse spray device. Allow surfaces to remain wet for 10 minutes. Then remove excess liquid. For sprayer applications, use a coarse (mist) spray. After use, all surfaces in the area must be thoroughly rinsed with potable water. *(Use of "mist" is not allowed in California.)*

To Disinfect Food Service Establishment Food Contact Surfaces: (For countertops, exterior surfaces of appliances, (and) tables add 3 ounces of this product to 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Before using this product, food products and packaging materials must be removed from area or carefully protected. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications use a coarse spray device. Allow surfaces to remain wet for 10 minutes. Then remove excess liquid and rinse the surface with potable water. (After use, all surfaces in the area must be thoroughly rinsed with potable water.)

For use on non-food contact surfaces as a general disinfectant in the brewery industry: Use 3 ounces of this product per 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Follow the general disinfectant directions above.

Directions for Fogging: (For use in) dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room

or carefully protected. After cleaning, fog desired areas using one quart per 1000 cubic feet of room area with a product solution containing 1.5 ounces product to 1 gallon of water (or equivalent use dilution) (1,200 ppm). Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging. All food contact surfaces must be sanitized with an EPA approved food contact sanitizer solution of 200 ppm active quaternary (1 ounce per 4 gallons) prior to use. Allow food contact surfaces to drain thoroughly before operations are resumed.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

Disinfecting Potato Storage Area and Equipment: Remove all potatoes prior to disinfection of potato storage area and equipment. Pre-clean hard surfaces by removing heavy soils or gross filth. Follow general disinfecting directions as outlined in that section. All treated surfaces must be thoroughly rinsed with potable water prior to reuse.

ANIMAL PREMISES

Animal Premise Virucidal Performance: At 4 ounces per 5 gallons use level (or equivalent use dilution) (625 ppm active quat), this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard, non-porous environmental surfaces:

- | | |
|---|--|
| Avian influenza A H9N2/Turkey/Wisconsin virus (ATCC VR-798) | Newcastle disease virus |
| Avian Reovirus (ATCC VR-2449) | Porcine Respiratory & Reproductive Virus (PRRSV) |
| Bovine Viral Diarrhea Virus (ATCC VR-534) | Porcine Rotavirus (ATCC VR-893) |
| Canine Coronavirus (ATCC VR-809) | Pseudorabies virus (ATCC VR-135) |
| Canine Distemper Virus | Transmissible Gastroenteritis (TGE) Virus |
| Equine Arteritis virus (ATCC VR-796) | |
| Infectious Bovine Rhinotracheitis virus (IBR) | |
| Infectious Laryngotracheitis Virus | |

The first step in any on-going sanitation program must be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated thoroughly clean all surfaces with soap or detergent and rinse with water.

Disinfection of Poultry/Turkey Equipment, Swine Quarters, Equine Quarters, Animal Quarters and Kennels: Prior to disinfection, remove all poultry, other animals and their feed from premises, vehicles (trucks and cars), and enclosures (coops and crates, kennels, stables). Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by poultry or other animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 3 ounces of this product per 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

Disinfection/Fungicide/Virucide of Poultry/Turkey Equipment, Swine Quarters, Equine Quarters, Animal Quarters and Kennels Directions: Remove all animals and feeds from premises, vehicles (trucks and cars) and enclosures (coops and crates, kennels, stables). Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 4 ounces of this product per 5 gallons of water (or equivalent use dilution) (625 ppm active quat). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After application, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent, then rinsed with potable water before reuse.

HATCHERIES: For general disinfection, use 3 ounces of this product per 5 gallons of water (or equivalent use dilution). For disinfection/fungicide/virucide, use 4 ounces of this product per 5 gallons of water (or equivalent use dilution). Apply to hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces wet (exposed to solution) for 10 minutes or more. Rinse surfaces with potable water. Allow to air dry.

VEHICLES: Clean all vehicles including mats, crates, cabs, and wheels with high-pressure water and this product. For sprayer applications, use a coarse (mist) spray. Use 3 ounces of this product per 5 gallons of water (or equivalent use dilution) for general disinfection. Use 4 ounces of this product per 5 gallons of water (or equivalent use dilution) for disinfection/fungicide/virucide. Leave all treated surfaces exposed to solution for 10 minutes or more and allow to air dry. (Use of "mist" is not allowed California.)

RENDERING PLANT/DRESSING PLANT DISINFECTANT DIRECTIONS (DRESSING PLANT AND RENDERING PLANT DISINFECTANT DIRECTIONS): Cover or remove all food and packaging material before disinfection. Remove gross soils. Disinfect walls and floors in poultry and animal dressing plants with a solution of 3 ounces of this product per 5 gallons of water (or equivalent use dilution) (469 ppm active quat). Surfaces must remain wet for 10 minutes then thoroughly rinsed with potable water before operations are resumed. Disinfect all rooms, exterior walls and loading platforms of dressing plants. Rinse all surfaces which contact food with potable water before reuse.

RENDERING PLANT/DRESSING PLANT DISINFECTANT/FUNGICIDE/VIRUCIDE DIRECTIONS (DRESSING PLANT AND RENDERING PLANT DISINFECTANT - FUNGICIDE - VIRUCIDE DIRECTIONS): Cover or remove all food packaging material before disinfection. Remove gross soils. Apply to walls and floors in poultry and animal

11/15

dressing plants with a solution of 4 ounces of this product per 5 gallons of water (or equivalent use dilution) (625 ppm active quat). Surfaces must remain wet for 10 minutes then thoroughly rinsed with potable water before operations are resumed. Disinfect all rooms, exterior walls and loading platforms of dressing plants. Rinse all surfaces which contact food with potable water before reuse.

VEAL, CALVING, HOG, CATTLE AND HORSE OPERATIONS : Between depopulations of facilities, cleaning and disinfection of pens, hutches, aisles and other environmental surfaces may be required. Empty all troughs, racks and other feeding and watering appliances. Flush soils from these areas. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with recommended disinfecting or Virucidal solution and allow to remain wet for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

FARM PREMISE USE : Do not use in milking stalls, milking parlors or milk houses. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, stables, chutes and other facilities and fixtures occupied or transversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with recommended disinfecting or virucidal solution and allow to remain wet for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

(OR)

FARM PREMISE DISINFECTION DIRECTIONS

For use in Equine, Dairy and Hog Farms:

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, (stalls,) stables, chutes and other facilities and fixtures occupied or transversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with recommended disinfecting or virucidal solution for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

For Use in Treatment of Animal Housing Facilities:

1. Remove all animals and feed from premises, vehicles and enclosures.

2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, stables, chutes and other facilities and fixtures occupied or transversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the recommended disinfecting and virucidal solution (4 ounces per 5 gallons of water or equivalent use dilution) (625 ppm active quat) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

Terrarium and Small Animal Cage and Cage Furniture Disinfection: Animals frequently defecate on hot rocks and other cage furniture items inside your terrarium. This can result in high bacteria and ammonia levels that can lead to possible infection/disease in your animals. When used regularly this product can eliminate these high bacteria/ammonia levels in your cage and on your cage furniture items.

1. Remove all animals.
2. Thoroughly clean all surfaces and objects (hot rocks, caves, cage furniture, feeding and watering dishes, and appliances) including the substrate in the terrarium or cage with soap or detergent and rinse with water.
3. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the recommended disinfecting and virucidal solution (4 ounces per 5 gallons of water or equivalent use dilution) (625 ppm active quat) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. Then wipe surfaces dry.
4. Saturate gravel as above and let stand for 10 minutes. Place in bucket of clean water and swirl for 15-30 seconds. Thoroughly air dry before returning to terrarium.
5. Do not return animals to the habitat until it is dry and ventilated.
6. Thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.
7. Clean terrarium at least once weekly or more as needed.

Note: Substrates for desert terrariums (i.e. gravel) must be completely dry before returning to terrarium to avoid high humidity levels. Always replace substrate if a foul odor persists.

Veterinary Clinics/Animal Life Science Laboratory/Animal Care Facilities/Animal Research Centers/Animal Quarantine Areas/Zoos/Pet Shop/Kennels/Animal Breeding Facilities/Breeding and Grooming Establishment/Tack Shops Disinfection Directions: For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates etc. Remove all litter, droppings and manure from floors, walls and surfaces of

12/16

facilities occupied or transversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 4 ounces per 5 gallons of water (or equivalent dilution) (625 ppm active quat) for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers used in removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

Special Instructions for Inactivating Avian influenza A : Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with the recommended disinfecting solution (4 oz per 5 gallons of water) (625 ppm active quat) (equivalent use dilution) for a period of 10 minutes. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

OTHER

For Use on Finished Floors: To limit gloss reduction, use 4 ounces of this product per 5 gallons of water (or equivalent use dilution). Apply with a damp mop or auto scrubber. Allow to air dry.

Cleansing of Body Surfaces and Body Orifices of Human Remains: To cleanse away skin secretions and accompanying malodor, to insure the removal of all soil and bloodstains, apply 3 ounces of this product to 5 gallon of water (or equivalent use dilution) (469 ppm active) to the surfaces and body openings, natural or artificial. Allow a 10 minute contact time for optimal results. Bathe the entire body using sponge or washcloth. A soft brush may be employed on surfaces other than the face. Allow a 10 minute contact time for optimal results. Prepare a fresh solution for application of each remains.

COMMERCIAL FLORIST USE DIRECTIONS (*Not for use in California*): To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare use solution by adding (mixing) 3 ounces per 5 gallons of water (or equivalent use dilution) (469 ppm active). For heavy-duty use, add (mix) 12 ounces per 5 gallons of water (or equivalent use dilution) (1,875 ppm active).

Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply Use solution to hard (inanimate) non-porous surfaces, thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge or sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist). (*Use of "mist" is not allowed in California.*)

Treated surfaces must remain wet for 10 minutes. Allow to air dry.

Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

FOR CONTROL OF SMALL FLIES ON NON-FOOD CONTACT SURFACES: For control of small flies: *Drosophila* spp. and the Phoridae family. To control flies on non-food contact surfaces such as floors, walls, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas. Remove food and food packaging prior to use. Cover exposed food-handling surfaces. After removing gross filth, apply a solution of 4 ounces of this product to 5 gallons of water (or equivalent use dilution) (625 ppm active) to surfaces and locations where flies may breed. Spray surfaces thoroughly or apply by pouring, mopping or sponging onto the surface. Repeat application 1-2 times per week or as needed. Do not contaminate food and food packaging.

FOR CONTROL OF SMALL FLIES IN DRAINS: For control of small flies: *Drosophila* spp. and the Phoridae family. Spray or pour solution of 4 ounces per 5 gallon of water (or equivalent use dilution) (625 ppm active) into drain during time of lowest level of drain use. Add 14 ounces of solution daily to each drain to maintain fly control. Apply around the edge of the drain and coat all sides of inside of drain.

Waterbed Conditioner: When used as a waterbed conditioner, controls the growth of odor-causing and slime-forming bacteria. Prevents bubbles, preserves plasticizers conditions vinyl interior, sequesters minerals.

If bed has not been treated properly, drain bed completely. Add 15 gallons of water, mix vigorously, drain bed again. Fill bed with water and follow dosage directions.

Dosage: To control growth of odor-causing and slime-forming bacteria, add 4 1/8 fluid ounces in a free flow waterbed of 90 to 180 gallon capacity. Add 8 1/4 fluid ounces in a waveless/fiber waterbed mattress of 90 to 180 gallon capacity. Repeat application every 4 to 6 months.

SANITIZER DIRECTIONS FOR NON-FOOD CONTACT SURFACES

Non-Food Contact Sanitizing Performance: Add 1 ounce of this product to 4 gallons of water (or equivalent use dilution) (200 ppm active) to sanitize hard, non-porous surfaces. At 200 ppm active, this product is an effective one-step sanitizer against the following:

Staphylococcus aureus (ATCC 6538)

Enterococcus aerogenes (ATCC 13048)

Preparation of sanitizer use solution: Add 1 ounce of this product to 4 gallons of water (or equivalent use dilution) (200 ppm active) to sanitize hard, non-porous surfaces. At 200 ppm active, this product is an effective one-step sanitizer against *Staphylococcus aureus* and *Enterococcus aerogenes*.

Apply sanitizer use solution to precleaned, hard non-porous surfaces with a cloth, mop, sponge, sprayer or by immersion.

For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface, (and) rub with brush, sponge or cloth. Do not breathe spray. (*Use of "mist" is not allowed in California.*)

Treated surfaces must remain wet for 3 minutes, then wipe with sponge, mop or cloth or allow to air dry.

FOOD PROCESSING

SHOE (BOOT) BATH SANITIZER DIRECTIONS: To prevent cross contamination from area to area in animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made sanitizing solution must be placed at all entrances to buildings, hatcheries and at all the entrances to the production and

packaging rooms. Scrape waterproof shoes and place in a ¼ ounce of this product per gallon of water (or equivalent use dilution) (200 ppm active) solution for 3 minutes prior to entering area. Change the sanitizing solution in the bath at least daily or when solution appears dirty.

SHOE FOAM DIRECTIONS: To prevent cross contamination from area to area in animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 to 1.5 ounces per gallon of water (or equivalent use dilution) (781 to 1172 ppm active) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacture of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 3 minutes prior to entering area. Foam area must be washed and replaced at least daily or when it appears dirty.

(For food processing or other facilities that have installed entryway sanitizing systems)

ENTRYWAY SANITIZING SYSTEMS: To prevent cross contamination from area to area, set the system to deliver (0.51 – 1.02 oz. per gallon of water) (1.5 – 3 oz. per 2 7/8 oz per gallon of water) (or equivalent use dilution) (400 – 800 ppm active) of sanitizing solution. The (spray) (foam) must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

Do not mix other foam additives to the sanitizing solution.

FOOD PROCESSING PLANTS USING FOGGING DEVICES

Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using 1 quart per 1000 cubic feet of room area with a solution containing 1½ ounces of product to 1 gallon of water (or equivalent use dilution) (1,200 ppm). Vacate the area of all personnel for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed prior to reuse with potable water.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

ANIMAL HOUSING

Dairy Cow Application Directions: The udders, flanks and teats of dairy cows can be washed with a solution of 1 ounce of this product in 4 gallons of warm water. Use a fresh towel for each cow. Avoid contamination of wash solution by dirt and soil. Do not dip used towel back into wash solution. When solution becomes visibly dirty, discard and provide fresh solution.

TO SANITIZE HOOF TRIMMING EQUIPMENT : Prior to application, pre-clean hoof trimming equipment and utensils before and after use on each animal with detergent and warm water or compatible cleaner to remove soil using a pre-scrape, pre-flush, or when necessary, pre-soak followed by a potable water rinse. To sanitize, prepare a ¼ oz of this product per 1 gallon of water solution. Apply to pre-cleaned hard surfaces using a cloth, mop, sponge or sprayer or by immersion to thoroughly wet surfaces. Allow surfaces to remain wet for at least 5 minutes followed by adequate draining or air drying. Prepare fresh solution at least daily or when solution becomes visibly dirty.

SANITIZING HATCHERY ROOMS USING FOGGING DEVICES

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Mix 96 ounces of this product to 320 ounces of water (or equivalent use dilution). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed ventilate buildings and other closed spaces. Do not allow people to contact or breathe this fog and do not enter until 2 hours after fogging is complete. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with soap or detergent and rinse with water.

SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES

Mix 10 ounces of this product to 125 ounces of water. Fog 3-8 ounces of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog. It is acceptable to fog setters and hatchers with a 1 ounce per gallon solution of this product (or equivalent use dilution) an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. When fogging is complete, ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

OTHER

LAUNDRY DEODORIZER (AGAINST ODOR CAUSING BACTERIA): Use 3.5 ounces per 100 pounds of dry weight fabric. Dilute the appropriate amount of this product in one to two gallons of water, then add to the wash wheel in the final rinse. Re-treat fabric after each washing. For activity against odor causing bacteria, conditions of high relative humidity or wet contamination are required. Laundered fabric may also be treated by soaking.

SALON/BARBER INSTRUMENTS AND TOOLS SANITIZER DIRECTIONS: Pre-clean soiled instruments and tools. Completely immerse water safe non-porous instrument and tools in a solution of ¼ ounce per gallon (or equivalent use dilution) (200 ppm active) of water for at least 3 minutes. Rinse thoroughly and dry before use. Fresh solution must be prepared at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Ultrasonic Bath Sanitizer Directions : Use this product to sanitize hard, (inanimate) non-porous, non-critical objects compatible with Ultrasonic cleaning units. Pour a fresh solution of ¼ ounce per gallon of water (or equivalent use dilution) (200 ppm active quat) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 3 minutes, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visible dirty or discolored.

Sanitization of Hard, Non-Porous Surfaces in Whirlpool Units: After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add ¼ ounce of this product for each gallon of (or equivalent use dilution) (200 ppm active quat) water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 3 minutes. After the unit has been thoroughly sanitized, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

Humidifier Directions (Not applicable in CA): Thoroughly clean water tank and filters. Add 0.1 ounces of this product per gallon of refill water. Not for use in heat or atomizing type humidifiers.

OR

Humidifier Bacteria/Algae Treatment (Not for use in California.): Formulated for use in portable humidifiers. Thoroughly clean water tank and filters before each heating season, or sooner if necessary. For every gallon of water in the humidifier tank add 0.1 ounces of this product. When you refill the tank, add 0.1 ounces of this product for every gallon of water added.

Not for use in heat vaporizing or atomizing type humidifiers.

WATER AND SMOKE DAMAGE RESTORATION (Not for use in California.)

Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use. This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim, frame lumber, tackless strip and paneling. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

Sewer backup & river flooding: During mitigation procedures, dilute 1-2 ounces of this product per gallon of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affective areas with a sprayer using a coarse spray tip before and after cleaning and extraction.

Carpets, carpet cushions and other porous materials such as sub floors, drywall, trim and frame lumber, tackless strip and paneling: For water damage from a clean water source, extract excess water. Test hidden area for colorfastness. Dilute 1-2 ounces of the product per gallon of water (781-1562 ppm active quat) allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a sprayer using a coarse spray tip, to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for 10 minutes. Follow with a thorough extraction. Dry rapidly and thoroughly.

Special Instructions for Cleaning Carpet Against Odor Causing Bacteria: This product may be used in industrial, institutional, commercial and residential areas such as homes, motels & hotel chains, nursing homes, schools and hospital(s). For use on wet, cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for color fastness.

For portable extraction units: Mix ½ ounce of this product per gallon of water.

For truck mounted extraction machines: Mix 12 ounces of this product per gallon of water and meter at 4 gallons per hour.

For rotary floor machines: Mix 1 ounce of this product per gallon of water and apply at the rate of 300-500 sq. ft. per gallon.

Do not mix this product with other cleaning products. Follow the cleaning procedures, specified by the manufacturer of the cleaning equipment. After using this product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the carpet manufacturer's directions.

CARPET DEODORIZATION/SANITIZER AGAINST ODOR-CAUSING BACTERIA, FOR HOME, INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE (Use as Sanitizer is not for use in California.)

This product cleans and deodorizes (sanitizes) the carpet by controlling/reducing the growth of odor-causing bacteria. It can be used in industrial, institutional, commercial and residential areas such as homes, motels, hotel chains, nursing homes and hospitals.

Vacuum carpet thoroughly prior to application. Mix ½ ounce of product per gallon of water. Follow the injection and/or extraction procedures as specified for any conventional steam cleaning equipment you are using. For rotary floor machines, mix 1 ounce per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon.

For use (Use this product) on washable synthetic fibers. Do not use on wool. Test color fastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to (the) cloth, a water-based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

Note: This product should not be mixed with other cleaning products.

Smoke Damage Restoration (Not for use in California.): Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use. This product is particularly suitable for use in smoke damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim and frame lumber, tackless strip and paneling. Follow directions as outlined in the Water Damage Restoration section. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

MUSHROOM FARM INDUSTRY USE DIRECTIONS

Site Preparation: The first step in any on-going sanitation program must be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

Disinfection: Use 3 ounces of this product per 5 gallons of water (0.60 ounces per gallon of water) (or equivalent dilution). Wet all surfaces thoroughly. Treated surfaces must be allowed to remain wet for 10 minutes. Let air-dry. For heavily soiled areas, pre-clean first. Prepare a fresh solution for each use.

For Heavy Duty Cleaning: When greater cleaning is desired, use 6 ounces of this product per 5 gallons of water (1.2 ounces of this product per gallon of water) (or equivalent dilution). Heavily soiled areas may require repeated cleaning before treatment.

DO NOT APPLY TO THE MUSHROOM CROP, COMPOST OR CASING. Rinse treated surfaces with potable water before they contact the crop, compost or casing.

15/16

DIRECTIONS FOR ALTERNATE CONTAINERS

BAG-IN BOX CONTAINERS

How to use this package: This package is designed to be used with dilution control systems only. Open package and connect to dispense according to directions on the box.

Trigger sprayers: Fill bottle from dispenser. Apply to surfaces according to directions above.

Mop Buckets: Fill bucket from dispenser. Set up "Wet Floor" signs. Mop floor surfaces as specified in directions above.

PACKETS (For pre-measured tear open packets)

(Simply) (Tear) open and pour contents into 4.5 gallons of water. Keep packets in box until ready to use.

OR

Pour contents of (3.5 oz) packet into 4.5 gallons of water. Keep packets in box until ready to use.

REFILLS

To Refill Concentrate From Large Containers Into Smaller Containers : This product may be used to fill and refill clean, properly labeled containers for dilution elsewhere within your facility. Make sure the small container has been cleaned, dried and properly labeled according to state and local regulations. Also make sure other items (funnels or hand pumps) are properly cleaned and dried. To refill, simply pour (or pump product) from the larger container directly into the smaller one being careful not to spill any product. Keep both containers sealed when not in use.

FOUR OUNCE PACKET LABEL TO BE USED WITH MASTER CONTAINER LABEL FOR DISINFECTION

FOR DISINFECTION

MIX EACH PACKET WITH 5 GALLONS OF WATER

MAQUAT® 10-PD

E.P.A. Reg. No. 10324-99 E.P.A. Est. No. 10324-IL-1

NET CONTENTS: 4.0 FLUID OZ.

ACTIVE INGREDIENTS:

Alkyl (C ₁₄ 60%, C ₁₆ 30%, C ₁₂ 5%, C ₁₈ 5%)	
dimethyl benzyl ammonium chloride	5.0%
Alkyl (C ₁₂ 68%, C ₁₄ 32%)	
dimethyl ethylbenzyl ammonium chloride.....	5.0%

INERT INGREDIENTS: 90.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE OUTER CONTAINER FOR PRECAUTIONARY STATEMENTS AND USE DIRECTIONS

DISPOSAL: Do not reuse container. Wrap and put in trash collection.

MASON CHEMICAL COMPANY

"THE QUATERNARY SPECIALISTS"

721 W. Algonquin Road
Arlington Heights, IL 60005

847-290-1621

Toll Free:

1-800-362-1855



9/1/91