

NOV 22 2005

Ms. Elizabeth Tannehill
Mason Chemical Company
721 W. Algonquin Road
Arlington Heights, IL 60005

Subject: Maquat 128PD
EPA Registration No. 10324-105
Amendment Date: June 24, 2005
EPA Receipt Date: July 11, 2005

Dear Ms. Tannehill,

The following amendment submitted in connection with registration under FIFRA section 3(c)(7)(A) is acceptable subject to the conditions listed below.

- Addition of new use directions and public health organisms
- Addition of new alternate formulations

Conditions

Revise the label as follows:

- 1) On page 5, revise the nomenclature for the Avian Influenza virus to read "Avian influenza A H9N2/Turkey/Wisconsin Virus."
- 2) On page 5, revise the nomenclature for Influenza A2/Japan, Rabies, and Feline leukemia by stating "Influenza A2/Japan Virus, Rabies Virus, and Feline leukemia virus."
- 3) On page 5, you must place the organisms, Canine Parvovirus, Minute virus of Mice, and Porcine Parvovirus, under a separate heading that states:

At 4 ounces per one gallon use level or equivalent dilution, this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard, non-porous environmental surfaces:

- 4) On page 6, revise the last statement under "General Deodorization," to state "Excess liquid should be wiped up or allowed to air dry."
- 5) On page 6, under "Cleaning of Body Surfaces and Body Orifices of Human Remains," revise the use dilution statement to read "apply 1 ounces of this product to a gallon of water (705 ppm active) to the ~~entire body~~ **EXPOSURES**"

| | | | | | | | |
|---------|--|--|--|--|--|--|--|
| SYMBOL | | | | | | | |
| SURNAME | | | | | | | |
| DATE | | | | | | | |

6) On page 7, delete the two statements, "Before using this product, food product ... rinsed with potable water, that appear above the "To disinfect Food Processing and Tobacco Premises."

7) On page 7, revise the heading "Directions for Fogging" to include "Dairies, beverage, and food processing plants."

8) Revise the "Effectiveness against Canine Parvovirus (CPV), Minute virus of Mice (Parvovirus) and Porcine Parvovirus" directions such that the directions are appropriate for the treatment of sites which would house dogs, mice, and pigs. The rate associated with these organisms is not acceptable for facilities that house other types of animals; therefore, all references to poultry must be deleted. The directions must be revised to include "... Prior to use, remove *all animals*, and their feeds from premise vehicles and enclosures *such as crates*; "... occupied or traversed *by animals*...;" and "after application. *ventilate buildings*, and other closed spaces. *Do not house animals* or employ equipment..."

9) Revise the "Disinfection/Fungicidal/Virucide of Poultry/Turkey Equipment, Swine Quarters, Animal Quarters and Kennel Directions" by deleting "kennels." The user should refer to the section that precedes this one. In addition, add the following statement so that users treating swine quarters will have appropriate information, "For effectiveness against Porcine Parvovirus, use this product at 4 ounces per gallon of water. This statement must appear after the use dilution statement.

10) Revise the heading, "Farm Premise" to read "Farm Premise Disinfection Directions." Also add the statement, "For effectiveness against Porcine Parvovirus, use this product at 4 ounces per gallon of water," to the "Farm Premise Disinfection Directions and "For Treatment of Animal Housing Facilities" directions. This statement must appear after the use dilution statement.

11) On page 8, under "Terrarium and Small Animal Cage and Cage Furniture Directions." add the statement, "For effectiveness against Minute virus of Mice, use this product at 4 ounces per gallon of water."

12) On page 8. Under "Rendering Plants" delete the sites. "equipment and utensils" because they are food contact surfaces comparable to dishes and cookware which may not be disinfected.

13) On page 9. Under "Veterinary Clinics/Animal Life Science, Laboratory/Zoos/Pet Shops/Kennels/Breeding and Grooming Establishments/Tack Shops Disinfection Directions. revise the rate to "1.5 oz per gallon of water" because many of the animal pathogens mitigated at this rate by your product are housed at these facilities.

14) On page 9. Under "For Control of Drosophila ssp" directions, revise the second statement to read "To control flies on non-food contact..."

| CONCURRENCES | | | | | | | |
|--------------|--|--|--|--|--|--|--|
| SYMBOL | | | | | | | |
| SURNAME | | | | | | | |
| DATE | | | | | | | |

- 15) On page 9, delete the "Laundry Sanitizer (Against Odor Causing Bacteria)" section because you have not cited data to support this claim.
- 16) On page 9, Under "Shoe Bath Sanitizer Directions, Shoe Foam Directions, and Entryway Sanitizing Systems" revise the first statement to read "*To prevent cross contamination from area to area in animal areas, and the packing and storage...*"
- 17) On page 10, delete the "Dairy Cow Application Directions" because approval for this use must be obtained from the Food and Drug Administration (FDA) since this use is considered an animal drug. You must submit to the Agency a letter from FDA approving your product for this use.
- 18) On page 10, under "Sanitization of Hard, Non-Porous Surfaces in Whirlpool Units, revise the phrase, "...has been thoroughly disinfected..." to read "...has been thoroughly sanitized."
- 19) On page 10, revise the heading, "Humidifier Sanitizer Directions" to read "Humidifier Directions" because the term, sanitizer, is inappropriate for this use. The directions state that the product is treating the water not the tank; this product is registered for disinfecting/sanitizing hard, non-porous surfaces.

Acceptable Data

The Confidential Statements of Formulas for alternate formulations #1 and #2 are acceptable. They are in compliance with PR Notice 91-2 and in agreement with the label.

General Comments

Correction to Data Matrix

The correct MRID No. for the efficacy study submitted for Feline Rhino Virus is 4600217. The submitted matrix has been corrected, and you should revise your matrix on file as well as any future matrixes that may cite this study.

A stamped copy of the accepted labeling with conditions is enclosed. Submit one (1) copy of your final printed label before distributing or selling this product bearing the revised labeling.

Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.

If the above conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions.

| CONCURRENCES | | | | | | | |
|--------------|--|--|--|--|--|--|--|
| SYMBOL | | | | | | | |
| SURNAME | | | | | | | |
| DATE | | | | | | | |

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

4/16

Should you have any questions concerning this letter, please contact Jacqueline McFarlane at (703) 308-6416.

Sincerely,

for Tracy Lantz

Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510C)

Enclosure: Stamped Label

CONCURRENCES

| SYMBOL | | | | | | | |
|---------|--|--|--|--|--|--|--|
| | | | | | | | |
| SURNAME | | | | | | | |
| DATE | | | | | | | |

**PRECAUTIONARY STATEMENT.
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER. KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

(If container is 5 gallons or larger the following statement must appear on the label.)

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents.

(If the container is greater than one gallon use the following storage and disposal statements.)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL (Larger than 1 gal.): Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. In addition, plastic containers may be disposed of by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(If container is one gallon or less use the following storage and disposal statements.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Store in original container in areas inaccessible to small children. Do not store on side. Avoid creasing or impacting of side walls. Do not reuse empty container. Wrap and discard in trash (or recycle).

MAQUAT[®] 128 PD

Cleaner - Disinfectant - Deodorizer - Fungicide - Mildewstat - Virucide* - With Organic Soil Tolerance For Hospitals, Nursing Homes, Whirlpool, Home, Hotels & Motels, Waterbed Conditioner, Institutional, Industrial, School, Dairy, Equine, Poultry/Turkey Farm, Veterinary, Restaurant, Food Processing and Process Areas, Federally Inspected Meat and Poultry Plants, Bar and Institutional Kitchen and for Sanitizing Ice Machines.

Non-Acid Bathroom Cleaner

Athletic Surface Disinfectant.

Formulated for effective Poultry Sanitation.

Formulated for effective Swine Premise Sanitation, Fungicide, and Rodenticide Act as

Formulated for effective Mushroom Farm Sanitation, oriented, for the pesticide,

Small Fly Ovicidal Treatment.

ACTIVE INGREDIENTS:

Alkyl (C₁₄ 60%, C₁₆ 30%, C₁₂ 5%, C₁₈ 5%)

dimethyl benzyl ammonium chloride 4.5%

Alkyl (C₁₂ 68%, C₁₄ 32%)

dimethyl ethylbenzyl ammonium chloride 4.5%

INERT INGREDIENTS: 91.0%

TOTAL: 100.0%

NOV 22 2005
Under the Federal Fungicide, and Rodenticide Act as registered under EPA Reg. No. 10324-105

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See left panel for additional precautionary statements.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

MASON CHEMICAL COMPANY

"THE QUATERNARY SPECIALISTS"

721 W. Algonquin Road

Arlington Heights, IL 60005

847-290-1621

Toll Free:

1-800-362-1855

E.P.A. Reg. No. 10324-105

E.P.A. Est. No. 10324-IL-1



NET CONTENTS

BATCH NO

This product contains no phosphorous.

This product is a phosphate-free formulation designed to provide effective cleaning, deodorizing and disinfection in areas where housekeeping is of prime importance in controlling the hazard of cross contamination on treated surfaces.

This product maximizes (improves) labor results by effectively controlling odors.

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

This product can be used to disinfect, clean and deodorize terrarium and small animal cages, hot rock, substrate and cage furniture (plastic terrarium ornaments, driftwood, heat caves and water dishes.

This product is for use in kitchens, bathrooms, and other household areas

This product is a multi-surface cleaner, deodorizer and disinfectant. Use on windows, mirrors, and other non-food contact glass surfaces.

When used as directed, this product will deodorize surfaces in restroom and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

Cross contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on treated surfaces, not only in hospitals, but also in schools, institutions and industry.

This product delivers non-acid disinfection performance in an economical concentrate.

This product is a concentrated Hospital Use disinfectant that is effective against a broad spectrum of bacteria, is virucidal*, (and) fungicidal, and eliminates odor causing bacteria when used as directed.

This product is an economical concentrate that can be used with a mop and bucket, trigger sprayers, sponge or by soaking.

This product will not leave a grit or soap scum.

Will not cause swelling of transducer membrane or harm compressor plates.

This product is a versatile Disinfectant & Sanitizer for Veterinarian, Veterinary Practice, Animal Care, Animal Laboratory, and Agricultural and Farm Premise applications.

This product is a complete, chemically balanced disinfectant/sanitizer that provides clear use solutions.

This product inhibits bacterial growth on moist surfaces and deodorizes by killing microorganisms that cause offensive odors. *(Not applicable in CA)*

This product is recommended for use as a disinfectant on hard, non-porous surfaces (at 703 ppm active). A potable water rinse is required after application on food contact surfaces.

This product is recommended for use in household and commercial humidifiers. Use of this product will control unpleasant (malodors) odors. *(Not applicable in CA)*

This product is a versatile sanitizer and broad-spectrum disinfectant formulated for use in Ultrasonic Baths (Ultrasonic cleaning units).

This product is a versatile cleaner, sanitizer and broad-spectrum disinfectant formulated for use on bath and therapy equipment (Whirlpools).

Use this product to sanitize and disinfect non-porous salon/barber tools and instruments such as combs, brushes, scissors, clippers, trimmers, razors, blades, tweezers and manicure instruments, and footbath surfaces.

This product is a one-step disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, HCV & HBV) and inhibits the growth of mold and mildew and their odors when used as directed.

This product may be applied through automatic washing systems, immersion tanks, foaming apparatus, low-pressure sprayers, and fogging (wet misting) systems.

This product neutralizes musty odors and tough odors from smoke, pet accidents, and spills on contact.

This product is specially formulated to effectively eliminate offensive odors caused by mold and mildew.

This product controls (reduces) (eliminates) (neutralizes) (destroys) odors to make your home (kitchen) (bathroom) sanitary.

This product is effective against household germs and odors caused by animal waste, septic tank or sewage backup, smoke and bathroom and kitchen odors.

Use this product on the multi-touch surfaces responsible for cross-contamination.

This product is effective at controlling mold and mildew odor on shower curtains.

This product is recommended for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

This product is a no-rinse disinfectant cleaner that disinfects, cleans and deodorizes in one labor saving step.

This product may be used in work areas such as tool rooms and garages for odor control and light duty cleaning.

This product provides long lasting freshness against tough (pet) odors such as odors from litter boxes and pet accidents

This product cleans, shines, deodorizes and disinfects all hard, non-porous surfaces listed on the label. It inhibits the growth of mold and mildew, leaving bathrooms and kitchens clean and fresh smelling.

This product is an effective antimicrobial cleaner designed for use by wholesale and retail florists, shippers and green houses. *(Not applicable in CA)*

When used as directed, this product will disinfect hard, non-porous surfaces such as flower buckets, floors and walls of coolers, design and packing benches and countertops. *(Not applicable in CA)*

This product may be relied on to deodorize coolers, buckets, garbage pails and other areas where obnoxious odors may develop. *(Not applicable in CA)*

Use this product to clean, disinfect and deodorize flower buckets, walls, floors of coolers, shippers, greenhouse packing areas, garbage pails and other areas where obnoxious odors may develop. *(Not applicable in CA)*

This product may be used as a general purpose antimicrobial detergent in florist shops, wholesale florist, shippers, green house packing areas and other commercial floriculture places for efficient cleaning and antimicrobial action against certain bacteria which cause: *(Not applicable in CA)*

1. Plugging of stems with slime, which reduces uptake of water for various flowers including roses, chrysanthemums, gladioli and tulips.

6/10

2. Production of ethylene gas, which may injure bloom of the various sensitive flowers including carnations, snapdragons, some orchids, baby's breath, sweet peas, freesia and alstroemeria.

Floor cleaner.

One-step cleaner.

Cleans everyday kitchen messes.

Cleans kitchen surfaces and food preparation areas.

Cleans and shines.

Great for use (on) (in) the (kitchen), (bathroom), (floors) and other household areas. Removes (eliminates) odors. Deodorizes. Safe for most surfaces.

Safe for most surfaces. For a cleaner, fresher household. Removes stains. Removes dirt. Non-staining. Clear formula.

Eliminates odors caused by bacteria (and) (mildew) (and) (non-fresh foods). Kills odor causing bacteria in the kitchen (bathroom).

Cleans everyday kitchen messes like dirt, grease and food stains. Cuts through tough grease and grime.

Disinfects (and sanitizes) kitchen surfaces (bathroom surfaces and floors). Sanitizes kitchen surfaces (bathroom surfaces and floors). Kills germs. Kills household bacteria.

Antibacterial.

Kills Athlete's Foot fungus on bathroom surfaces.

Hospital use Disinfectant. Institutional Disinfectant (Sanitizer).

Kills common kitchen (bathroom) germs (and viruses*).

Maquat[®] 128 PD for use in:

- Hospitals, nursing homes, medical and dental offices and clinics, physician offices, operating rooms, isolation wards, and medical research facilities.
- Patient care rooms & facilities, recovery (rooms), anesthesia, Emergency Rooms, X-ray cat labs, newborn nurseries, orthopedics, whirlpool surfaces, footbath surfaces, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms.
- EMS & fire facilities, emergency vehicles, ambulance(s), ambulance equipment/surfaces, police cars.
- Day care centers and nurseries, sick rooms.
- Acute care institutions, alternate care institutions, home healthcare institutions.
- Life care retirement communities.
- Restaurants, restaurants and bars, bars, cafeterias, institutional kitchens, fast food operations and food storage areas.
- Supermarkets, convenience stores, retail and wholesale establishments, department stores, shopping malls, gift shops, video stores, bookstores, dressing rooms and laundries, photocopy centers, bicycle shops, auto repair centers.
- Computer manufacturing sites, toy factories.
- Food establishments, coffee shops, donut shops, bagel stores, pizza parlors, liquor stores.

- Crime scenes
- funeral homes, mortuaries, burial vaults, mausoleums, autopsy rooms.
- Police stations, courthouses, correctional facilities, jails, prisons, municipal government buildings, penitentiaries, correctional institutions, bus stations, train stations.
- Institutional facilities, laboratories, factories, business and office buildings, restrooms, hotels and motels and transportation terminals.
- Public restrooms, public facilities, waysides, travel rest areas, shower rooms, shower stalls, bathrooms.
- Hotel, motels, dormitories.
- Kitchens, bathrooms and other household areas.
- Homes (households).
- Institutions, schools and colleges, churches, classrooms, community colleges, universities, athletic facilities and locker rooms, exercise rooms, exercise facilities, gyms, gymnasiums.
- Cosmetic manufacturing facilities, medical device manufacturing facilities, biotechnology firms, pharmaceutical manufacturing facilities.
- Heath clubs, spas, tanning spas, tanning beds, footbath surfaces, massage/facial salons, hair/nail/pedicure salons, barber/beauty shops, salons, tattoo parlors.
- Museums, art galleries, post offices, performance/theater centers, banks, libraries, movie houses, bowling alleys.
- Recycling centers.
- Humidifier water tanks. *(Not for use in California)*
- Campgrounds, playgrounds, recreational facilities, picnic facilities, sports arenas, sports complexes.
- Food processing plants, USDA inspected food-processing facilities, dairy farms, hog farms, equine farms, poultry and turkey farms and egg processing plants, meat/poultry processing plants, meat/poultry producing establishments, mushroom farms, rendering plants.
- Processing facilities for Fish, Milk, Citrus, Ice Cream and Potatoes.
- Tobacco plant premise.
- Veterinary clinics, animal life science laboratories, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops and other animal care facilities.
- Animal holding areas, waiting rooms, examination rooms, x ray, operating rooms.
- Household and automotive garages, boats, ships, barges, campers, trailers, mobile homes, cars, trucks, buses, trains, taxis and airplanes.
- Cruise lines, airline terminals, airports, shipping terminals, public transportation.
- Commercial florist and flower shops.
- Basements, cellars, bedrooms, attics, living rooms, porches.

This product may be used on washable hard, non-porous surfaces such as:

- Countertops (counters), stovetops (stoves), sinks (bathroom kitchen), tub surfaces, and exterior surfaces of Slurppy[®] machines, appliances, refrigerators, and ice machines.
- Glass, metal, stainless steel, glazed porcelain, glazed ceramic, granite, marble, plastic (such as polystyrene or polypropylene), sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome and vinyl.
- Enameled surfaces, painted woodwork (finished), Formica[®], vinyl and plastic upholstery.
- Foundations, steps, plumbing fixtures, finished baseboards and window sills.
- Examination tables, X ray tables, washing areas, animal grooming areas.
- Tables, chairs, desks, bed frames, lifts, washable walls, cabinets, doorknobs and garbage cans, cuspidors and spittoons.
- Exhaust fans, refrigerated storage and display equipment, coils and drain pans of: air conditioning, refrigeration equipment and heat pumps.
- Large inflatable, non-porous plastic and rubber structures: animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment.
- Hard, non-porous surfaces of picnic tables and outdoor furniture furniture or Non-wooden Picnic tables and outdoor furniture except cushions and wood frames. (Note: Only one statement will be used on label.)
- Telephones and Telephone booths
- Highchairs
- Drinking fountains.
- Shower stalls, shower doors and curtains, bathtubs and glazed tiles, chrome plated intakes, toilets, toilet bowls, toilet bowl surfaces, urinals, empty diaper pails, portable and chemical toilets and latrine buckets, porcelain tile and restroom fixtures.
- Ultrasonic baths, whirlpools, whirlpool bathtubs.
- Kennels, kennel runs, cages, kennel/cage floors, conductive flooring.
- Wrestling and gymnastic mats, athletic training tables, physical therapy tables.

Use this product to clean, sanitize and disinfectant non-porous personal protective safety equipment, protective headgear, athletic helmets, wrestling/boxing headgear, athletic shoe soles, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors and ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at the recommended 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

Use this product to clean, sanitize and disinfectant non-porous ambulance equipment and surfaces by rinsing all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at the recommended 120°F temperature will avoid overheating and distortion of the ambulance equipment and surfaces that would necessitate replacement.)

This product is recommended for Poultry Premise Sanitation (Hatcheries)

| | | |
|--------------------|--------------------|-----------------------|
| Egg Receiving Area | Tray Dumping Area | Chick Processing Area |
| Egg Holding Area | Chick Holding Area | Chick Loading Area |
| Setter Room | Hatchery Room | Poultry Buildings |

This product is recommended for Swine Premise Sanitation:

| | | |
|---------------------------|-------------------|------------|
| Farrowing Barn. and Areas | Dressing Plants | Blocks |
| Waterers and Feeders | Loading Equipment | Creep Area |
| Hauling Equipment | Nursery | Chutes |

This product meets AOAC Use – Dilution Test Standards for hospital disinfectants.

This product meets AOAC efficacy standards for hard surface non-food contact sanitizers.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing (*Duck*) Hepatitis B Virus.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Hepatitis C Virus. (*Not for use in California*)

OR

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus).

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Human and Canine Coronavirus.

This product has passed the Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces utilizing Feline Calicivirus (surrogate for Norwalk virus).

This product is effective against the control of small flies on floors, walls, drains, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas.

For larger areas, such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfection.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

(Note for reviewer: For labels that list medical premises and metal and/or stainless steel surfaces, one of the following statements must be used:)

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

Or

This product is not for use on Medical device surfaces.

Apply this product with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray.

8/18

NOTE: With spray applications, cover or remove all food products. Prepare fresh solution at least daily or when use solution becomes visibly dirty. For heavily soiled areas, a preliminary cleaning is required.

DISINFECTION

Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions: For disinfecting floors, walls, countertops, bathing areas, lavatories, bed frames, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 1 ounce of this product to one gallon of water (703 ppm active quat) OR mix a 1 ounce packet with one gallon of water. Treated surfaces must remain wet for 10 minutes. At this use level, this product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum against:

| | |
|-------------------------------------|--|
| <i>Campylobacter jejuni</i> | <i>Salmonella choleraesuis</i> |
| <i>Corynebacterium ammoniagenes</i> | <i>Salmonella schottmuelleri</i> |
| <i>Enterobacter aerogenes</i> | <i>Salmonella typhi</i> |
| <i>Enterococcus faecalis</i> | <i>Shigella dysenteriae</i> |
| <i>Escherichia coli</i> | <i>Staphylococcus aureus</i> |
| <i>Klebsiella pneumoniae</i> | <i>Staphylococcus aureus (Methicillin resistant)</i> |
| <i>Listeria monocytogenes</i> | (10324-59) |
| <i>Pseudomonas aeruginosa</i> | <i>Streptococcus salivarius</i> |

Virucidal Performance: At 1 ounce per gallon use level (703 ppm active quat), this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard non-porous environmental surfaces:

| | |
|------------------------------------|--|
| Avian influenza A/Turkey/Wisconsin | Infectious Bronchitis Virus |
| Avian Reovirus | Norwalk/Norovirus (<i>new organism</i>) |
| Bovine Viral Diarrhea virus | Infectious Avian Laryngotracheitis |
| Canine Distemper | Infectious Bovine Rhinotracheitis virus (IBR) |
| Canine Coronavirus | Influenza A2/Japan |
| Human Coronavirus | Newcastle disease virus |
| Equine Arteritis Virus | Porcine Respiratory & Reproductive Virus (PRRSV) |
| Hepatitis B Virus (HBV) | Porcine Rotavirus |
| Hepatitis C Virus (HCV) | Pseudorabies virus |
| Herpes Simplex Type 1 | Transmissible Gastroenteritis (TGE) |
| HIV-1 (AIDS Virus) | Vaccinia virus |

At 1.5 ounces per gallon use level, this product is found to be effective against the following viruses on hard, non-porous environmental surfaces in the presence of 5% serum with a 10 minute contact time:

| | |
|---|-----------------------------------|
| Canine Adenovirus | Canine Parainfluenza Virus |
| Infectious Canine Hepatitis Virus | Feline Calicivirus |
| Feline Infectious Peritonitis | Feline leukemia |
| Feline Panleukopenia | Feline Picornavirus |
| Feline Rhinotracheitis | Murine Parainfluenza virus type 1 |
| Rabies | |
| Canine Parvovirus (CPV) at 4 oz. per gallon | |
| Minute virus of Mice (Parvovirus) at 4 oz. per gallon | |
| Porcine Parvovirus at 4 oz. per gallon | |

Fungicidal Performance:

| | |
|--------------------------|------------------------------------|
| <i>Aspergillus niger</i> | <i>Trichophyton mentagrophytes</i> |
|--------------------------|------------------------------------|

Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions
Disinfection Fungicidal and Virucidal Dilution Chart

| Ounces of Product | Amount of Water |
|-------------------|-----------------|
|-------------------|-----------------|

| | |
|--------------|---------------|
| 1 ounce | 1 quart |
| 1/2 ounce | 1/2 Gallon |
| 1.0 ounces | 1 gallon |
| 2 1/2 ounces | 2 1/2 gallons |
| 5.0 ounces | 5 gallons |
| 10.0 ounces | 10 gallons |

At 1 ounce per one gallon (or equivalent use dilution) (703 ppm active quat) in the presence of 5% organic load this product was proven to be effective against (Duck) Hepatitis B Virus with a contact time of 10 minutes.

At 1 ounce per one gallon (or equivalent use dilution) (703 ppm active quat) in the presence of 5% organic load this product was proven to be effective against Hepatitis C Virus (HCV) with a contact time of 10 minutes. (*Not for use in California*)

OR

At 1 ounce per one gallon (or equivalent use dilution) (703 ppm active quat) in the presence of 5% organic load this product was proven to be effective against (Duck) Hepatitis B Virus (HBV) and Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus [HCV]) with a contact time of 10 minutes.

At 1 ounce per one gallon in the presence of 5% blood serum for a two-minute contact time this product was found to be effective against HIV-1 (AIDS Virus).

***KILLS HIV, HCV and HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C virus (HCV).

"SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS."

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, or eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

Contact Time: Leave surfaces wet for two minutes for HIV and ten minutes for HBV and HCV with 1 ounce per gallon (or equivalent use dilution) (703 ppm active quat) use solution. This 2 minute contact time will not control other common types of viruses and bacteria.

Disinfection/Fungicidal/Virucidal Directions: Use at a rate of 1 ounce per gallon. Allow surfaces to remain wet for 10 minutes. Apply use solution to hard inanimate, non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist).

01/10

Add 1 ounce per gallon (8 milliliters per liter) of water to disinfect hard, non-porous surfaces. Treated surfaces must remain wet for 10 minutes. Prepare a fresh solution at least daily or when use dilution becomes diluted or soiled.

At 2 ounces per one gallon (or equivalent use dilution) (703 ppm active quat) in the presence of 5% organic load this product was proven to be effective against Feline Calicivirus (surrogate for Norwalk virus) with a contact time of 10 minutes.

For Use on Finished Floors: To limit gloss reduction, use 1 ounce of this product per one (1) gallon of water. Apply with a damp mop or auto scrubber. Allow to air dry.

General Deodorization: To deodorize, add 1 ounce of this product to one gallon of water (or equivalent use dilution) (703 ppm active quat). Allow to air dry. (Alternate statement: Excess material should be wiped up or allowed to air dry.)

For Deodorizing Garbage Cans, Garbage Trucks, Industrial Waste Receptacles and Garbage Handling Equipment: It is especially important to pre-clean for the product to perform properly. Then, apply a wetting concentration of 3.5 ounces of this product per gallon of water (or equivalent use dilution) (2,460 ppm quat active).

For Odors Caused by Dogs, Cats and Other Domestic Animals: Use on rugs, floors, walls, tile, cages, crates, litter boxes, mats, floor coverings, or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area. Then follow directions for "General Deodorization".

Fungicidal Activity: At 1 ounce per gallon use-level, (or equivalent use dilution) (703 ppm active quat) this product is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) and Aspergillus niger on inanimate surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities.

Mold and Mildew Control: At 1 ounce per gallon (or equivalent use dilution) (703 ppm active quat) this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat treatment every seven days or when new growth appears.

Cleansing of Body Surfaces and Body Orifices of Human Remains: To cleanse away skin secretions and accompanying malodor, to insure the removal of all soil and bloodstains, apply 1 ounce of this product to a gallon of water (703 ppm active) to the surfaces and body openings, natural or artificial. Allow a 10 minute contact time for optimal results. Bathe the entire body using sponge or washcloth. A soft brush may be employed on surfaces other than the face. Prepare a fresh solution for application of each remains.

To control the growth of mold and mildew on large inflatable non-porous plastic and rubber structures (animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 1 ounce per gallon (703 ppm active quat) of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

To control the growth of mold and mildew on non-porous athletic equipment (wrestling and gymnastic mats, athletic training tables, physical therapy tables, athletic helmets, wrestling/boxing headgear, athletic shoe soles): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 1 ounce per gallon (703 ppm active quat) of water (or equivalent dilution) for a period of

10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

Disinfection/Fungicide/Virucide for Barber/Salon Tools Directions: Immerse pre-cleaned barber/salon tools, such as combs, brushes, razors, clipper and trimmer blades, tweezers, manicure/pedicure tools and scissors, in a 1 ounce per gallon solution of the product (or equivalent use dilution) (703 ppm active quat). Completely immerse instruments and tools for at least 10 minutes. Rinse thoroughly and dry before use. Prepare a fresh solution at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Disinfection of Hard, Non-Porous Surfaces in Footbaths: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 1 ounce per gallon of water (or equivalent use dilution) (703 ppm active quat) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 10 minutes. For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly disinfected, rinse all cleaned surfaces with fresh water. Do not use equipment until treatment has been absorbed, set or dried.

NON-ACID TOILET BOWL (AND URINAL) DISINFECTION/CLEANER DIRECTIONS

Remove gross filth prior to disinfection.

From Concentrate: Add 1 ounce (8 milliliters per liter) to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop or sponge. Repeat and allow to stand for 10 minutes and flush.

From Use Solution: Empty toilet bowl or urinal and apply 1 ounce per gallon (8 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow to stand for 10 minutes and flush. For sprayer applications, use a coarse spray device.

For Heavy Duty Cleaning: Empty toilet bowl or urinal and apply 4 ounces per gallon (32 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow to stand for 10 minutes and flush.

Cleaning/Deodorizing Directions: Add ½ to 2 ounces per gallon (4-16 milliliters per liter) of water to clean and deodorize windows, mirrors and glass surfaces. Use a coarse spray device. Spray 6-8 inches from surface. Rub with sponge or cloth. Do not breathe spray (mist).

Ultrasonic Bath Disinfectant Directions: Use this product to disinfect hard (inanimate) non-porous, non-critical objects compatible with Ultrasonic cleaning units. Pour (a) fresh solution of 1 ounce per gallon of water (or equivalent use dilution) (703 ppm active quat) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visible dirty or discolored.

For Disinfecting Hard Non-porous Bath (Fiberglass) and Therapy Equipment: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate

surfaces after using the whirlpool unit, drain the water and refill with fresh water to just cover the intake valve. Add 1 ounce of this product for each gallon of water (703 ppm active quat) in the unit at this point. Briefly start the pump to circulate the solutions. Turn off pump. Wash down the unit sides, seat of the chair lift, and any/all related equipment with a clean swab, brush or sponge. Product to surface contact time must be at least 10 minutes for proper disinfection. After the unit has been thoroughly disinfected, drain solution from the unit and rinse any/all disinfected (cleaned) surfaces with fresh water. The unit is ready for reuse.

Disinfection of Hard Non-Porous Surfaces in Whirlpool Units: After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 1 ounce of this product for each gallon (or equivalent use dilution) (703 ppm active quat) of water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

Before using this product, food products and packaging materials must be removed from the room or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

To Disinfect Food-Processing and Tobacco Premises: For floors, walls and storage areas, add 1 ounce of this product to one gallon of water (or equivalent use dilution) (703 ppm active quat). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications, use a coarse spray device. Allow to remain wet for 10 minutes. Then remove excess liquid. Before using this product, food products and packaging materials must be removed from area or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

To Disinfect Food Service Establishment Food Contact Surfaces: For countertops, exterior surfaces of appliances, and tables, add 1 ounce of this product to one gallon of water (or equivalent use dilution) (703 ppm active quat). Before using this product, food products and packaging materials must be removed from area or carefully protected. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications, use a coarse spray device. Allow to remain wet for 10 minutes. Then remove excess liquid and rinse the surface with potable water. After use, all surfaces in the area must be thoroughly rinsed with potable water.

For Use on Non-Food Contact Surfaces as a General Disinfectant in the Brewery Industry: Use 1 ounce of this product per gallon of water (or equivalent use dilution) (703 ppm active quat). Follow the Disinfectant directions above.

Directions for Fogging: For use in dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using one quart per 1000 cubic feet of room area with a product solution containing 1¼ ounces product to 1 gallon of water (or equivalent use dilution) (1,230 ppm). Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging. All food contact surfaces must be sanitized with an EPA approved food contact sanitizer solution of 200 ppm active quaternary prior to use. Allow food contact surfaces to drain thoroughly before operations are resumed.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of

the actual fogging. The building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

POULTRY (AND SWINE) PREMISE SANITATION SITE PREPARATION

The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated thoroughly clean all surfaces with soap or detergent and rinse with water.

Effectiveness against Canine Parvovirus (CPV), Minute virus of Mice (Parvovirus) and Porcine Parvovirus: At 4 ounces per gallon of water or equivalent dilution this product is found to be effective against Canine Parvovirus (CPV), Minute virus of Mice (Parvovirus) and Porcine Parvovirus on hard, non-porous environmental surfaces in the presence of 5% serum with a 10 minute contact time. Prior to use, remove all poultry, other animals, and their feeds from premises vehicles and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by poultry and other animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 4 ounces of this product per gallon of water (or equivalent use dilution). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. After application, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

OR

For Canine Parvovirus Activity: This product is effective against Canine Parvovirus at 4 oz. per gallon. Follow Veterinary Clinics/Animal Life Science Laboratory/Zoos/Pet Shop/Kennels/Breeding and Grooming Establishment/Tack Shops Disinfection Directions.

Disinfection/Fungicide/Virucide of Poultry/Turkey Equipment, Swine Quarters, Animal Quarters and Kennels Directions: Prior to use, remove all poultry, other animals and their feeds from premises vehicles and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by poultry, other animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 1 ounce of this product per gallon of water (or equivalent use dilution) (703 ppm active quat). Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After application, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

HATCHERIES: Use 1 ounce of this product per gallon of water to treat the following hard non-porous surfaces: hatcher, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more. Rinse with potable water before reuse. Then allow to air dry.

VEHICLES: Clean all vehicles including mats, crates, cabs, and wheels with high-pressure water and this product. For sprayer applications, use a coarse (mist or) spray. Use 1 ounce per gallon to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more and allow to air dry.

DRESSING PLANT USE: Disinfect walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the recommended use solution of 1 ounce of this product per gallon of water. Scrub to loosen all soils. Allow to soak for 10 minutes and thoroughly rinse all wetted and cleaned surfaces with potable water.

OR

RENDERING PLANT/DRESSING PLANT DISINFECTANT/FUNGICIDE/VIRUCIDE DIRECTIONS (DRESSING PLANT AND RENDERING PLANT DISINFECTANT - FUNGICIDE - VIRUCIDE DIRECTIONS): Cover or remove all food packaging material before disinfection. Remove gross soils. Apply to walls and floors in poultry and animal dressing plants with a solution of 1 ounce of this product per 1 gallon of water (or equivalent use dilution) (703 ppm active quat). Equipment and utensils must remain wet for 10 minutes then thoroughly rinsed with potable water before operations are resumed. Disinfect all rooms, exterior walls and loading platforms of dressing plants. Thoroughly rinse all wetted and cleaned surfaces with potable water.

FARM PREMISE USE: Do not use in milking stalls, milking parlors or milk houses. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or transversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with recommended disinfecting or virucidal solution and allow to remain wet for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

(OR)

FARM PREMISE DISINFECTION DIRECTIONS

For use in Equine, Dairy and Hog Farms:

1. Remove all animals and feed from premises, vehicles and enclosures.

2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, chutes and other facilities and fixtures occupied or transversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the recommended disinfecting or virucidal solution for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

For Use For Treatment of Animal Housing Facilities:

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, chutes and other facilities and fixtures occupied or transversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the recommended disinfecting and virucidal solution (1 ounce per gallon of water or equivalent use dilution) (703 ppm active quat) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

Terrarium and Small Animal Cage and Cage Furniture Disinfection: Animals frequently defecate on hot rocks and other cage furniture items inside your terrarium. This can result in high bacteria and ammonia levels that can lead to possible infection/disease in your animals. When used regularly this product can eliminate these high bacteria/ammonia levels in your cage and on your cage furniture items.

1. Remove all animals.
2. Thoroughly clean all surfaces and objects (hot rocks, caves, cage furniture, feeding and watering dishes, and appliances) including the substrate in the terrarium or cage with soap or detergent and rinse with water.
3. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the recommended disinfecting and virucidal solution (1 ounce per gallon of water or equivalent use dilution) (703 ppm active quat) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. Then wipe surfaces dry.

12/10

4. Saturate gravel as above and let stand for 10 minutes. Place in bucket of clean water and swirl for 15-30 seconds. Thoroughly dry before returning to terrarium.
5. Do not return animals to the habitat until it is dry and ventilated.
6. Thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.
7. Clean terrarium at least once weekly or more as needed.

Note: Substrates for desert terrariums (i.e. gravel) must be completely dry before returning to terrarium to avoid high humidity levels. Always replace substrate if a foul odor persists.

CLEANING AND DISINFECTION: For all general cleaning and disinfection use 1 ounce of this product per gallon of water. Apply this product using a cloth, mop, or pressure sprayer so as to thoroughly wet surface to be disinfected. Allow to remain wet for 10 minutes and then let air dry. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution after each use. For sprayer applications, use a coarse spray device.

Disinfecting Potato Storage Area and Equipment: Remove all potatoes prior to disinfection of potato storage area and equipment. Pre-clean hard surfaces by removing heavy soils or gross filth. Follow general disinfecting directions as outlined in that section. All treated surfaces must be thoroughly rinsed with potable water prior to reuse.

Veterinary Clinics/Animal Life Science Laboratory/Zoos/Pet Shop/Kennels/Breeding and Grooming Establishment/Tack Shops Disinfection

Directions: For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or transversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 1 ounce per gallon of water (or equivalent dilution) (703 ppm active quat) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

COMMERCIAL FLORIST USE DIRECTIONS (Not applicable in CA): To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare use solution by adding (mixing) 1 ounce per gallon of water (1:128). For heavy-duty cleaning, add (mix) 4 ounces per gallon of water (1:32).

Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply use solution to hard (inanimate) non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge or sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse (mist) pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist).

Treated surfaces must remain wet for 10 minutes. Allow to air dry.

Prepare a fresh solution at least daily or sooner if use solution becomes visibly dirty.

FOR CONTROL OF SMALL FLIES ON NON-FOOD CONTACT SURFACES: For control of small flies: *Drosophila* spp. and the Phoridae family. To control flies on non-food contact surfaces such as floors, walls, countertops, metal surfaces, painted

surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas. Remove food and food packaging prior to use. Cover exposed food-handling surfaces. After removing gross filth, apply a solution of 1 ounce of this product per gallon of water (or equivalent use dilution) (703 ppm active quat) to surfaces and locations where flies may breed. Spray surfaces thoroughly or apply by pouring, mopping or sponging onto the surface. Repeat application 1-2 times per week or as needed. Do not contaminate food and food packaging.

FOR CONTROL OF SMALL FLIES IN DRAINS: For control of small flies: *Drosophila* spp. and the Phoridae family. Spray or pour solution of 1 ounce per gallon of water (or equivalent use dilution) (703 ppm active quat) into drain during time of lowest level of drain use. Add 14 ounces of solution daily to each drain to maintain fly control. Apply around the edge of the drain and coat all sides of inside of drain.

Waterbed Conditioner: When used as a waterbed conditioner, controls the growth of odor-causing and slime-forming bacteria. Prevents bubbles, preserves plasticizers, conditions vinyl interior, sequesters minerals.

If bed has not been treated properly, drain bed completely. Add 15 gallons of water, mix vigorously, drain bed again. Fill bed with water and follow dosage directions.

Dosage: To control growth of odor-causing and slime-forming bacteria, add 4.56 fluid ounces in a free flow waterbed of 90 to 180 gallon capacity. Add 9.125 fluid ounces in a waveless/fiber waterbed mattress of 90 to 180 gallon capacity. Repeat application every 4 to 6 months.


SANITIZER DIRECTIONS FOR NON-FOOD CONTACT SURFACES

Preparation of sanitizer use solution: Add 1 ounce of this product to 1 gallon of water (or equivalent use dilution) (703 ppm active) to sanitize hard, non-porous surfaces. At 703 ppm active, this product is an effective one-step sanitizer against *Staphylococcus aureus* and *Klebsiella pneumoniae*.

Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces with a cloth, mop, sponge, sprayer or by immersion.

For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray.

Treated surfaces must remain wet for 60 seconds, then wipe with sponge, mop or cloth or allow to air dry.



SHOE BATH SANITIZATION DIRECTIONS: To prevent cross contamination into treated animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made sanitizing solution should be placed at all entrances to buildings, hatcheries and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and place in 1 ounce of this product per gallon (8 milliliters per liter) of water solution for 60 seconds prior to entering area. Change the sanitizing solution in the bath at least daily or sooner if solution appears dirty.

13/10

SHOE FOAM DIRECTIONS: To prevent cross contamination in treated animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 to 1.5 ounces per gallon of water (or equivalent use dilution) (703 to 1051 ppm active) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacture of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 60 seconds prior to entering area. Foam area should be washed and replaced at least daily or when it appears dirty.

(For food processing or other facilities that have installed entryway sanitizing systems.)

ENTRYWAY SANITIZING SYSTEMS: To prevent cross contamination of harmful organisms from area to area, set the system to deliver (1oz. per gallon of water) (703 ppm active) of sanitizing solution. The (spray) (foam) should cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

Do not mix other foam additives to the sanitizing solution.

SALON/BARBER INSTRUMENTS AND TOOLS SANITIZER DIRECTIONS: Pre-clean soiled instruments and tools. Completely immerse water safe non-porous instruments and tools in a solution of 1 ounce per gallon (703 ppm active) (or equivalent use dilution) of water for at least 60 seconds. Rinse thoroughly and dry before use. Fresh solution should be prepared at least daily or more often if solution becomes cloudy or soiled.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

Ultrasonic Bath Sanitizer Directions: Use this product to disinfect hard, (inanimate) non-porous, non-critical objects compatible with Ultrasonic cleaning units. Pour a fresh solution of 1 ounce per gallon of water (or equivalent use dilution) (703 ppm active quat) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 60 seconds, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visible dirty or discolored.

Sanitization of Hard, Non-Porous Surfaces in Whirlpool Units: After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 1 ounce of this product for each gallon of (or equivalent use dilution) (703 ppm active quat) water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 60 seconds. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

Footbath Sanitizer Directions: To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 1 oz per gallon of water (or equivalent use dilution) (703 ppm active quat) to exposed surfaces with a cloth, mop,

sponge or spray. Brush or swab thoroughly and allow solution to stand for 60 seconds. For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly sanitized, rinse all cleaned surfaces with fresh water. Do not use equipment until treatment has been absorbed, set or dried.

Humidifier Sanitizer Directions *(Not applicable in CA):* Thoroughly clean water tank and filters. Add 1 ounce of this product per gallon of refill water. Not for use in heat or atomizing type humidifiers.

OR

Humidifier Bacteria/Algae Treatment *(Not applicable in CA):* Formulated for use in portable humidifiers. Thoroughly clean water tank and filters before each heating season, or sooner if necessary. For every 1 gallon of water in the humidifier tank add 1 ounce of this product. When you refill the tank, add ¼ ounce of this product for every 2 gallons of water in the tank.

Not for use in heat vaporizing or atomizing type humidifiers.

WATER AND SMOKE DAMAGE RESTORATION *(Not applicable in CA)*

Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use. This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim, and frame lumber, tackless strip and paneling. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

Sewer backup & river flooding: During mitigation procedures, dilute 1-2 ounces of this product per gallon of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affective areas with a sprayer using a coarse spray tip, before and after cleaning and extraction.

Carpets, carpet cushions and other porous materials such as sub floors, drywall, trim and frame lumber, tackless strip and paneling: For water damage from a clean water source, extract excess water. Test hidden area for colorfastness. Dilute 1-2 ounces of the product per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a sprayer using a coarse spray tip, to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for 10 minutes. Follow with a through extraction. Dry rapidly and thoroughly.

Special Instructions for Cleaning Carpet Against Odor Causing Bacteria: This product may be used in industrial and institutional areas such as homes, motels & hotel chains, nursing homes, schools and hospital(s). For use on wet, cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for color fastness.

For Portable Extraction Units: Mix ½ ounce of this product per gallon of water.

For Truck Mounted Extraction Machines: Mix 12 ounces of the product per gallon of water and meter at 4 gallons per hour.

For Rotary Floor Machines: Mix 1 ounce of this product per gallon of water and apply at the rate of 300-500 sq. ft. per gallon.

Do not mix this product with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using this product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the carpet manufacturer's directions.

CARPET SANITIZER AGAINST ODOR-CAUSING BACTERIA, FOR HOME, INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE (Not applicable in CA)

This product sanitizes the carpet by controlling/reducing the growth of odor-causing bacteria. It can be used in industrial and institutional areas such as homes, motels, hotel chains, nursing homes and hospitals.

Vacuum carpet thoroughly prior to application. Mix ½ ounce of product per gallon of water. Follow the Injection and/or Extraction procedures as specified for any conventional steam cleaning equipment you are using. For rotary floor machines, mix 1 ounce per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon.

For use (Use this product) on washable synthetic fibers. Do not use on wool. Test color fastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to (the) cloth, a water-based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

Note: This product should not be mixed with other cleaning products.

Smoke Damage Restoration (Not applicable in CA): Effective against odor causing bacteria and fungi for home, institutional, industrial and hospital use. This product is particularly suitable for use in smoke damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim and frame lumber, tackless strip and paneling. Follow directions as outlined in the Water Damage Restoration section. Using solutions recommended, saturate affected materials with enough product to remain wet for at least 10 minutes. Use proper ventilation.

SANITIZING HATCHERY ROOMS USING FOGGING DEVICES

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Mix 155 ounces this product to 2.5 gallons water (or equivalent use dilution). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed ventilate buildings and other closed spaces. Do not allow people to contact or breath this fog and do not enter until 2 hours after fogging is complete. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with soap or detergent and rinse with water.

SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES

Mix 16 ounces of this product to 128 ounces of water. Fog 3-8 ounces of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breath this. It is acceptable to fog setters and hatchers with 1 ounce per gallon solution of this product (or equivalent use dilution) on an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. When fogging is completed ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Do not allow people to contact or breath this fog and do not enter until 2 hours after fogging is complete. Thoroughly scrub all treated feed racks,

mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

FOOD PROCESSING PLANTS USING FOGGING DEVICES

For use in dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using 1 quart per 1000 cubic feet of room area with a solution containing 1¾ ounces of product to 1 gallon of water (1,230 ppm) (or equivalent use dilution). Vacate the area of all personnel during fogging and for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed prior to reuse with potable water.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.

MUSHROOM FARM INDUSTRY USE DIRECTIONS

Site Preparation: The first step in any on going sanitation program should be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

Disinfection: Use 1 ounce of this product per gallon of water (or equivalent dilution). Wet all surfaces thoroughly. Treated surfaces should be allowed to remain wet for 10 minutes. Let air-dry. For heavily soiled areas, preclean first. Prepare a fresh solution for each use.

For Heavy Duty Cleaning: When greater cleaning is desired, use 2 ounces of this product per gallon of water. (or equivalent dilution). Heavily soiled areas may require repeated cleaning before treatment.

DO NOT APPLY TO THE MUSHROOM CROP, COMPOST OR CASING. Rinse treated surfaces with potable water before they contact the crop, compost or casing.

(Alternative language for bag in the box dispensing equipment.)

DIRECTIONS FOR BAG-IN BOX CONTAINERS

How to use this package: This package is designed to be used with dilution control systems only. Open package and connect to dispense according to directions on the box.

Trigger sprayers: Fill bottle from dispenser. Apply to surfaces according to directions above.

Mop Buckets: Fill bucket from dispenser. Set up "Wet Floor" signs. Mop floor surfaces as specified in directions above.

5110

**ONE OUNCE PACKET LABEL TO BE USED WITH MASTER
CONTAINER LABEL FOR DISINFECTION**

FOR DISINFECTION

MIX EACH PACKET WITH 1 GALLON OF WATER

MAQUAT[®] 128 PD

E.P.A. Reg. No. 10324-105 E.P.A. Est. No. 10324-IL-1

NET CONTENTS: 1.0 FLUID OZ.

ACTIVE INGREDIENTS:

Alkyl (C₁₄ 60%, C₁₆ 30%, C₁₂ 5%, C₁₈ 5%)
dimethyl benzyl ammonium chloride.....4.5%

Alkyl (C₁₂ 68%, C₁₄ 32%)
dimethyl ethylbenzyl ammonium chloride4.5%

INERT INGREDIENTS:91.0%

TOTAL:100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

**SEE OUTER CONTAINER FOR PRECAUTIONARY STATEMENTS AND USE
DIRECTIONS.**

DISPOSAL: Do not reuse container. Wrap and put in trash collection.

**MASON CHEMICAL COMPANY
"THE QUATERNARY SPECIALISTS"**



721 W. Algonquin Road
Arlington Heights, IL 60005

847-290-1621

Toll Free:
1-800-362-1855

16/16