



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 4, 2021

Ryan J. Connair
REGISTRATION SPECIALIST
Mason, a Pilot Chemical Company
9075 Centre Pointe Dr., Suite 400
West Chester, OH 45069

Subject: Label Amendment – Updating Master Label
Product Name: Maguard 1522
EPA Registration Number: 10324-230
Received Date: 01/20/2021
Action Case Number: 00216634

Dear Ryan Connair:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. Pursuant to 40 CFR 156.10(a)(6) you must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. See FIFRA section 2(p)(2). If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process, FIFRA section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Assurance.

Page 2 of 2
EPA Reg. No. 10324-230
Action Case Number: 00216634

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Terria Northern via email at northern.terria@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Steven Snyderman". The signature is written in a cursive style with a horizontal line underneath the name.

Steven Snyderman, Product Manager 33
Regulatory Management Branch II
Antimicrobials Division (7510P)
Office of Pesticide Programs

Enclosure: Accepted label

MAGUARD® 1522

(Note to Reviewer: Marketing claims may be used on the front panel.)

{Peroxyacetic-Acid Based{:}}

Disinfectant • {Food Contact} Sanitizer • Bactericide • Virucide* • Fungicide • Tuberculocide
• Cleaner • Deodorizer

ACTIVE INGREDIENTS:

Hydrogen peroxide.....	22.0%
Peroxyacetic acid	15.0%

OTHER INGREDIENTS: 63.0%

TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
DANGER {PELIGRO}**

{See [{left} {back} {side} {right} {insert} {panel} {of label} {below}] for {additional} {precautionary statements}.

(Note to Reviewer: In accordance with 40 CFR 156.68(d), all first aid statements, as prescribed, will appear on the front panel of the product label.)

FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.


{For [{chemical} {and} {or} {medical} {and} {or} {environmental}] emergencies, call {insert name and/or number of emergency contact} {hours of operation} {24 hours a day} {7 days a week}}.



{See {additional} {sheet} {insert} {inside} {outer container} for {other} {directions for use} {information} {claims} {organisms}.}

Net Contents:

{{Batch} {Lot} No} {Manufacturing Date}:
{Product of USA} {Made in the USA}



MAGUARD® 1522

ORGANISM LIST

(Note to Reviewer: The list of organisms can be formatted into paragraph form using a comma to separate organisms.)

DISINFECTION PERFORMANCE: This product kills the following bacteria in 1 minute with 5% organic soil load and 400 ppm hard water on hard, non-porous surfaces:

Acinetobacter baumannii {(ATCC 19606)}
Clostridium perfringens {(ATCC 13124)}
Enterococcus faecalis {Vancomycin-Resistant} {(VRE)} {(ATCC 51575)}[§]
Escherichia coli O157:H7 {(ATCC 35150)}
Klebsiella pneumoniae {(ATCC 4352)}
Listeria monocytogenes {(ATCC 19111)}
Mycobacterium bovis (Tuberculosis surrogate) ‡
Pseudomonas aeruginosa {(ATCC 15442)}
Salmonella enterica {(ATCC 10708)}
Salmonella enterica serovar *Typhimurium* {(ATCC 13311)}
Serratia marcescens {(ATCC 13880)}
Shigella dysenteriae serotype 1 {(ATCC 29026)}
Staphylococcus aureus {(ATCC 6538)}
Staphylococcus aureus {Community-Associated Methicillin Resistant} {(CA-MRSA)} {(Genotype USA300)} {(CI 08001)}
Staphylococcus aureus {Methicillin-Resistant} {(MRSA)} {(ATCC 33592)}
Staphylococcus aureus {Vancomycin-Intermediate} {(VISA)} {(ATCC 700787)}
Staphylococcus epidermidis {Methicillin-Resistant} {(MRSE)} {(ATCC 51625)}
Streptococcus pneumoniae {(ATCC 6304)}[§]
Streptococcus pyogenes {(ATCC 19615)}
Vibrio cholerae {(ATCC 14035)}
Yersinia enterocolitica {(ATCC 35669)}

VIRUCIDAL* PERFORMANCE: This product kills the following viruses in 1 minute with 5% organic soil load and 400 ppm hard water on hard, non-porous surfaces:

Hepatitis B Virus {(HBV)} {(Duck Hepatitis B Virus)} {(Hepadna Virus Testing)}
Hepatitis C Virus {(HCV)} {(Bovine Viral Diarrhea Virus)} {(American Bioresearch Laboratories)}
Herpes Simplex Type 1 Virus {(ATCC VR-260)}
Herpes Simplex Type 2 Virus {(ATCC VR-734)}
Human Coronavirus Strain 229e {(ATCC VR-740)}
Human Immunodeficiency Virus Type 1 {(HIV-1)} {(Zeptomatrix Corporation)}
Human Rotavirus {(ATCC VR-2018)}
Influenza A {(A/Hong Kong/8/68-H3N2)} Virus {(SPAFAS)}
Norovirus {(Norwalk-like Virus)} {(Feline Calicivirus)} {(University of Ottawa)}
Poliovirus Type 1 {(ATCC 1562)} ‡
Respiratory Syncytial Virus {(RSV)} {(ATCC VR-26)}
Rhinovirus Type 37 {(ATCC 1147)} {(Organon Teknika Corp)}
Vaccinia Virus {(ATCC VR-156)}

ANIMAL VIRUCIDAL* PERFORMANCE: This product kills the following viruses in 1 minute with 5% organic soil load and 400 ppm hard water on hard, non-porous surfaces:

Avian Influenza A Virus {(Turkey/Wis/66-H9N2)} {(SPAFAS)}

FUNGICIDAL PERFORMANCE: This product kills the following fungi in 1 minute with 5% organic soil load and 400 ppm hard water on hard, non-porous surfaces:

Candida albicans {(ATCC 10231)}
Trichophyton interdigitale {(formerly *mentagrophytes*)} {(ATCC 9533)}

FOOD CONTACT SURFACE SANITIZING PERFORMANCE: This product is an effective food contact sanitizer in 1 minute and 400 ppm hard water on hard, non-porous surfaces against the following organisms:

Staphylococcus aureus {(ATCC 6538)}
Escherichia coli {(ATCC 11229)}

[§] Indicates a 3-minute contact time is necessary for this claim.

[‡] Indicates a 10-minute contact time is necessary for this claim.



TABLE OF CONTENTS

(Note to Reviewer: The Table of Contents is optional and may appear on labeling with the page numbers altered as necessary to reflect the pagination of the final printed label.)

ORGANISM LIST	2
TABLE OF CONTENTS	3
MARKETING CLAIMS	4
DIRECTIONS FOR USE	4
SANITIZATION.....	5
DISINFECTION	6
WATER TREATMENT	8
AGRICULTURAL OR HORTICULTURAL USES.....	9
POST-HARVEST TREATMENT	9
OIL FIELD, GAS PRODUCTION AND TRANSMISSION PIPELINE AND SYSTEMS (Not for use in CA.).....	10
ALTERNATE CONTAINER/DELIVERY SYSTEMS.....	10
STORAGE AND DISPOSAL	11
PRECAUTIONARY STATEMENTS	11
{SPANISH ADVISORY STATEMENTS}	12
GRAPHICS AND ICONS	12
{WARRANTY STATEMENT}	12



MARKETING CLAIMS

(Note to Reviewer: Marketing text is considered optional. Commas and the words “and” “or” can be added to phrases to make text grammatically correct.)

(Note to Reviewer: In the case where a location, surface, or marketing claim is not registered in the State of California the statement “(Not for use in CA.)” may be added to the relevant text.)

**THIS PRODUCT IS {A PEROXYACETIC ACID-BASED}
[{{SANITIZER} {/} {DISINFECTANT}}] [{{FORMULATED} {DEVELOPED}}] FOR {THE FOLLOWING USES}{:}**

(Note to Reviewer: The use sites under each heading can be used in any order, and may be placed in a paragraph or used as a list. Appropriate punctuation and an “and,” “&,” or “or” may be used to link use sites.)

INSTITUTIONAL/INDUSTRIAL SANITIZING OF PRE-CLEANED NON-POROUS FOOD CONTACT SURFACES IN:

- Dairies
- Wineries
- Breweries
- Food and Beverage Plants
- Disinfecting Poultry Premises
- Poultry Hatcheries
- Animal Housing Facilities
- Reverse Osmosis and Ultra Filtration Membranes and Distribution Systems (Not for use in CA.)

HARD SURFACE DISINFECTION IN:

- Hospitals
- Health Care Facilities
- Schools
- Colleges
- Veterinary Clinics
- Animal Life Science Laboratories
- Industrial Facilities
- Office Buildings
- Recreational Facilities
- Retail and Wholesale Establishments

BACTERIA, FUNGI, AND SLIME CONTROL IN:

- Pulp and Paper Mill Systems (Not for use in CA.)
- Cooling Water Systems (Not for use in CA.)



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on hard, non-porous surfaces.

(Note to Reviewer (General Considerations): Numbered instructions will be used if label space permits, otherwise they may appear in paragraph format. The list of organisms can be formatted into paragraph form using a comma to separate organisms. Unit abbreviations can be spelled out. Note symbols such as asterisks () and daggers (†) may be replaced with equivalent symbols. When choosing optional text, appropriate punctuation can be inserted or deleted. Equivalent use dilution ratios may be substituted within the directions.)*

{Please read entire label and use strictly in accordance with precautionary statements and directions.}

SANITIZATION

NOTE: FOR MECHANICAL OPERATIONS prepared use solution may not be used for subsequent sanitizing but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATIONS prepare a fresh sanitizing solution daily, or more often if the solution becomes diluted or soiled.

This product is {a peroxyacetic acid sanitizer} for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, conveyor belts, and pasteurizers in dairies, breweries, wineries, beverage and food processing/packing plants, egg processing/packing equipment surfaces, and eating establishments. This product is effective as a sanitizer when solution is prepared in water of up to 400 ppm hardness as CaCO₃. {This product has demonstrated greater than a 99.999% reduction of survivors after a 30 second exposure period in the AOAC Germicidal and Detergent Sanitizing Action of Disinfectants study.}

SANITIZING FOOD CONTACT SURFACES

{Effective against *Staphylococcus aureus* and *Escherichia coli*.}

Prior to sanitizing, remove gross food particles, then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 0.33 - 1.87 oz. of this product dissolved in 5 gal. of water (0.053% v/v concentration). This will provide 88 - 500 ppm of peroxyacetic acid. {At this dilution this product is effective against *Staphylococcus aureus* and *Escherichia coli*.} Use immersion, coarse spray, or circulation techniques as appropriate to the equipment. Allow surfaces to remain visibly wet for at least 60 seconds, or longer if specified by governing sanitary code. Drain thoroughly. Do not rinse.

SANITIZING, EATING, DRINKING, AND FOOD PREP UTENSILS

Remove gross food particles by a prescrape, a preflush and, when necessary, a presoak treatment. Wash with a recommended detergent. Rinse with clean water. Sanitize in a solution of 0.33 - 1.87 oz. of this product dissolved in 5 gal. of water. Immerse all utensils for at least 60 seconds, or longer if specified by governing sanitary code. Do not rinse. Drain and air dry.

SANITIZING TABLEWARE

For sanitizing tableware in low temperature warewashing machines, inject this product into the final rinse water at a concentration of 0.33 - 1.87 oz. of this product dissolved in 5 gal. of water. This will provide 88 - 500 ppm of peroxyacetic acid. Do not rinse. Air dry.

To ensure that the concentration of this product does not fall below 88 ppm peroxyacetic acid, periodically test the rinse solution with a suitable test kit and adjust the dispensing rate accordingly. Consult your technical service representative for assistance and further information on sanitizing tableware in warewashing machines.

FINAL SANITIZING BOTTLE RINSE

This product may be used as a final sanitizing rinse for returnable and non-returnable bottles at a 0.053% dilution (0.33 oz. of this product dissolved in 5 gal. of water). This will provide 88 ppm of peroxyacetic acid.

BATCH SANITIZATION (NON-FOOD CONTACT SURFACES) OF ULTRA FILTRATION AND REVERSE OSMOSIS (RO) MEMBRANES (Not for use in CA.)

This product can be used for the sanitization of ultra filtration, medical, and non-medical institutional/industrial reverse osmosis (RO) membranes and their associated distribution systems.

{This product has been shown to be an effective disinfectant when tested by AOAC and EPA methods.} {This product may not eliminate all vegetative microorganisms in reverse osmosis membranes and their associated piping systems due to their construction and/or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed.} Check with equipment manufacturer for membrane compatibility with this product.

Remove biological or organic fouling from the membrane or other parts of the system with an appropriate cleaner. Flush the system with RO permeate or similar quality water. Remove mineral deposits with suitable acidic cleaner prior to sanitizing the membranes with this product. Flush the system again with the RO permeate or similar quality water. Prepare an appropriate volume of 0.33% solution of the product (0.33 gal. of this product to 100 gal. of water). This will provide 568 ppm of peroxyacetic acid and 834 ppm hydrogen peroxide. Fill the entire water circuit to be sanitized with the dilute solution and allow the solution to reach a minimum of 20°C (68°F). Recirculate the dilute solution of this product for a minimum of 10 minutes. Allow membrane elements to soak in the solution for a minimum of 20 minutes. Rinse the RO system and test for residuals to ensure that there is less than 3 ppm peroxygen. Diverting product water to drain can reduce residuals.

BATCH SANITIZATION (NON-FOOD CONTACT SURFACES) OF PIPING SYSTEMS ASSOCIATED WITH RO MEMBRANES (Not for use in CA.)

Isolate incompatible equipment from piping system. This includes activated carbon filters and ion exchange equipment. Turn off power to ultraviolet light units. Estimate total volume of water contained in the system (tanks, rinse stations and piping). Prepare an appropriate volume of 0.33 - 0.5% of this product by adding 0.33 - 0.5 gal. of the product for every 100 gal. of solution prepared. Use RO permeate or similar quality water for dilution. This will provide 554 - 840 ppm peroxyacetic acid and 813 - 1232 ppm hydrogen peroxide. Recirculate the use solution through the system for a minimum of 4 hours. Process usage valves should be opened and closed to expose internals to the product. Completely drain the system of use solution. Thoroughly rinse the system by filling with RO permeate or similar quality water and recirculate before drainage. Repeat the process until test for residuals indicates there is less than 3 ppm peroxygen.

CONTINUOUS/INTERMITTENT ADDITION TO MINIMIZE THE ACCUMULATION OF BIOLOGICAL MATTER BETWEEN INTERMITTENT SANITIZING EPISODES IN PIPING SYSTEMS ASSOCIATED WITH RO MEMBRANES (NON-FOOD CONTACT SURFACES) (Not for use in CA.)

This product, as received or diluted, may be added continuously to the feed water system, between system sanitizing episodes, to aid in minimizing the regrowth/ accumulation of biological matter. The peroxygen residual in the system which will be effective will vary with the design and usage characteristics of the system. Adjust the addition rate of this product or the solution and periodically monitor residual peroxygen so that the desired effect is obtained. For continuous addition, do not exceed 7 ppm (0.33 oz. of product per 440 gal. of water) of this product. This will give 1 ppm peroxyacetic acid and 1.4 ppm hydrogen peroxide. For intermittent feed, do not exceed 750 ppm (8.5 oz. of product per 100 gal. of water) of this product. This will give 110 ppm peroxyacetic acid and 160 ppm hydrogen peroxide.

DISINFECTION

(Note to Reviewer: If hospitals, medical devices, and/or stainless-steel surfaces are listed on the label, one of the following FDA/EPA Memorandum of Understanding statements must be used.)

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

(OR)

This product is not for use on medical device surfaces.

FOR HEALTH CARE, INSTITUTIONAL, AND INDUSTRIAL USE.

{This product disinfects as it cleans in one operation.} This product can be used to disinfect floors, walls and other hard non-porous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, doors, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, non-porous glazed porcelain, glazed ceramic, plastic (such as polypropylene and polyethylene), {stainless steel,} glass, aluminum, non-porous baked enamel, chrome, laminated or painted surfaces or sealed stone. This product should not be used on marble or brass surfaces.

{{[Areas of use in hospitals:] {This product may be used for}} surgical and obstetrical suites; housekeeping surfaces; physical therapy departments; nursing services; dental facilities; autopsy facilities; intensive care units; pharmacies; and clinical laboratories.} {This product may {also} be used in nursing homes, {other} healthcare facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, office buildings, recreational facilities, industrial facilities; hotels; retail facilities; office buildings; retail and wholesale establishments.}

Dilute this product with the appropriate amount of water to an effective concentration of 1135 ppm peroxyacetic acid and 1665 ppm hydrogen peroxide (0.85 oz. per gal. of water). Pre-clean visibly soiled areas. Apply solution with a cloth, mop, sponge, auto-scrubber, or hand pumped trigger sprayer and allow to remain visibly wet for 1 minute to kill bacteria, viruses*, and fungi as cited on the label. Use a 3-minute contact time for *Streptococcus pneumoniae* and Vancomycin Resistant *Enterococcus faecalis*. Use a 10-minute contact time for Poliovirus Type 1 and Tuberculocidal activity. This product is effective against tuberculosis (*Mycobacterium bovis*) at ambient temperature (at least 22°C / 72°F). Allow surface to air dry. Prepare a fresh solution daily or more often if the use solution becomes visibly soiled, clouded, or diluted.

BLOODBORNE PATHOGEN INSTRUCTIONS (*Note to Reviewer: Heading is optional. If instructions used, all indented text must be included.*):

***KILLS HIV-1 (AIDS VIRUS), HBV, AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS**

PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of surfaces/ objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus, or Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HBV, AND HCV ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: {{Wear protective latex gloves, gowns, masks, and eye protection} {Specific barrier protection items to be worn when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, and eye protection}}.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application. Contact Time: HIV-1, HBV, and HCV are inactivated after 1 minute of contact.

Infectious material: Blood and other bodily fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

HARD SURFACE DISINFECTION

{This product disinfects as it cleans in one operation.} This product can be used to disinfect floors, walls and other hard non-porous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, non-porous glazed porcelain, plastic (such as polypropylene and polyethylene), {stainless steel,} or glass.

{{{Areas of use in hospitals:} {This product may be used for}} surgical and obstetrical suites; housekeeping services; physical therapy departments; nursing services; autopsy facilities.} {This product may {also} be used in {nursing homes,} {{other} {health-care facilities,}} schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, dietary areas, office buildings, recreational facilities, retail and wholesale establishments.}

Dilute this product with the appropriate amount of water to an effective concentration of 1135 ppm peroxyacetic acid and 1665 ppm hydrogen peroxide (0.85 oz. per gal. of water). Pre-clean visibly soiled areas. Apply solution with a cloth, mop, sponge, auto-scrubber, or hand pumped trigger sprayer and allow to remain visibly wet for 1 minute to kill bacteria, viruses*, and fungi as cited on the label. Use a 3-minute contact time for *Streptococcus pneumoniae* and Vancomycin-Resistant *Enterococcus faecalis*. Use a 10-minute contact time for Poliovirus Type 1 and Tuberculocidal activity. This product is effective against tuberculosis (*Mycobacterium bovis*) at ambient temperature (22°C / 72°F). Allow surface to air dry. Prepare a fresh solution daily or more often if the use solution becomes visibly soiled, clouded, or diluted.

COMBINATION DISINFECTION AND CLEANING

This product is effective against *Staphylococcus aureus*, *Salmonella enterica*, *Pseudomonas aeruginosa*, *Trichophyton interdigitale*, and *Escherichia coli* O157:H7 at 0.08% (0.5 oz. of this product per 5 gal. of water) {in hard water (400 ppm as CaCO₃)} {and 5% soil {(fetal bovine serum)}} on hard, non-porous surfaces. Pre-clean visibly soiled areas. Apply solution with mop, cloth, sponge, brush, scrubber, or coarse spray device, or by soaking to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted.

{This product is designed for use in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries, and livestock quarters.} {When used as directed, this product is specifically designed to disinfect, deodorize and clean inanimate, hard, surfaces such as walls, floors, sink tops, furniture, operating tables, kennel runs, cages, and feeding and watering equipment.} {{In addition,} this product will deodorize those areas which are generally hard to keep smelling fresh such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.}

Rinse with potable water after use on surfaces that contact food, feed, or drinking water. Prepare a fresh solution for each use.

DISINFECTION OF POULTRY PREMISES, TRUCKS, COOPS, AND CRATES

POULTRY HATCHERY DISINFECTION

Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.08% (0.5 oz. of this product per 5 gal of water.) solution of this product and allow to remain visibly wet for 10 minutes. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse. See your technical representative for specific recommendations for all cleaning and rinsing requirements.

DISINFECTION AND DEODORIZING OF ANIMAL HOUSING FACILITIES {(BARN, KENNELS, HUTCHES, ETC.)}

Do not use in milking stalls, milking parlors, or milk houses. Remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter, and gross soils from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering equipment. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.08% (0.5 oz. of this product per 5 gal. of water) solution of this product and allow to remain visibly wet for 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

WATER TREATMENT

BIOFOULING CONTROL IN PULP AND PAPERMILL SYSTEMS (Not for use in CA.)

For use in the manufacture of paper and paperboard intended for food contact and non-food contact uses. This product can be used to control bacteria, fungi, and freshwater organisms in paper, paperboard, or nonwoven process water and influent water systems. Suitable dosing points include but are not limited to: stock chests, pulpers, the white water loop and white water storage systems, and influent water streams.

Contamination with other chemicals could result in product decomposition.

- Add this product at a point in the system where uniform mixing and even distribution will occur.
- Use 0.1 - 0.5 lb. of this product per 1000 gal. of solution as a continuous or intermittent slug treatment. This will provide 1.8 - 9 ppm peroxyacetic acid (12 - 60 ppm of this product). Repeat treatment as required to maintain control.

ANTIMICROBIAL RINSE OF PRE-CLEANED OR NEW RETURNABLE OR NON-RETURNABLE CONTAINERS

To reduce the number of non-pathogenic beverage spoilage organisms. Effective against *Aspergillus versicolor* (ATCC 9577), *Byssoschlamys fulva* (ATCC 10099), *Pediococcus damnosus* (ATCC 29358), *Lactobacillus buchneri* (ATCC 4005), and *Saccharomyces cerevisiae* (ATCC 47058).

1. Prepare use solution by adding 9.85 oz. of this product to 5 gal. potable water. This provides 2,632 ppm peroxyacetic acid.
2. Apply antimicrobial rinse at a temperature of 40°C - 60°C (104°F - 140°F) and allow a minimum 7-second contact period.
3. Allow containers to drain thoroughly, and then rinse with sterile or potable water.

TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS

This product can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables for the control of bacteria and fungi in commercial operations and packinghouses.

BATCH, CONTINUOUS OR SPRAY SYSTEM PROCESSES: Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method. Add this product at a rate no more than 80 ppm peroxyacetic acid to the use solution. This can be accomplished by initially adding 1.0 oz. of this product per 16.4 gal. of water. The fruits and vegetables can be continuously sprayed (using coarse spray) or submerged (dipped) in the resulting solution. Periodic or continuous addition of this product to maintain the required concentration may be added as necessary. Contact time of 60 seconds is recommended to ensure efficacy. A potable water rinse is not required. This product is not intended for use in primary flumes prior to the point of the first dewatering stage.

FOR DISINFECTION OF SEWAGE AND WASTEWATER EFFLUENTS IN TREATMENT PLANTS

Use this product to treat sewage and wastewater effluent related to public and private wastewater treatment plants. This product can be applied directly to the effluent or may be used with an appropriate activator such as hydrogen peroxide or other technology. This product may be applied to effluent water discharged from trickle bed or percolating fluidized bed filters. The application rate for individual facilities will depend on the degree of bioloading of the effluent stream to be discharged and the local microbial discharge limit. Adjust application rate to meet the need of the individual facility.

1. Add this product to effluent water at a concentration of 0.5 - 15 ppm. Allow contact time of approximately 15 - 60 minutes.
2. The maximum amount of peroxyacetic acid that can be discharged from the treatment facility is 1 ppm. Use an appropriate peroxyacetic acid test kit analyzer to ensure that this level is not exceeded.

INFLUENT WATER SYSTEMS (Not for use in CA.)

This product should be fed continuously to incoming fresh water streams (non-potable use only) at dosages ranging from 10 - 975 ppm peroxyacetic acid (65 - 6500 ppm of this product).

MILL PROCESS WATERS (Not for use in CA.)

- **Continuous Feed:** This product should be fed continuously at dosages ranging from 10 - 975 ppm peroxyacetic acid (65 - 6500 ppm of this product). This range is equivalent to 0.13 - 13 lbs. of this product per ton (dry basis) of pulp or paper produced.
- **Intermittent Feed:** This product should be fed intermittently (6 - 8 times per day) at dosages ranging from 10 - 975 ppm peroxyacetic acid (65 - 6500 ppm of this product). This range is equivalent to 0.13 - 13 lbs. of this product per ton (dry basis) of pulp or paper produced.
- **Shock Dose:** This product should be shock dosed at dosages ranging from 98 - 2048 ppm peroxyacetic acid (648 - 13,638 ppm of this product). This range is equivalent to 1.3 - 27.3 lbs. of this product per ton (dry basis) of pulp or paper produced.

CONTROL OF SLIME FORMING BACTERIA AND BIOFOULING IN ONCE-THROUGH AND RECIRCULATING COOLING WATER (COOLING TOWERS, EVAPORATIVE CONDENSERS, AIR WASHERS) AND ORNAMENTAL OR RECREATIONAL WATER FEATURES (Not for use in CA.)

Severely fouled systems must be cleaned before adding this product. This product must be added in the water system directly, and not mixed with any other chemicals or additives. Never add this product into any feeding device, such as shot feeders, filter housings, bypass feeders, or miscellaneous piping of any kind, because dangerous acute decomposition can occur. Discontinue the use of chlorine or bromine products prior to using this product. Contamination with other chemicals could result in product decomposition. Add this product to only water at a point in the system where uniform mixing and even distribution will occur.

For shock (slug) treatment for moderately to severely fouled systems add 5 - 20 oz. of this product per 1000 gal. of process water (7 - 27 ppm peroxyacetic acid). Repeat as necessary until microbiological control is evident. Thereafter, to maintain control use (1.5 - 7.5 oz.) of this product per 1000 gal. of process water (2 - 10 ppm of peroxyacetic acid) as a continuous treatment method. Continuous dosing methods usually require 1.5 - 5 oz. per 1000 gal. of water (2 - 7 ppm peroxyacetic acid) to achieve adequate results.

Intermittent dosing treatment usually require dose cycles of a minimum once per every other day, up to 6 times per 24 hours. Recommended rates for intermittent dose cycles are 5 - 10 oz of this product per 1000 gal. of process water (7 - 14 ppm peroxyacetic acid).

AGRICULTURAL OR HORTICULTURAL USES

There is a Restricted-Entry Interval of zero (0) hours after the use of this product. This product should never be mixed or combined with any other pesticide or fertilizer. Upon soil contact this diluted product decomposes rapidly to oxygen, carbon dioxide, and water. This product may be harmful to fish if exposed on a continuous basis at concentrations of 1 ppm or more of active peroxyacetic acid. Meter this product into pressurized pipes using a plastic or stainless steel injection/backflow device installed far enough upstream from the equipment to insure thorough mixing. For open flowing bodies of water, apply this product as far upstream as possible to allow adequate mixing prior to the flow entering any larger body of water. If open pouring of this product is required, pour product as close to the surface of the water as possible to reduce odor exposure.

TREATMENT OF AGRICULTURAL OR IRRIGATION WATER SYSTEMS (SAND FILTERS, HUMIDIFICATION SYSTEMS, STORAGE TANKS, PONDS, RESERVOIRS, CANALS):

For the control of sulfides, odor, slime and algae in water systems, apply this product at 2 - 10 ppm active peroxyacetic acid. This feed rate equals 15 - 75 oz. of this product per 10,000 gal. of water. Repeat dose as necessary to maintain control, which will vary with seasonal conditions. For prevention of algae, some systems may require continuous low-level dosing during warm sunny periods (2 - 5 ppm peroxyacetic acid).

Drip Irrigation Systems: To clean slime and algae from drip system filters, tapes and emitters, meter this product at the rate of 7.5 - 15 oz. of this product per 1000 gal. of water (10 - 20 ppm peroxyacetic acid). When required during normal irrigation cycles, use this product at the recommended dose for a minimum of 30 minutes. Thereafter, the irrigation cycle should be discontinued and the line should not be flushed.

POST-HARVEST TREATMENT

FOR TREATMENT OF RAW, UNPROCESSED FRUIT AND VEGETABLE SURFACES

This product can be applied as a dip or spray to control the growth of non-public health microorganisms that may cause decay and/or spoilage on raw, post-harvest fruits and vegetables during the washing process. This product can be applied during physical cleaning processes, including at the roller spreader, washer manifold, dip tank, on the brushes or elsewhere in the washing process prior to, simultaneously with, or after detergent wash.

1. Prepare use solution by diluting 1 oz. of this product per 16 gal. of potable water. This will provide 85 ppm peroxyacetic acid and 125 ppm hydrogen peroxide.
2. Apply the use solution using a coarse spray directed at the fruits or vegetables, or by soaking the fruits and vegetables in the use solution. Allow a contact time of at least 45 seconds.
3. The treated produce can be drain dried without a potable water rinse.
4. Do not reuse solution after treatment.

FOR THE TREATMENT OF PROCESSED FRUITS AND VEGETABLES AND PROCESS WATERS TO CONTROL GROWTH OF NON-PUBLIC HEALTH MICROORGANISMS THAT CAN CAUSE SPOILAGE

1. Prepare use solution by diluting 1.5 oz. of this product per 25 gal. of potable water. This will provide 80 ppm peroxyacetic acid and 117 ppm hydrogen peroxide.
2. Apply the use solution as a spray or dip. Allow a contact time of at least 45 seconds. No rinse following application is required. This use complies with the requirements of 21 CFR 173.315 (a) 5.
3. The treated produce can be drain dried without a potable water rinse.
4. Do not reuse solution after treatment.

OIL FIELD, GAS PRODUCTION AND TRANSMISSION PIPELINE AND SYSTEMS (Not for use in CA.)

For antimicrobial use with aqueous treatment fluids in subterranean oil and gas field well operations such as well drilling, formation fracturing, productivity enhancement and secondary recovery.

This product can be used for control of slime forming and spoilage bacteria, yeast and fungi, and anaerobic sulfate reducing bacteria that lead to reservoir souring and metal corrosion. This product must be introduced through a closed mixed/loading and delivery transfer system equipped with a metering device that is appropriate for its intended uses.

DRILLING MUDS, FRACTURING FLUIDS, WELL SQUEEZED FLUIDS

For the preservation of drilling muds, workover and completion fluids and other products susceptible to contamination, pre-mix with the fluid or add directly at the point of use at 3.75 – 75 oz. of this product per 1000 gal. of water (5 – 100 ppm peroxyacetic acid) as required. Depending on the severity of the contamination, initial application may be added up to 749 oz. of this product per 1000 gal. of water (1000 ppm peroxyacetic acid).

FLOODING, INJECTION AND PRODUCED WATER

For water flooding operations, add initially at 3.75 - 75 oz. of this product per 1000 gal. of water (5 - 100 ppm) and repeat until control is achieved. Subsequent treatment may be continued on a weekly basis or as required. Injection wells associated with gas storage systems may be treated with up to 100 ppm active peroxyacetic acid when diluted in the formation water. Any additional top-up water should be treated as required.

For hydrostatic systems, apply 3.75 - 75 oz. of this product per 1000 gal. of water (5 - 100 ppm peroxyacetic acid) depending on the water quality and the duration of the shut-in.

PIPELINE AND TANK MAINTENANCE

For microbial control in water-bottoms in crude and refined hydrocarbon storage tanks, piping and transportation systems. Apply 3.75 - 75 oz. of this product per 1000 gal. of water (5 - 100 ppm peroxyacetic acid) in the aqueous phase, directly injected into the water-bottom, pipeline or may be added to the hydrocarbon phase. Treatment may be applied daily or monthly for both storage and transportation systems as needed.

ALTERNATE CONTAINER/DELIVERY SYSTEMS

{eraDOCate Closed System Cap Instructions:}

DILUTION CHART

For hard surface disinfection {/virucide*}, add the contents of one concentrate container to the following amount of water: (Note to Reviewer: Equivalent use dilutions may be substituted.)

Concentrate Container Size {(eraDOCator-60)}	{Potable} Water Dilution	Contact Time
0.107 oz.	16 oz.	1 minute
0.213 oz.	32 oz.	1 minute
0.85 oz.	1 gal.	1 minute
2.13 oz.	2.5 gal.	1 minute
4.25 oz.	5 gal.	1 minute

MIXING INSTRUCTIONS

WARNING: Although this is a closed mixing system, all PPE requirements contained in the precautionary statements section of this label still apply. NEVER remove the closed system cap from the concentrate container.

1. Fill mix container with the proper amount of potable water, as specified in the above chart.
2. Remove one concentrate container with the closed system cap from the storage case.
3. Keep the mix container in an upright position and couple the concentrate container to the mix container by threading the closed system cap onto the mix container.
4. Remove the safety tab from the closed system cap.
5. Rotate the concentrate container in a clockwise motion until the threads bottom out, indicating a fully coupled position.
6. In the fully coupled position the concentrate will flow into the mix container. Once all concentrate has moved into the mix container, invert the mix container several times, allowing the diluted solution to rinse the concentrate container thoroughly.

7. Return the mix container to the upright position and allow enough time for the concentrate container to fully empty into mix container.
8. Remove the closed system cap from the mix container by rotating the cap counterclockwise and dispose of the concentrate container with the closed system cap in accordance with this label.
9. Apply the 1135 ppm peroxyacetic acid and 1665 ppm hydrogen peroxide solution, contained in the mix container, in accordance with this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original containers in a cool, well-vented area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

(Note to Reviewer: One or more of the following paragraphs for Container Handling will be selected, depending on packaging use/type.)

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different (concentrated) product to be diluted in the container.

{For non-refillable containers equal to or less than 5 gal.}

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

{For non-refillable containers greater than 5 gal.}

Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand container on its end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat the procedure two more times. Then offer for recycling or dispose in a sanitary landfill, or by incineration, if allowed by state and local authorities by burning.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. CORROSIVE. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through the skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes on skin or on clothing. Wear goggles and/or face shield, coveralls over long-sleeved shirt and long pants, socks, chemical resistant footwear, and chemical-resistant gloves when handling. Wear a minimum of a NIOSH approved respirator with an organic vapor (OV) cartridge with any combination N, R, or P filter with NIOSH approval number prefix TC-84A. Higher level respirators with OV filters that are NIOSH approved for particulates can also be used. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

STRONG OXIDIZING AGENT. Corrosive. Mix only with water. Product must be diluted in accordance with label directions prior to use. This product is not combustible; however, at temperatures exceeding 156°F (69°C), decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish, and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the US Environmental Protection Agency.



{SPANISH ADVISORY STATEMENTS}

(Note to Reviewer: This statement is optional except when used on labels with agricultural uses.)

{SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE.

IF YOU DO NOT UNDERSTAND THE LABEL, FIND SOMEONE TO EXPLAIN IT TO YOU IN DETAIL.}



GRAPHICS AND ICONS

(Note to Reviewer: These are representative icons for use sites/application methods listed in the location/surfaces section of this label that may appear on the label with the appropriate directions for use, PPE or package type.)

{Baby Drowning in Bucket
Warning Graphic}

{Made in USA Logo/Flag}

{Recycling Logo}

{Mixing Chemical Warning
Graphic}



{WARRANTY STATEMENT}

(Note to Reviewer: This statement is optional.)

Read Product Safety Data Sheet prior to use. PRODUCT WARRANTY, DISCLAIMER AND LIMITATION OF LIABILITY ARE FOUND on the Product Safety Data Sheet. Unless inconsistent with applicable law, use of Product signifies agreement with these provisions.

Lea la Hoja de Seguridad del Producto antes de usarlo. LA GARANTIA DEL PRODUCTO, DECLINACION Y LIMITACION DE RESPONSABILIDAD SE ENCUESTRAN en la Hoja de Seguridad del Producto. A menos de que sea inconsistente con la ley, el uso del producto significa acuerdo con estas disposiciones.