



010352-00035-060899

Printed: 07:38:46 Tuesday, 13 Jul, 1999 # 15 / 3526

Systems Integration Group, Inc.

PM 33

10352-35

6-8-99

1072



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

JUN 08 1999

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Union Carbide Corporation, Inc.
P. O. Box 670
Bound Brook, NJ 08805

Attention: Joan E. Young

Subject: UCARCIDE 750 Antimicrobial
EPA Registration Number 10352-35
Your Submission Dated May 12, 1999
EPA Received Date May 18, 1999

The submitted final printed labels have been placed in your product file. They have not been reviewed in accordance with the PR Notice 82-2.

If you have any questions concerning this letter, please contact Karen M. Leavy-Munk at (703)-308-6237.

Sincerely yours,

A handwritten signature in cursive script, appearing to read "Martha Swindell".

Marshall Swindell
Product Manager 33
Regulatory Management Branch I
Antimicrobial Division(7505C)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PIGMENTS AND FILLER SLURRIES FOR PAPER AND PAPERBOARD (For use in food contact pigments and filler slurries)

UCARCIDE® 750 Antimicrobial may be used to control the growth of microorganisms which contaminate various systems used in production of paper and paper products. UCARCIDE® 750 Antimicrobial should be added at a point of uniform mixing.

Use from 0.1 to 0.6 lbs (1.4 to 8.2 fluid ounces) of UCARCIDE® 750 Antimicrobial per 1000 pounds of dry powder, to produce a concentration from 100 to 600 ppm (based on slurry solids) in the mixed slurry.


WATER BASED COATINGS FOR PAPER AND PAPERBOARD (For use in non-food contact coatings only)

Use from 0.1 to 0.6 lbs (1.4 to 8.2 fluid ounces) of UCARCIDE® 750 Antimicrobial per 1000 pounds of dry powder, to produce a concentration from 100 to 600 ppm (based on slurry solids) in the mixed slurry.

GENERAL PRESERVATIVE APPLICATIONS (Except food contact applications)

UCARCIDE® 750 Antimicrobial is recommended for use in solutions, emulsions, slurries, adhesives and other aqueous-based liquid products to control the growth of bacteria and fungi.

For effective preservation, add UCARCIDE® 750 Antimicrobial to the product formulation at a rate of 200 to 2000 ppm based on the water content of the product (0.2 to 2.0 lbs UCARCIDE® 750 Antimicrobial/ 1000 lbs water content). Mix uniformly.



_____ 50.0%
_____ 50.0%
_____ 100.0%

CHILDREN

TREATMENT

Get immediate medical attention.
al attention.

en. If symptoms persist, call a physician.
drink. Seek medical advice with urgency.
mucosal damage may contraindicate

ONARY STATEMENTS.

PORATION

EPA Est. No. 10352-WV-2
UCC-L90209 (G04/99)
Made In USA
PE0499

OSIVE LIQUID, ACIDIC, ORGANIC, N.O.S.
(CONTAINS GLUTARALDEHYDE)

ID NUMBER
UN 3265

PG II

1-800-UCC-HELP (1-800-822-4357)
E USA 01-304-744-3487

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or your Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal Containers or Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures approved by state and local authorities. **Plastic Containers:** May be incinerated, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Metal Containers:** Must not be incinerated. Do not cut or weld on or near metal containers.

NOT REVIEWED

~~In~~ Accordance with FR Notice 82-2.
~~Based~~ on Draft Labeling Dated

2072