



010352-00038-060899

Printed: 07:39:43 Tuesday, 13 Jul, 1999 # 18 / 3529

Systems Integration Group, Inc.

ACCEPTED

JUN 08 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. **10352-38**

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRESERVATIVE USE

UCARCIDE® 250 Preservative is recommended for use in aqueous or water containing products and systems, including industrial, institutional and consumer in-can processes and products, to control the growth of bacteria and fungi. For effective preservation, add UCARCIDE® 250 Preservative to the product formulation at a rate of 0.02% to 0.20% based on the water content of the product (0.2 to 2.0 lbs UCARCIDE® 250 Preservative/1000 lbs water content). Mix uniformly.

PRESERVATIVE FOR CONCENTRATES

For use in concentrates where effective preservation is needed after dilution, add UCARCIDE® 250 Preservative to the product formulation at a rate such that the diluted end-use product will contain 0.02% to 0.20% UCARCIDE® 250 Preservative.

At no time during the preservation process should the level of UCARCIDE® 250 Preser. exceed 2.0%.

REVERSE OSMOSIS MEMBRANES

For effective preservation of reverse osmosis elements (where approved for compatibility by membrane manufacturer), immerse element in a tank containing 0.2% to 2.0% UCARCIDE® 250 Preservative.

UCARCIDE® 250 Preservative can also be added to in-line recirculating systems for preservation of installed out-of-service reverse osmosis equipment (where approved for compatibility by membrane manufacturer). Add 0.2% to 2.0% UCARCIDE® 250 Preservative to the tank in the circulating system. Maintain UCARCIDE® 250 Preservative concentration by periodic addition to counteract any system leakage.

CONCRETE ADMIXTURES

For effective preservation of concrete admixtures, add UCARCIDE® 250 Preservative to the product formulation at a rate of 2000 to 8000 ppm based on the weight of the admixture (2.0 to 8.0 lbs UCARCIDE® 250 Preservative/1000 lbs. concrete admixture). Mix uniformly.

UCARCIDE® 250 PRESERVATIVE

INDUSTRIAL, AND CONSUMER

_____ 50.0%

_____ 50.0%
_____ 100.0%

CHILDREN

TREATMENT

Get immediate medical attention.

Get medical attention.

If symptoms persist, call a physician.

Seek medical advice with

Medical damage may contraindicate the

HAZARD STATEMENTS

PRECAUTION

EPA Est. No. 10352-WV-2
UCC-L90290 (E)
Made In USA
PE1097

TOXIC LIQUID, ACIDIC, ORGANIC, N.O.S.
CONTAINS GLUTARALDEHYDE

UN 3285

PG II

1-800-UC-C-HELP (1-800-822-4357)

SA 01-304-744-3487

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or your Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal Containers or Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures approved by state and local authorities. **Plastic Containers:** May be incinerated, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Metal Containers:** Must not be incinerated. Do not cut or weld on or near metal containers.

PM 33

10352-38

6-8-99

1081