

MAY 10 1990

Frances Janison  
Ford's Chemical & Service, Inc.  
2739 Pasadena Boulevard  
Pasadena, CA 77502

Subject: Ford's General Outdoor Spray  
EPA Registration Number 10370-32  
Your submission dated 4-5-90  
Revised Labeling

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Signal word Caution should be of a larger type size and placed above the "Keep Out of Reach of Children Statement".
2. Add a preharvest interval of 1 day for apples.
3. Delete the preharvest interval for tomatoes, peppers, cucumbers, summer squash.

Submit five copies of final printed labeling before you release this product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Dennis H. Edwards, Jr.  
Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

51667.2:I/C:Kumar:WP4.2:C.Disk:KENCO:05/09/90:EK:SW:VO:DD:EK

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

294

FORD'S LIQUID SEVIN® 2L  
BRAND CARBARYL INSECTICIDE

For Use on Ornamentals, Flowers, Lawns,  
Fruit Trees and Vegetables

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate.....	23.00%	✓
INERT INGREDIENTS.....	77.00%	✓
TOTAL.....	100.00%	✓

\*\*\*\*\*

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

\*Registered trademark of Union Carbide Agricultural  
Products, Inc.

MAY 10 1977

KEEP OUT OF REACH OF CHILDREN - (CAUTION)

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

10370-32

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing vapors or  
spray mist. Avoid contact with eyes, skin or clothing. Wash  
thoroughly after handling. Do not take internally. Keep children  
and animals off treated lawns until these areas are dry. Avoid  
contamination of food.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible,  
cholinesterase inhibitor. Atropine is antidotal.

\*\*\*\*\*

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not  
apply directly to water or wetlands. Do not contaminate water by  
cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment  
or residue on blooming crops or weeds. Do not apply this product  
or allow it to drift to blooming crops or weeds if bees are  
visiting the treatment area.

\*\*\*\*\*

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner  
inconsistent with its labeling. Subsequent planting of food and  
feed crops in treated soil must be made with crops on this or  
other Carbaryl labels. Use 3 tablespoons of concentrated Liquid  
Sevin 2L for 1 gallon of water and apply thoroughly. Do not  
store diluted spray. Make new dilution for each use.

\*\*\*\*\*

**BEST AVAILABLE COPY**

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not reuse empty bottle. Wrap and put in trash.

\*\*\*\*\*

**ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES**

Japanese Beetles, Rose Aphids, Bagworms, Elm Leaf Beetles, Birch, Boxwood and Oak Leafminers, Rose Slugs, Periodical Cicadae (17 Year Locust), Scale Insects, Eastern Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips - Spray when first signs of damage appear and repeat every 7 to 10 days as necessary. **IMPORTANT - DO NOT USE ON BOSTON IVY.**

**APPLES**

**IMPORTANT:** To avoid possible fruit thinning on apples, do not apply within 15 days after petal fall. Apple Maggot, Rosy Apple Aphid, Fruit Tree Leaf Rollers, Eastern Tent Caterpillar, European Apple Sawfly, Apple Rust Mite. *May be applied within 1 day of harvest.*

**PEARS**

Pear Psylla, Japanese Beetle, Green Fruitworm, Pear Leaf Blister Mite, Pear Rust Mite - Make application between bud break and bloom, and a second at petal fall and then 3 to 4 week intervals for 3 to 4 months as necessary. May be applied within 1 day of harvest.

**PEACHES, APRICOTS and NECTARINES**

Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Codling Moth, Fruit Tree Leaf Roller, June Beetles, Japanese Beetle, European Earwig - Do not apply within 1 day of harvest for Peaches, and within 3 days of harvest for Apricots and Nectarines. For Peach Tree Borer, apply 2 or 3 sprays to trunk from the ground to scaffold limbs, timed with moth flight. For Codling Moth, Oriental Fruit Moth and Plum Curculio, apply at petal fall and every 10 - 14 days thereafter until control is achieved.

**LAWN INSECTS**

Sod Webworms, Earwigs, Grasshoppers, Fleas and Ants - Use 2 tablespoons per gallon of water for each 50 sq. ft. of established lawn covered. For best results, use a hose-end sprayer and apply after mowing and watering area to be treated. Do not water for two days after treatment. Repeat in 2 to 3 weeks as necessary.

**BEST AVAILABLE COPY**

494

**TOMATOES, PEPPERS, CUCUMBERS, SUMMER SQUASH**

Pickleworm, Fruitworms, Melon Worms, Hornworms, Fall Armyworms, Corn Borer - Apply when insects first appear. Repeat at 7 to 10 day intervals. ~~May be applied within 1 day of harvest.~~

**NOTICE:** Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

FORD'S CHEMICAL & SERVICE, INC.  
2739 Pasadena Blvd. Pasadena, Texas 77502

031687FCS  
10/87