

10370-33

1042

ACCEPTED

SEP 21 1988

10370-33

FOAMSPRAY PRODUCTS 57% MALATHION
Emulsifiable Concentrate

CONTROLS: Aphids, Bagworm, Cabbage Worm, Codling Moth, Cucumber Beetle, Fleas, Flea Beetle, Mealybugs, Pecan Nut Casebearer, Plum Curculio, Potato Leafhopper, Red Spiders, Rose Leafhopper, Spider Mites, Thrips, Vine Borer and Whiteflies

ACTIVE INGREDIENT:

Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.....57.00%
INERT INGREDIENTS**.....43.00%
TOTAL.....100.00%

**This product contains xylene range aromatics

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with soap and water.

IF IN EYES: Flush with water for 15 minutes. Contact a physician.

IF INHALED: Remove victim to fresh air and apply artificial respiration if indicated.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing vapors or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat and open flame.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store product in original labeled container in a cool, dry, locked place out of the reach of children.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

PESTICIDE DISPOSAL: Securely wrap container in several layers of newspaper and discard in trash.

ORNAMENTALS

For control of aphids, spider mites, whiteflies, mealybugs, thrips, Japanese beetle adults, four-lined leaf bug, tarnished plant bug, rose leafhopper, and potato leafhopper, use 1-1/2 pints per 100 gallons of water or 2 teaspoons per 1 gallon of water. Repeat in 12 to 14 days if necessary. Do not use on Boston, Maidenhair or Pteris ferns, or Crassula. Do not apply to Petunias.

VEGETABLES

For aphids, spider mites, cucumber beetles, flea beetles, leafhoppers, and cabbage worms on cole crops; pickle worms and vine borer on squash - spray thoroughly with 2 pints concentrate per 100 gallons water, or 2 teaspoons per gallon. Make no application to any vegetable within 7 days of harvest.

FRUITS AND NUTS

For control of codling moth on apples and pears; plum curculio on peaches, plums, quinces - spray with 2 pints concentrate in 100 gallons of water, or 2 teaspoons per gallon. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Make no application within 7 days of harvest. For pecan nutcase bearers, apply 1 quart concentrate per 100 gallons of water, or 2 teaspoons per gallon. Start treatment after eggs begin to hatch. Make no treatment within 7 days of harvest.

FLEAS IN DOG KENNELS AND YARDS

Spray infested areas with 10 tablespoons concentrate in 1 gallon of water, at the rate of 1 gallon of mixture to 1,000 sq. ft. Remove manure and debris before spraying.

SHOULD AFFORD PROTECTION UP TO 30 DAYS. REPEAT APPLICATION IF NECESSARY. DO NOT USE ON PETS AND OTHER ANIMALS. TO CONTROL FLEAS ON PETS AND OTHER ANIMALS, USE AN INSECTICIDE RECOMMENDED FOR THIS PURPOSE.

Manufactured by

FORD'S CHEMICAL & SERVICE, INC.
2739 Pasadena Blvd.
Pasadena, TX 77502

AVAILABLE COPY