

10370-37 2-25-91

DURSBAN® 2E

INSECTICIDE
—Controls—

Chinch Bugs, Sod Webworms (Lawn Moth), Bagworms,
Cutworms, Grasshoppers,
INDOOR PEST CONTROL

To be applied only by or under the Supervision of Pest Control Operators and other Trained Personnel responsible for Insect Control Programs.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 24.29%
Xylene Range Aromatic Solvent ~~65.71%~~

INERT INGREDIENTS:

Total 100.0%

Contains 2 pounds Chlorpyrifos per gallon
DURSBAN® — Trademark of The Dow Chemical Company

U.S. Patent No. 3,244,586

* CONTAINS XYLENE RANGE AROMATIC SOLVENT

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **IF ON SKIN:** Wash thoroughly with soap and warm water. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically, atropine is an antidote. 2-PAM is also antidote and may be administered in conjunction with atropine.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not take internally. Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing. Avoid breathing vapors and spray mist. Avoid spraying food crops. Handle concentrate in a ventilated area. Keep children and pets off treated area until spray has dried.

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

EPA Est. No. 10370-TX-1

EPA Reg. No. 10370-37 NET CONTENTS: ONE GALLON

Manufactured By

Chemical & Service, Inc.

2739 PASADENA BLVD.
PASADENA, TEXAS 77502

ford's

ACCEPTED
AUG 28 1984
Un...
Fung...
as...
registered under
EPA Reg. No. 10370-37

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL AND CHEMICAL HAZARDS

Keep container closed. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Handle concentrate in a ventilated area.

GENERAL INFORMATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Use to control the pests indicated in the areas listed below by application as a 0.25% or .5% chlorpyrifos (Dursban) spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute with water or with a suitable base oil.*

*Use only deodorized kerosene for oil-based sprays to be applied in food handling establishments.

SPRAY DILUTION CHART*

Gallons of Spray Mixture Desired	Fluid Ounces* of DURSBAN 1-E To Use	
	0.25% Spray	0.50% Spray
Water-base spray:		
1	1 1/4	2 1/2
3	4	8
5	6 1/2	13 1/2
10	13	26 1/2

*Oil-Base Spray:

1	1	2
3	3	6
5	5	10
10	10	21

Note: 1 fluid ounce = 2 tablespoonsful = 6 teaspoonful

*Calculated on basis of deodorized kerosene

NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrin may be added to the spray mixture. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays. Consult labeling for the dichlorvos or pyrethrin product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use in combination with dichlorvos (DDVP) or pyrethrin for treatment of food areas of food handling establishments.

CONTINUED ON BACK PANEL

**DIRECTIONS FOR USE,
CONTINUED FROM SIDE PANEL**

**APPLICATION WITHIN FOOD
HANDLING ESTABLISHMENTS**

(Places other than private residences in which food is held, processed, prepared or served), including but not limited to restaurants, grocery stores, bakeries, bottling plants, canneries and grain mills: Use a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 PSI or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated, using special care to avoid unnecessary runoff. Do not introduce spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations. For ants, apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs, bases, or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), DURSBAN may be applied after 2 days from last treatment but emergency use should be limited to once per month. Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

APPLICATION WITH RESIDENTIAL BUILDINGS

Including homes and apartment buildings and within non-food areas of industrial, institution, and commercial buildings including hospitals, stores, manufacturing plants and warehouses: For spot treatment only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only.

Note. A period of 4 to 7 days is normally required for maximum effect on cockroaches.

FOR CONTROL OF CARPET BEETLES, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

FOR CONTROL OF FLEAS AND BROWN DOG TICKS, thoroughly apply a 0.5% spray as a spot treatment of infested areas such as pet beds and resting quarters, near-by cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. **DO NOT TREAT PETS WITH THIS PRODUCT.**

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas, use only oil-base sprays. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

ZOYSIA LAWNS: Chinch Bug damage appears as a yellowing and final browning. The bug sucks the plant juices from the grass blades and eventually plants die. The turf should be free of thatch. Mow to 1 inch height and remove the clippings before spraying. Preventive sprays may be applied starting May 1st, repeated at June 15th and finally on August 1st or as recommended by your local Agricultural Extension Service specialist for your area. If damage occurs in unsprayed areas, treat immediately and plan on a second treatment within 30 to 45 days.

Use Rate & Dilution Table for Ornamental Turf

AREA Sq. Ft.	AMOUNT Chemical	Use Hose-End Sprayer. Dilute with Water to Give
1,000 sq. ft.	3 tbsp.	3-6 gals.
5,000 sq. ft.	15 tbsp.	15-30 gals.
10,500 sq. ft.	1 pint	30-60 gals.

Small 1½ and 3 gallon pressure sprayers are not well adapted for this treatment.

BLUEGRASS LAWNS: Sod Webworms are capable of destroying large areas of turf. New worms appear in late April and early May with successive generations hatching each month through August. For this reason, it is wise to plan on three applications spaced 30 to 45 days apart, starting May 1st. Apply to the turf early or late in the day when worms are active. This rate spray will also control any earwigs, cutworms or grasshoppers present.

Before spraying, turf should be free of troublesome thatch which encourages insects. Rake the area and remove the debris to expose insect hiding places. Water the lawn and cut to a height of 2 to 2½ inches.

PESTS OF ORNAMENTALS: Use DURSBAN insecticide to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed), and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute DURSBAN insecticide with water according to following directions and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants. To control bagworms, grasshoppers, mites, spittlebugs, whiteflies, use 1 tbsp. in 3 gals. water or 1 pint in 100 gals. of water. For cutworms, mealybugs, leafhoppers, thrips (exposed), use 2 tbsp. per 3 gals. or 1 qt. in 100 gals.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

BEST AVAILABLE COPY

corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations. For ants, apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs, bases, or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back) DURSBAN may be applied after 2 days from last treatment but emergency use should be limited to once per month. Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

APPLICATION WITH RESIDENTIAL BUILDINGS

Including homes and apartment buildings and within non-food areas of industrial, institution, and commercial buildings including hospitals, stores, manufacturing plants and warehouses. For spot treatment only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides) ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sink, stoves, refrigerators and cabinets, around plumbing, and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only.

Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

use only oil-base sprays. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

ZOYSIA LAWNS: Chinch Bug damage appears as a yellowing and final browning. The bug sucks the plant juices from the grass blades and eventually plants die. The turf should be free of thatch. Mow to 1 inch height and remove the clippings before spraying. Preventive sprays may be applied starting May 1st, repeated at June 15th and finally on August 1st or as recommended by your local Agricultural Extension Service specialist for your area. If damage occurs in unsprayed areas, treat immediately and plan on a second treatment within 30 to 45 days.

Use Rate & Dilution Table for Ornamental Turf

AREA Sq Ft	AMOUNT Chemical	Use Hand-End Sprayer: Dilute with Water to Give
1,000 sq ft	3 fl oz	34 gal
5,000 sq ft	15 fl oz	153 gal
10,500 sq ft	1 1/2 gal	326 gal

Small 1 1/2 and 3 gallon pressure sprayers are not well adapted for this treatment.

Dilute DURSBAN insecticide with water according to following directions and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants. To control bagworms, grasshoppers, mites, spittlebugs, whiteflies, use 1 tbsp in 3 gals water or 1 pint in 100 gals of water. For cutworms, mealybugs, leafhoppers, thrips (exposed), use 2 tbsp per 3 gals or 1 qt in 100 gals.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

BEST DOCUMENT AVAILABLE

Reg # 10370-37
DURSBAN® 2E
 4/4/91

PM-19

192

INSECTICIDE
—Controls—

Chinch Bugs, Sod Webworms (Lawn Moth), Bagworms,
 Cutworms, Grasshoppers,
INDOOR PEST CONTROL

To be applied only by or under the Supervision of Pest Control Operators and other Trained Personnel responsible for Insect Control Programs.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro 2-pyridyl) phosphorothioate] 24.29%
 Xylene Range Aromatic Solvent ~~67.71%~~

***INERT INGREDIENTS:**

..... ~~28.0%~~ 75.71%
 Total 100.0%

Contains 2 pounds Chlorpyrifos per gallon
 DURSBAN® — Trademark of The Dow Chemical Company

U.S. Patent No. 3,244,586

* CONTAINS XYLENE RANGE AROMATIC SOLVENT

KEEP OUT OF REACH OF CHILDREN
WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **IF ON SKIN:** Wash thoroughly with soap and warm water. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically, atropine is an antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not take internally. Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing. Avoid breathing vapors and spray-mist. Avoid spraying food crops. Handle concentrate in a ventilated area. Keep children and pets off treated area until spray has dried.

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

EPA Est. No. 10370-TX-1

EPA Reg. No. 10370-37 NET CONTENTS: ONE GALLON

Manufactured By

Ford's

Chemical & Service, Inc.

2739 PASADENA BLVD.
 PASADENA, TEXAS 77502

ACCEPTED
 AUG 28 1984
 Un...
 as...
 registered under
 EPA Reg. No. 10370-37

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL AND CHEMICAL HAZARDS

Keep container closed. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Handle concentrate in a ventilated area.

GENERAL INFORMATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Use to control the pests indicated in the areas listed below by application as a 0.25% or .5% chlorpyrifos (Dursban) spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute with water or with a suitable base oil.*

*Use only deodorized kerosene for oil-based sprays to be applied in food handling establishments.

SPRAY DILUTION CHART*

Gallons of Spray Mixture Desired	Fluid Ounces* of DURSBAN 1 E To Use	
	0.25% Spray	0.50% Spray
Water-base spray:		
1	1 1/2	2 1/2
3	4	8
5	6 1/2	13 1/2
10	13	26 1/2
Oil-Base Spray:		
1	1	2
3	3	6
5	5	10
10	10	21

Note: 1 fluid ounce = 2 tablespoonsful = 6 teaspoonful
 *Calculated on basis of deodorized kerosene
 NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

CONTINUED ON BACK PANEL

Not Reviewed. Registrant claims to be in accordance with EPA's Product Guidance for Phase 2...
 to Inert Change in Status.

**DIRECTIONS FOR USE,
CONTINUED FROM SIDE PANEL**

**APPLICATION WITHIN FOOD
HANDLING ESTABLISHMENTS**

(Places other than private residences in which food is held, processed, prepared or served), including but not limited to restaurants, grocery stores, bakeries, bottling plants, canneries and grain mills: Use a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 PSI or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated, using special care to avoid unnecessary runoff. Do not introduce spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations. For ants, apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs, bases, or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), DURSBAN may be applied after 2 days from last treatment but emergency use should be limited to once per month. Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

APPLICATION WITH RESIDENTIAL BUILDINGS

Including homes and apartment buildings and within non-food areas of industrial, institutional, and commercial buildings including hospitals, stores, manufacturing plants and warehouses: For spot treatment only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only.

Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

FOR CONTROL OF CARPET BEETLES, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

FOR CONTROL OF FLEAS AND BROWN DOG TICKS, thoroughly apply a 0.5% spray as a spot treatment of infested areas such as pet beds and resting quarters, near-by cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. **DO NOT TREAT PETS WITH THIS PRODUCT.**

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas, use only oil-base sprays. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

ZOYSIA LAWNS: Chinch Bug damage appears as a yellowing and final browning. The bug sucks the plant juices from the grass blades and eventually plants die. The turf should be free of thatch. Mow to 1 inch height and remove the clippings before spraying. Preventive sprays may be applied starting May 1st, repeated at June 15th and finally on August 1st or as recommended by your local Agricultural Extension Service specialist for your area. If damage occurs in unsprayed areas, treat immediately and plan on a second treatment within 30 to 45 days.

Use Rate & Dilution Table for Ornamental Turf

AREA Sq. Ft.	AMOUNT Chemical	Use Hand-End Sprayer, Dilute with Water to Give
1,000 sq. ft.	3 tbsp.	3-8 gals.
5,000 sq. ft.	15 tbsp.	15-30 gals.
10,500 sq. ft.	1 pint.	30-60 gals.

Small 1 1/2 and 3 gallon pressure sprayers are not well adapted for this treatment.

BLUEGRASS LAWNS: Sod Webworms are capable of destroying large areas of turf. New worms appear in late April and early May with successive generations hatching each month through August. For this reason, it is wise to plan on three applications spaced 30 to 45 days apart, starting May 1st. Apply to the turf early or late in the day when worms are active. This rate spray will also control any earwigs, cutworms or grasshoppers present. Before spraying, turf should be free of troublesome thatch which encourages insects. Rake the turf and remove the debris to expose insect hiding places. Water the lawn and cut to a height of 2 to 2 1/2 inches.

PESTS OF ORNAMENTALS: Use DURSBAN insecticide to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed), and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute DURSBAN insecticide with water according to following directions and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants. To control bagworms, grasshoppers, mites, spittlebugs, whiteflies, use 1 tbsp. in 3 gals. water or 1 pint in 100 gals. of water. For cutworms, mealybugs, leafhoppers, thrips (exposed), use 2 tbsp. per 3 gals. or 1 qt. in 100 gals.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

Not Recommended. Registrant claims to be in accordance with
Revised Edition of Pesticide Product Guidance for Phase 2
Registration, Page 2.12, Active to Inert Change in Status.

DURSBAN® 2E

INSECTICIDE
—Controls—

Chinch Bugs, Sod Webworms (Lawn Moth), Bagworms,
Cutworms, Grasshoppers,
INDOOR PEST CONTROL

To be applied only by or under the Supervision of Pest Control Operators and other Trained Personnel responsible for Insect Control Programs.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 24.29%
Xylene Range Aromatic Solvent ~~82.77%~~

INERT INGREDIENTS:

Total 100.0%

Contains 2 pounds Chlorpyrifos per gallon
DURSBAN® — Trademark of The Dow Chemical Company

U.S. Patent No. 3,244,506

* CONTAINS XYLENE RANGE AROMATIC SOLVENT

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **IF ON SKIN:** Wash thoroughly with soap and warm water. **IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically, atropine is an antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not take internally. Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing. Avoid breathing vapors and spray mist. Avoid spraying food crops. Handle concentrate in a ventilated area. Keep children and pets off treated area until spray has dried.

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

EPA Est. No. 10370-TX-1

EPA Reg. No. 10370-37 NET CONTENTS. ONE GALLON

Manufactured By

Ford's

Chemical & Service, Inc.

2739 PASADENA BLVD.
PASADENA, TEXAS 77502

ACCEPTED
AUG 28 1984
U.S. Environmental Protection Agency
EPA Reg. No. 10370-37

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**PHYSICAL AND
CHEMICAL HAZARDS**

Keep container closed. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Handle concentrate in a ventilated area.

GENERAL INFORMATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Use to control the pests indicated in the areas listed below by application as a 0.25% or .5% chlorpyrifos (Dursban) spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute with water or with a suitable base oil.*

*Use only deodorized kerosene for oil-based sprays to be applied in food handling establishments.

SPRAY DILUTION CHART*

Gallons of Spray Mixture Desired	Fluid Ounces* of DURSBAN 2E To Use	
	0.25% Spray	0.50% Spray
Water base spray		
1	1 1/2	2 1/2
3	4	8
5	6 1/2	13 1/2
10	13	26 1/2
*Oil Base Spray:		
1	1	2
3	3	6
5	5	10
10	10	20

Note: 1 fluid ounce = 2 tablespoonsful = 6 teaspoonsful

*Calculated on basis of deodorized kerosene

NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

CONTINUED ON BACK PANEL

BEST AVAILABLE COPY

**DIRECTIONS FOR USE,
CONTINUED FROM SIDE PANEL**

**APPLICATION WITHIN FOOD
HANDLING ESTABLISHMENTS**

(Places other than private residences in which food is held, processed, prepared or served), including but not limited to restaurants, grocery stores, bakeries, bottling plants, canneries and grain mills: Use a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 PSI or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated, using special care to avoid unnecessary runoff. Do not introduce spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations. For ants, apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs, bases, or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a plume of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), DURSBAN may be applied after 2 days from last treatment but emergency use should be limited to once per month. Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

APPLICATION WITH RESIDENTIAL BUILDINGS

Including homes and apartment buildings and within non-food areas of industrial, institution, and commercial buildings including hospitals, stores, manufacturing plants and warehouses. For spot treatment only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets, arc and plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only.

Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

FOR CONTROL OF CARPET BEETLES, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

FOR CONTROL OF FLEAS AND BROWN DOG TICKS, thoroughly apply a 0.5% spray as a spot treatment of infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. **DO NOT TREAT PETS WITH THIS PRODUCT.**

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas, use only oil-base sprays. Prior to application, de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

ZOYSIA LAWNS: Chinch Bug damage appears as a yellowing and final browning. The bug sucks the plant juices from the grass blades and eventually plants die. The turf should be free of thatch. Mow to 1 inch height and remove the clippings before spraying. Preventive sprays may be applied starting May 1st, repeated at June 15th and finally on August 1st or as recommended by your local Agricultural Extension Service specialist for your area. If damage occurs in unsprayed areas, treat immediately and plan on a second treatment within 30 to 45 days.

Use Rate & Dilution Table for Ornamental Turf

AREA Sq. Ft.	AMOUNT Chemical	Use Nozzle End Sprayer, Dilute with Water to 600
1 000 sq. ft.	3 tbsp.	3.0 gals.
5 000 sq. ft.	15 tbsp.	15.0 gals.
10 000 sq. ft.	1 pint	30.0 gals.

Small 1 1/2 and 3 gallon pressure sprayers are not well adapted for this treatment.

BLUEGRASS LAWNS: Sod Webworms are capable of destroying large areas of turf. New worms appear in late April and early May with successive generations hatching each month through August. For this reason, it is wise to plan on three applications spaced 30 to 45 days apart, starting May 1st. Apply to the turf early or late in the day when worms are active. This rate spray will also control any earwigs, cutworms or grasshoppers present.

Before spraying, turf should be free of troublesome thatch which encourages insects. Rake the area and remove the debris to expose insect hiding places. Water the lawn and cut to a height of 2 to 2 1/2 inches.

PESTS OF ORNAMENTALS: Use DURSBAN insecticide to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed), and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute DURSBAN insecticide with water according to following directions and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. **NOTE:** Do not use on azaleas, camellias, poinsettias, rose bushes or variegated hvy because of possible injury to these plants. To control bagworms, grasshoppers, mites, spittlebugs, whiteflies, use 1 tbsp. in 3 gals. water or 1 pint in 100 gals. of water. For cutworms, mealybugs, leafhoppers, thrips (exposed), use 2 tbsp. per 3 gals. or 1 qt. in 100 gals.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

BEST AVAILABLE COPY

292

Not Reviewed. Reprinted from the original...