

Ford's

Diazinon 5 Granules

ACCEPTED
NOV - 2 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10370-44

Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and in the home garden only.

Ready-to-use Granules. Contains 5% diazinon insecticide.

Treats up to 5000 square feet.

Active Ingredient:
Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate... 5%

Inert Ingredients: 95%

Total: 100%

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements on back of bag.

EPA Reg. No. 10370-44

kills:

Ants, Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers, Chinch bugs
Crickets, Cutworms
Earwigs, Fire ants*
Fleas, Onion maggots
Root maggots
Sod webworms
White grubs of:
• European chafer
• Japanese beetle
• Southern chafer
Wireworms
...and others

* Aids in control

See Directions for Use

Statement of Practical Treatment

If swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration, if indicated.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

For lawns, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes.

Net Weight

ROUSSEL UCLAF

Roussel Uclaf Corporation
95 Chestnut Ridge Road
Morrisville, New Jersey 07645

BEST AVAILABLE COPY

Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

In Lawns kills

Ants	Crickets	Sod webworms
Armyworms	Cutworms	Sowbugs
Bermudagrass mites	Digger wasps	(Pillbugs)
Billbugs	Earwigs	Springtails
Brown dog ticks	Fire ants*	White grubs of:
Chiggers	Fleas	• European chafer
Chinch bugs	Lawn moths	• Japanese beetle
Clover mites	Leafhoppers	• Southern chafer
	Millipedes	* Aids in control

In Gardens kills

Cutworms	Onion maggots	Wireworms
Mole crickets	Root maggots	

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

OVERALL LAWN APPLICATION

Special Precautions To Reduce Risk To Birds

Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

Diazinon 5 Granules controls certain lawn insects on both grass and dichondra lawns. Apply Granules with fertilizer spreader. Refer to the tables that follow for use rates and spreader settings.

Pest	Rate*	Timing/Special Directions
Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (Pillbugs), Springtails (Collembola)	2 lbs. per 1,000 sq. ft.	Apply when grass is dry. Water lawn lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.
Billbugs, White grubs of Japanese beetle, European chafer, Southern chafer, <i>Atraxius spretulus</i> (dung beetle)	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or Summer when damage is first noticed. Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat again in late Summer or early Fall if a second generation occurs.
Hyperodes weevils	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

* 1,000 sq. ft. = an area 20 ft. x 50 ft.

100 sq. ft. = an area 10 ft. x 10 ft.

2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to achieve correct rate of application.

LAWN SPREADER SETTING NO. **
(To deliver 2 lbs. per 1,000 sq. ft.)

Spreader Make and Model	Setting
Central Imperial Mark No. 1662-G	6½
Central Suburbia Lawn Spreader No. 1622-0	6½
Cyclone Model 81 Rotary	4½
Greenfield Model 105	6½
ORTHO Drop Spreader Model 3000	4½
ORTHO Broadcast Spreader Model 4000	4½
J. C. Penney Model 6005	7¼
J. C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 671.1919P	6

** For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

Lawn Spot Treatment Diazinon 5 Granules

Ants: Apply 1½ teaspoons of **A** over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply ½ cup over, and 2 ft. around, each fire ant mound. Immediately after application, water-in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gal. of water should be applied to each mound area. Reapply as necessary.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear. **Diazinon 5 Granules**

Digger wasps: Apply 1½ teaspoons of **A** in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation

The following pests are commonly found around or near house foundations, and may enter the house:

Ants	Crickets	Millipedes
Brown dog ticks	Earwigs	Sowbugs (Pillbugs)
Clover mites	Fleas	Springtails (Collembola)

Apply a 2-foot band of Granules next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. If necessary, repeat application at 3-week intervals.

VEGETABLES - INSECT CONTROL

Special Precautions To Reduce Risk To Birds

Birds feeding on treated areas may be killed. To reduce risk to birds:

1. Thoroughly mix Granules into the soil during or immediately after application. Refer to the Soil Incorporation table that follows.
2. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

Diazinon 5 Granules controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Do not apply this product to food crops grown in greenhouses.

Unless directions under Timing/Special Directions state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

SOIL INCORPORATION

Insect	Depth of Mixing or Incorporation into Soil
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-8 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

* If type of cutworm cannot be identified, mix into soil 4-8 inches deep.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole and Snap) (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	

Note: Do not use on dried bean varieties, such as pintos, dried limas, or navy beans.

Beets, Red (Table)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Broccoli	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Brussels Sprouts	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

BEST AVAILABLE COPY

Crop	Pest	Rate	Timing/Special Directions
Cabbage	Root Maggots	7-10 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Carrots	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cauliflower	Root Maggots	7-10 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Celery	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
Collards	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cucumbers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
Endive (Escarole)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Kale	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 oz. per 500 sq. ft.	
	Mole Crickets	3 1/2 oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

Lettuce	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, and Hybrids of these), and Watermelons	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Mustard	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (Bulb and Green)	Onion Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Parsley	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Peas (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Note: Do not use on dried pea varieties such as dried black-eyed peas.			
Peppers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Potatoes	Cutworms (Surface and Subterranean)	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Radishes	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Squash (Summer and Winter)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	

5 7 5

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Seed Corn Maggots	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil 2 inches just before planting.
Sweet Potatoes	Wireworms, Flea Beetles	10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting
Swiss Chard	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Tomatoes	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Turnips	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

BEST AVAILABLE COPY