

10370-49

PM 12

103

Mr. R.B. Boyd, Jr.  
Ford's Chemical & Service, Inc.  
2739 Pasadena Boulevard  
Pasadena, TX 77502

OCT 10 1986

Dear Mr. Boyd:

Subject: Add Insects to Perimeter Treatment Directions  
Ford's Roach Bait  
EPA Registration No. 10370-49  
Your Application Dated June 9, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

The reason that we requested that cockroaches be deleted from the perimeter treatment directions is because previously submitted data on cockroaches have indicated very poor performance in reducing infestations with band or perimeter treatments.

Sincerely yours,

Lawrence J. Schnaubelt <sup>LJ</sup>  
Acting Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosure

BEST AVAILABLE COPY

85297:Schnaubelt:E-8:KENCO:8/19/86:8/28/86:de:vo

CONCURRENCES


1320-1 (4-81)

OFFICIAL FILE

7's Date ..06/09/86 REVISED LABELING IN RESPONSE TO 2/5/86 EPA LETTER.

### NATURE OF AMENDMENT

EXPAND THE FOLLOWING APPROVED COPY TO READ AS THE COPY BELOW UNDER NO.1

For Millipedes and Pillbugs scatter the bait in the foraging area around the foundation of buildings. Repeat treatment as needed.

1. ....

#### OUTDOOR INSECT CONTROL AROUND HOMES AND BUILDING FOUNDATIONS

Scatter the bait evenly in a band 5 feet wide around the structure next to the foundation as a Bait Barrier against insects invading the dwelling or place of business. Use 7½ ounces of product to treat a band 5 ft. wide and 20 ft. long (100 sq.ft.) One pound will treat up to 200 Sq.Ft.; (5 ft.x 40 Ft. band) which is equivalent to the rate of 4.6 lbs. per 1,000 Sq.Ft.

This treatment will aid in the reduction of, reduce annoyance from, & help to prevent entry into homes from commonly found insects which may feed upon the bait such as Millipedes, Earwigs, Pillbugs(Sowbugs), Cutworms, Mole Crickets, and Field Crickets,

Apply when insects are present and are causing problems, especially in the late spring or early fall. Repeat as necessary to keep fresh bait in the area or every 30 days when insects are a problem.

2. ... REVISED STORAGE & DISPOSAL STATEMENT- DOES NOT REQUIRE AGENCY APPROVAL.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in it's original labeled container in a cool, dry, locked place out of reach of children. DISPOSAL: Do not re-use empty bag or canister. Wrap and put in trash.

OCT 10 1986

10370-49

BEST AVAILABLE COPY

# ROACH BAIT

A ready-to-use bait insecticide to aid in the control of roaches in homes, apartment buildings, stores, restaurants (non-food areas), hospitals, hotels and motels.

## ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-dimethyl-O-(3,5,6-trichloro-2-pyridyl) phosphorothioate)

0.50%

## INERT INGREDIENTS

99.50%

TOTAL

100.00%

## KEEP OUT OF REACH OF CHILDREN CAUTION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For the control of American, Oriental, Brown-Banded, Smoky Brown, & German cockroaches inside buildings use approximately 2 ounces (1/4 cup measure) of Roach Bait per 100 square feet of space. Place this bait in cracks and crevices or in suitable containers behind and underneath fixtures, in closets, and other areas where roaches may hide. Place only in or near infested areas. All bait placements should be inaccessible to children or pets. Replenish bait as necessary.

For control of roaches in attics use approximately 2 ounces (1/4 cup measure) of Roach Bait per 100 square feet. Scatter bait in inaccessible areas of the attic.

For the control of American, Oriental, Brown Banded, Smoky Brown, & German cockroaches in Food Processing Plants, use only when plant is not in operation and food is not exposed. Use only in suitable bait containers. Locate these bait containers on the floor near walls, in storage areas, under appliances, and other areas where roaches have been observed. Maintain strict account of location and number of bait containers. Remove all bait containers from edible product areas before resuming operations. Do not use in edible product areas of food processing plants, restaurants or other serving areas where food is commercially prepared or processed while plant is in operation. Do not use in serving areas while food is exposed.

Roach Bait are best used as a supplemental residual treatment preceded by recommended roach sprays and good sanitation practices.

For Harvester Ants, Black Crazy Ant, Little Black Ant, Pyramid Ant, Pharaoh Ant, Cornfield Ant, Thief Ant, and Certain Other Ants scatter 1 to 2 tablespoons of bait within 6 to 12 inches of the nest entrance. Additional bait may be scattered in the foraging area and around the foundations of the buildings. Control of the colony may take 2 to 6 weeks. Repeat treatment after 6 weeks if necessary. Treatment should be made only when ants are active.

For Millipedes and Pillbugs scatter the bait in the foraging area around the foundation of buildings. Repeat treatment as needed.

EPA Reg. No. 10370-49

### STORAGE AND DISPOSAL

Do not re-use empty container. Wrap container and put in trash collected.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep out of reach of children. May be harmful if swallowed. Do not take internally. Avoid eye and skin contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

**IF ON SKIN:** In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water.

**IF ON EYES:** Flush with plenty of water.

**NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

**NOTICE:** Seller guarantees that he complies with the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of the use or handling and accepts the product on these conditions.

EPA Est. No. 10370-1a 1

MANUFACTURED BY

**ford's**

CHEMICAL & SERVICE, INC.

2739 PASADENA BLVD.  
PASADENA, TEXAS 77502

**BEST AVAILABLE COPY**