

ford's

CONTINUED FROM BACK PANEL

Cutworms, Lespedeza — Cutworms (Surface and Subterranean) — Broadcast 2.4 lbs. per acre... Peanuts — Cutworms apply 2.0 lbs. per acre... Soybeans — Cutworms (Surface and Subterranean) — Broadcast 2.4 lbs. per acre... Sugar Beets — Cutworms broadcast 3.4 lbs. per acre... Tobacco — Cutworms broadcast 2.3 lbs. per acre...

RANGE, PASTURE AND GRASSLAND INSECTS — Unless otherwise stated, Diazinon AG500 should always be applied as a spray... Bermudagrass and Forage Grasses (Other than Range and Pasture Grasses) — Insects: Wintered Grasshoppers apply 1.0 lb. per acre... Ornamental Insect Control — To control certain insects on Azalea, Azalea, Birch, Boxwood, Camellia, Gardenia, Chrysanthemum, Dogwood, Elm, Gardenia, Hawthorn, Holly, Juniper, Lead, Locust, Magnolia, Pine, Plum, Poinsettia, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated below.

Table with 3 columns: Insect Name, Rate per 300 sq. ft., and Rate per 1000 sq. ft. Lists insects like Japanese Beetles, European Spruce Sawflies, etc.

Do not use on certain Ferns, Fuchsia, Hibiscus, Papaya, Pilea and Gardenia since injury to the plants may occur... Do not use on certain Ferns, Fuchsia, Hibiscus, Papaya, Pilea and Gardenia since injury to the plants may occur.

LAWN INSECT CONTROL — To control insects listed below, apply recommended rates for control of Ants, Beetles, Caterpillars, Crickets, Earwigs, Fleas, Mosquitoes, Sawflies and Coleoptera (Springtails)...

Table with 4 columns: Insect Name, Rate per 1000 sq. ft. (Diazinon AG500), Rate per 1000 sq. ft. (Water), and Remarks. Lists insects like Leafhoppers, Fleas, etc.

Do not use on certain plants... Do not use on certain plants...

FOREST INSECTS — Pine, Seed Chewing Bark Beetles, Bark Beetles, Bark Beetles, Bark Beetles... FIRE ANTS — To control fire ants, apply Diazinon AG500 as a spray...

DIAZINON® AG500 INSECTICIDE

FOR AGRICULTURAL USE ONLY FOR THE CONTROL OF CERTAIN INSECTS ON FRUITS, NUTS, VEGETABLES, FIELD CROPS AND ACTIVE INGREDIENTS:

- 0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate
Xylene
INERT INGREDIENTS

DIAZINON® Registered Trademark of Ciba-Geigy Corp. DIAZINON AG500 contains 4 lbs. of D

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center immediately. Gastric lavage is indicated if material was taken... IF INHALED: Remove victim to fresh air. Apply respiration if indicated... IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water... IF IN EYES: Flush eyes for at least 15 minutes with soap and water.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing after handling...

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding on treated areas may be killed... This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area.

PHYSICAL AND CHEMICAL HAZARDS FLAMMABLE MIXTURE

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. PESTICIDE DISPOSAL: Pour or rinse that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides...

EPA Reg. No. 10370-52 NET CONTENTS: ONE GALLON MANUFACTURED BY

ford's Chemical & Service, Inc. 2739 PASADENA BLVD. PASADENA, TEXAS 77502

ACCEPTED  
COMMENTS  
EPA Letter Dated:

NOV 29 1982

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

500

10370-52

FIELD CROPS AND ORNAMENTALS

48.0%  
31.83%  
20.17%  
TOTAL 100.00%

contains 4 lbs. of Diazinon per gallon

DREN

NT  
material was taken internally DO NOT IN-  
duce vomiting. Do not induce vomiting or give anything

persists  
red in conjunction with atropine sulfate

MALS

es on skin or on clothing Wash thoroughly  
or pets Do not permit children or pets to  
contaminate feed or use on household pets

may be killed Keep out of lakes streams  
to water that will be used for recreational  
recommended on this label Do not apply  
used in areas where impact on threatened  
Do not apply when weather conditions  
Apply this product only as specified

ed area Do not apply when bees are actively  
to provide the maximum possible interval

open flame

ICIDE DISPOSAL: Pesticide spray mixture  
pesticides or buried in a safe place away  
reconditioning, or puncture and dispose

EPA Est. No. 10370-TX-1

6-82

DIRECTIONS FOR USE

GENERAL INFORMATION

NOTE: Unless otherwise stated, the rates listed on this label should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

**FRUIT AND NUT CROPS** — NOTE: Unless otherwise stated, the rates listed on this label should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

**Almonds** — Mites: Two Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, etc. Apply as a dormant spray in early spring when trees are still on the ground and buds are closed. Make a second application when the buds are well developed but still closed. Do not apply within 7 days of harvest. Raspberry, Crown Borer (Raspberry Root Borer), apply 2 lbs. per acre in a minimum of 100 gallons of water as a trench to the crown area and to the lower limbs. Apply either in the spring before berries begin to form, or in the fall after harvest. **Cherries** — European Red Mites, Fruit Tree Leaf Rollers, mix 1 qt. per 100 gal. of water. Apply in cover sprays as necessary, beginning with petal fall. Leafhoppers, apply 1 qt. per 100 gal. of water in 3-4 applications beginning at petal fall and continuing through the growing season or apply as infestations develop. Cherry Fruit Flies, mix 1 qt. per 100 gal. of water. Make 3 or 4 applications beginning after adult females begin to emerge and repeat at 10-day intervals. Bark Cherry Aphids, mix 1 qt. per 100 gal. of water. Apply as a cover spray when aphids appear and repeat after 10 days. San Jose Scale Crawlers, mix 1 qt. per 100 gal. of water. Apply during the pre-bloom stage of when crawlers first appear. Do not apply within 10 days of harvest. Cherry Rust Mites, mix 1 qt. per 100 gal. of water. Apply as a dormant spray after harvest. **Citrus Fruits** — Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangerines, Citrus Citrus Kumquats and hybrids of these — Citrus Aphids, Fruit Tree Leaf Rollers, etc. Apply as a dormant spray after harvest. **Cranberries** — B and B Headed Fleaworms, Spangarmite, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. Cranberry Fruitworms, apply 3 lbs. in 40 gal. of water per acre. Do not apply within 7 days of harvest. **Figs** — Fig Aphid, Fig Wasp, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Ferrets** — Ferret Leaf Miner, apply 2 lbs. in 40 gal. of water per acre. **Grapes** — Grape Aphid, Grape Leafhopper, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Olives** — Olive Scale Crawlers, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Peaches and Nectarines** — Aphids, Brown Mites, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Pineapples** — Scale, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Plums and Prunes** — Brown Mites, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Strawberries** — Aphids, etc. Apply 2 lbs. per acre in 40 gal. of water per acre. **Walnuts** — Aphids, etc. Apply 2 lbs. per acre in 40 gal. of water per acre.

CONTINUED FROM SIDE PANEL

**VEGETABLE FOLIAR INSECTS** — Unless otherwise stated, foliar applications of Diazinon AG-500 should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

**VEGETABLE SOIL INSECTS** — Unless otherwise stated, soil applications of Diazinon AG-500 should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

**FIELD AND FORAGE FOLIAR INSECTS** — Unless otherwise stated, foliar applications of Diazinon AG-500 should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

**FIELD AND FORAGE SOIL INSECTS** — Unless otherwise stated, soil applications of Diazinon AG-500 should be applied as a minimum. For more information on the use of this pesticide, consult the label for the specific crop. The following information is provided for your information only. It is not intended to replace the label for any specific crop. The user should always refer to the label for the specific crop. The user should always refer to the label for the specific crop.

Row Spacing	Linear foot of Corn row/acre	Rate per acre	Planting Time application	Pestemergence basal treatment
30"	10,800	25.56 lb.		14.28 lb.
40"	8,100	34.28 lb.		19.05 lb.
48"	6,750	41.14 lb.	1.75 lbs.	22.86 lb.
60"	5,400	51.43 lb.	2.25 lbs.	28.57 lb.
72"	4,500	62.86 lb.	2.85 lbs.	34.29 lb.
84"	3,600	79.71 lb.	3.65 lbs.	42.86 lb.
96"	2,700	104.76 lb.	4.65 lbs.	56.43 lb.
108"	1,800	139.64 lb.	6.05 lbs.	75.14 lb.
120"	1,350	185.71 lb.	7.95 lbs.	99.99 lb.

NOTE: For spacings not listed, calculate the appropriate dosage based on the 40" row figure

CONTINUED ON BACK PANEL



10370-53

- 2 -

d. Revise the Pesticide Disposal text to read:

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

e. Correct the following typographical errors:

- (1) Revise "windlife" to "wildlife" in the second sentence of the Environmental Hazards.
- (2) Revise "Cranberries, Blackberries, etc." to "Caneberries--Blackberries, etc." in the Fruit and Nut Crops directions.
- (3) Revise "mushroom u~~s~~" to "mushrooms houses" in the Mushroom directions of the Vegetable Foliar Insects section.
- (4) Revise "broadhast" to "broadcast" in the cowpeas directions of the Field and Forage Soil Insects section.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosure