

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

APR 1 3 1994

Christina M. Lawes ROUSSEL UCLAF CORP. 95 Chestnut Ridge Road Montvale, NJ 07645416

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

SUBJECT:

Label Amendment Submissions of 11/08/93 in Response to PR Notice 93-7

EPA Reg. No.: SEE ATTACHED /0370-54

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully REMOVES THE PRODUCT FROM THE SCOPE of PR Notice 93-7 one the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements), the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

Sincerely,

James Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Enclosures

BEST COPY AVAILABLE



Printed with Soy/Canola ink on paper that contains at least 50% recycled liber

FOUSSEL UCLAF CORP. 10370 11/08/93

Original Submissions

FORD'S LIQUID SEVIN 2L BRAND CARBARYL INSECTICIDE - 10370-32
FORD'S DURSBAN 2.5% G - 10370-54
FORD'S 50% MALATHION EC - 10370-147
FORD'S CRABGRASS AND FOXTAIL KILLER - 10370-169
LAWN FUNGICIDE GRANULES - 10370-202
ORGANICIDE DIP - 10370-207
CLEAN CROP SEVIN BRAND CARBARYL INSECTICIDE LIQUID - 10370-285

DURSBAN 2.5% G

* CONTROLS CHINCH BUGS, SOD WEBWORMS (LAWN MOTHS),
ANTS, FLEAS, COCKROACHES, BROWN DOG TICKS, CRICKETS,
CERTAIN WHITE GRUBS, **CUTWORMS, EARWIGS, SOWBUGS
(PILBUGS), AND HYPERODES (TURFGRASS) WEEVILS ***INFESTING HOME LAWNS AND OTHER ORNAMENTAL TURF
AREAS

"Dursban is a registered trademark of the Dow Elanco. Contains Dursban Brand Insecticide. U.S. Patent No. 3,244,586. Gold Crest is a registered trademark of Roussel Uclaf Corporation.

EPA REG. NO. 10370-54

EPA EST. NO.

KEEP OUT OF THE REACH OF CHILDREN -- CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and wash affected areas with soap and water.

IF ON EYES: Flush eyes with plenty of water.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

See Side Panel For Additional Precautionary Statements

ACCEPTED NET CONTENTS with COMMENTS In EPA Letter Dated

JUN 30 1994

Under the Federal Insecticide. Fundicide, and Redenticide Act no amended. for the pesticide registered under EPA Reg. No.



ROUSSEL UCLAF

Roussel Uclaf Corporation 95 Chestnut Ridge Road Montvale, New Jersey 07645

f54s 050493tk

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

2 ...

MAY BE HARMFUL IF SWALLOWED. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. This product is toxic to fish and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water, and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not re-use empty container. Wrap container and put in trash collection.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply Dursban 2.5%G uniformly to home lawns and other omamental and recreational turf areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear, using Dursban 2.5%G at the rate of 40 pounds per acre. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of Dursban 2.5%G and water needed for small sized areas. Re-treat as necessary.

RATES TO USE

Areas to be Treated	Pounds of Dursban 2.5%G	Gals. of Water to Sprinkle on Turf after Treatment
1,000 sq. ft.	1	30
5,000 sq. ft.	5	150
1,000 sq. ft. 5,000 sq. ft. 10,700 sq. ft.	10	325

FLEAS AND BROWN DOG TICKS Thoroughly apply Dursban 2.5%G as a spot treatment to infested areas such as:

PET BEDS: Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment of pet areas. Do not apply this product indoors. To control the source of your flea infestation, treat pets with an E.P.A. approved product registered for direct application to pets.

YARD AREAS: Apply Dursban 2.5%G uniformly to home lawns, under shrubbery, and other ornamental recreational turf grass areas. Treat when insects first appear using granules at the rate of one pound per 1,000 sq. ft. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application, sprinkle the turf with 50 gallons of water per 1,000 sq. ft. so as to wash the granules into the zone of insect infestation. Re-treat as necessary. DO NOT TREAT PETS WITH THIS PRODUCT.

COCKROACHES: To help prevent infestation of buildings by cockroaches. Treat a band 6 to 10 feet wide around and adjacent to the building where pets are active and may find entrance. Treat at a rate of one pound per 1,000 square feet.

**TO CONTROL CERTAIN WHITE GRUBS (larvae of European chafer and Japanese beetle) use two to four times the amounts of Dursban 2.5%G shown in the table. (i.e., 2-4 lbs. per 1,000 sq.ft.). After application, apply enough water to thoroughly wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

***TO CONTROL HYPERODES (TURFGRASS) WEEVILS, use twice the rate of Dursban 2.5%G shown in the table. Treat suspected problem areas in Mid-April and again in Mid-May. consult your local Agricultural Extension Service specialist for the best time to treat in your area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.