

Ford's

DURSBAN 2.5% G

GRANULAR INSECTICIDE

* **CONTROLS CHINCH BUGS, SOIL WEBWORMS (LAWN MOTHS), ANTS, FLEAS, COCKROACHES, BROWN DOG TICKS, CRICKETS, CERTAIN WHITE GRUBS, CUTWORMS, EARWIGS, SOWBUGS (PILBUGS), AND HYPERODES (TURFGRASS) WEEVILS INFESTING HOME LAWNS AND OTHER ORNAMENTAL TURF AREAS**

* **Kills Fire Ants**

ACTIVE INGREDIENT:

Chlorpyrifos(0,0-diethyl-0-(3,5,6-trichloro-2-pyridyl)phosphorothioate) 2.5%

INERT INGREDIENTS: 97.5%
100.0%

*Dursban is a registered trademark of the Dow Elanco.
Contains Dursban Brand Insecticide.
U.S. Patent No. 3,244,586.

EPA REG. NO. 10370-54

EPA EST. NO.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and wash affected areas with soap and water.

IF ON EYES: Flush eyes with plenty of water.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

**See Side Panel For Additional
Precautionary Statements**

ACCEPTED
SEP 27 1994
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No.

NET CONTENTS

ROUSSEL UCLAF

Roussel Uclaf Corporation
95 Chestnut Ridge Road
Montvale, New Jersey 07645
154s 121493xk



PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION

MAY BE HARMFUL IF SWALLOWED. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water, and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water.

Do not apply where runoff is likely to occur.

Do not contaminate water by cleaning of equipment or disposal of equipment washwater.
Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not re-use empty container. Wrap container and put in trash collection.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Apply Dursban 2.5%G uniformly to home lawns and other ornamental and recreational turf areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear, using Dursban 2.5%G at the rate of 40 pounds per acre. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle.

For best results sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of Dursban 2.5%G and water needed for small sized areas. Re-treat as necessary.

RATES TO USE

Areas to be Treated	Pounds of Dursban 2.5%G	Gals. of Water to Sprinkle on Turf after Treatment
1,000 sq. ft.	1	30
5,000 sq. ft.	5	150
10,700 sq. ft.	10	325

FLEAS AND BROWN DOG TICKS Thoroughly apply Dursban 2.5%G as a spot treatment to infested areas such as:

PET BEDS: Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment of pet areas. Do not apply this product indoors. To control the source of your flea infestation, treat pets with a product registered for direct application to pets.

BEST AVAILABLE COPY

YARD AREAS: Apply Dursban 2.5%G uniformly to home lawns, under shrubbery, and other ornamental recreational turf grass areas. Treat when insects first appear using granules at the rate of one pound per 1,000 sq. ft. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. After application, sprinkle the turf with 50 gallons of water per 1,000 sq. ft. so as to wash the granules into the zone of insect infestation. Re-treat as necessary. **DO NOT TREAT PETS WITH THIS PRODUCT.**

COCKROACHES: To help prevent infestation of buildings by cockroaches. Treat a band 6 to 10 feet wide around and adjacent to the building where pets are active and may find entrance. Treat at a rate of one pound per 1,000 square feet.

TO CONTROL CERTAIN WHITE GRUBS (larvae of European chafer and Japanese beetle) use two to four times the amounts of Dursban 2.5%G shown in the table. (i.e., 2-4 lbs. per 1,000 sq.ft.). After application, apply enough water to thoroughly wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

TO CONTROL HYPERODES (TURFGRASS) WEEVILS, use twice the rate of Dursban 2.5%G shown in the table. Treat suspected problem areas in Mid-April and again in Mid-May. consult your local Agricultural Extension Service specialist for the best time to treat in your area.

← Add Attachment ①

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

BEST AVAILABLE COPY

ATTACHMENT 1

IMPORTED FIRE ANT TREATMENT: As an aid in the control of the *imported fire ant* apply 4 tablespoons of this product to the center of the mound and an additional 2 tablespoons to the two feet surrounding it. Do not water in or disturb the mound. Apply when temperatures are between 65° and 80°F. Allow 3 to 4 days for maximum control, then repeat if necessary.

ANTS NESTING OUTDOORS: This product may be used outdoors for control of ants, including; *Acrobat, Argentine, Big Head, Cornfield, Field, Fire, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants*. If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, slabs or in stumps or logs. Dursban 2.5% G should be applied to thoroughly cover the nest and irrigate to properly water in the product. In addition to treating ant nests or mounds, entry points of buildings (used by ants) and the soil surface along the perimeter of the building should be treated.

When ants are scattered or numerous in a lawn or garden, or where colonies are very large (especially with the Argentine or fire ants), a broadcast treatment may be necessary. Apply granules to thoroughly cover the infested areas, paying particular attention to the areas around trees, bushes and fences, and irrigate with the appropriate amount of water. Apply at a rate of 1 2/3 pounds per 1000 square feet.

BEST AVAILABLE COPY