

ford's

CONTINUED FROM BACK PANEL

materials. The use of large quantities in any area may leave an objectionable odor. Care should be taken to avoid deposits which could frequently be contacted by children. Do not allow children in treated areas until surfaces are dry. Do not contaminate food, food containers or cooking utensils.

ANIMALS

BEEF CATTLE: Lice — 1 gal. to 100 gals. of water or 6 1/2 oz. per 5 gals. of water. Ticks — 1 1/2 gals. per 100 gals. of water or 6 1/2 oz. per 5 gals. of water. Horn Fly — 1 1/4 gals. per 100 gals. of water or 6 1/2 oz. per 5 gals. Apply complete coverage spray. One treatment may be sufficient. Repeat application only as needed at 2 week intervals.

DAIRY CATTLE (Non Lactating): Lice — 6 1/2 oz. per 5 gals. of water. Ticks — 6 1/2 oz. per 5 gals. of water. Repeat applications only as needed at 2 week intervals. DO NOT spray during lactation or within two weeks of freshening. DO NOT apply to calves under one month of age. DO NOT apply to lactating cattle. DO NOT dip within two months of freshening.

SWINE: Lice — 1 gal. to 100 gals. of water or 6 1/2 oz. per 5 gals. of water. Apply as a complete spray to animals pens and litter. One treatment may be sufficient. Repeat application only as needed. When applying spray, avoid contamination of feed, food containers and watering troughs. Sprayed hogs should be kept out of sun and wind for a few hours after treatment.

DOGS AND CATS — DIRECT APPLICATION

For control of fleas, apply spray of 1 oz. of Malathion 57% Emulsifiable Liquid per 1 gal. of water. Wet the animal thoroughly as a hand sprayer. If the animal has free run treatment may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas before retreating. If the animal is confined to an area free of fleas, one treatment may be sufficient. DO NOT treat animals under one month of age.

ANIMAL QUARTERS

For control of fleas in kennels, pens, yards, lawns and under houses. Remove manure or debris. Use spray of 5 oz. of 57% Emulsifiable Liquid per 1 gal. of water at rate of 1 gal. per 1000 sq. ft. Use garden sprayer.

ORNAMENTALS

For Oyster Shell Scale and Lace Bugs, use 1 pint. For Spider Mites, Aphids, White Flies, Mealybugs, Thrips, Japanese Beetle, European Pine Shoot Moth, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scrub Scale and Euonymus Scale, use 1 1/2 pints. For Bagworm, Tent Caterpillar, Boxwood Leaf Miner, Oak Barkworm, Florida Red Scale, Fletcher Scale, Juniper, Azalea and Magnolia Scale, use 2 pints. For Black Scale, Citrus, Salt Brown and Monterey Scales, use 2 1/2 pints. Apply thorough full coverage sprays as needed. Note: DO NOT apply to petunias, certain ferns (including Boston Maidenhair, Pteris) and some species of Grasses since plant injury may occur.

At above rates per 100 gals. of water

NOTICE: For complete information on the use of Malathion 57% Emulsifiable Liquid, see the label on the container.

MALATHION 57%

PREMIUM GRADE
An Emulsifiable Concentrate containing
5 lbs. of Actual Malathion per Gallon

ACTIVE INGREDIENTS

Malathion (0,0 dimethyl dithiophosphate of diethyl mercaptosuccinate) 57.00%

Xylene 29.27%

INERT INGREDIENTS

TOTAL 100.00%

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR OTHER CAUTIONS

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or poison treatment center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one of two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. If on skin — Wash with soap and water. If in eyes — Flush with water for 15 minutes. Contact a physician. If inhaled — Remove victim to fresh air and apply respiration if indicated.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

EPA Reg. No. 10370 XX NET CONTENTS: ONE GALLON EPA Est. No. 10370 TX 1

MANUFACTURED BY

ford's

Chemical & Service, Inc.

2739 PASADENA BLVD. PASADENA, TEXAS 77502
(713) 477-5525

283

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION

May be fatal if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing vapors or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding on treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL AND CHEMICAL HAZARDS

FLAMMABLE MIXTURE Do not use, pour, spill, or store near heat or open flame.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

Pesticide Disposal: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture, and dispose of in a sanitary landfill or by other approved state and local procedures.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CRACK AND CREVICE TREATMENT

FOOD AND FEED HANDLING ESTABLISHMENTS — Places other than private residences in which exposed food or feed is held, processed, prepared or served.

FOOD AREA—APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems such as milks, dairies, bakeries, restaurants or anywhere food or feed is stored, processed, prepared or served. Serving areas (when food is exposed and facility is in operation) also would be considered a food area.

Apply in small amounts directly into cracks and crevices.

CONTINUED FROM SIDE PANEL

CONTINUED FROM SIDE PANEL

using a bulbous duster or other suitable equipment capable of applying insecticide directly into cracks and crevices, in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases, conduits, motor housing and electrical switch boxes where cockroaches hide.

Care should be taken to avoid depositing the product onto exposed food and feed processing, preparation and serving surfaces or introducing the material into the air. Avoid contamination of food and feed processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AND FEED AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

SERVING AREAS — Facilities where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held.

Apply as a spot treatment to selective surfaces such as baseboards, under elements of construction into cracks and crevices. Avoid treating surfaces likely to be contacted by food. (Do not apply when facility is in operation or foods are exposed.)

NON FOOD AREAS — Includes garbage rooms, lavatories, floor drains (to sewer entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, incinerators and storage (after canning or bottling).

Apply to baseboard areas around water pipes, surfaces behind and beneath sinks, lockers, tables, pantries and similar areas where insects hide or through which they may enter.

RECOMMENDATIONS AND DIRECTIONS FOR USE

PROTECTION OF STORED GRAINS For the protection of stored grains such as wheat, oats, rice, corn, grain sorghum and field or garden seeds against Confused Flour Beetle, Rice Weevil, Granary Weevil, Saw Toothed Grain Beetle, Lesser Grain Borer and Indian Meal Moth. Apply 1 part 57% EMULSIFIABLE EMULSION MALATHION in 2 to 5 gals. of water per 1,000 bushels. Apply as the grain is being loaded or turned into final storage.

FOR PROTECTION OF STORED PEANUTS against Red Flour Beetle, Indian Meal Moth, Confused Flour Beetle, Rice Weevil, Flat Grain Beetle, Rusty Grain Beetle, Lesser Grain Beetle, Granary Weevil and Saw Toothed Grain Beetle.

Residual Warehouse Spray (before storing peanuts): Clean the warehouse thoroughly of old peanuts 1 to 2 weeks before new peanut crop is stored. Then, thoroughly spray the interior of the empty warehouse, especially cracks and protected places. Treat outside walls to a height of 6 to 8 ft. and the ground to a distance of 6 ft. from the warehouse. Use 1 pint of Malathion Premium Grade 57% Emulsifiable Emulsion in sufficient water to make 2 1/2 gals. of spray or 1 gal. with 1 1/2 gals. of water. Apply as a coarse spray at the rate of 2 gals. per 1,000 sq. ft. of surface or to runoff.

Bulk Spray Treatment (peanuts going into storage): Use Malathion Premium Grade 57% Emulsifiable Emulsion at the rate of 2 1/2 pints per 5 gals. of water for each 15 tons of farmers stock peanuts as they go into storage. Use good spray equipment. Preferably use a suitable mechanical spray applicator that regulates the rate of application to the flow of peanuts. Adjust the operating pressure of spray pump and size of nozzle opening to correlate the amount of spray delivery with the rate of flow of peanuts being treated. Avoid spraying with a fine mist that drifts away by using low nozzle pressure. Shield the nozzle against wind and air currents. After peanuts have been bulk treated use Malathion Premium Grade 25% Wettable Powder as a supplemental surface spray as follows: use 1 1/4 lb. of Malathion Premium Grade 25% Wettable Powder in 2 gals. of water for each 1,000 sq. ft. of surface. Apply the first surface treatment as soon as the bins are filled and leveled, but not later than the first week in October. Apply the second surface treatment one month later, followed by subsequent treatment at 2 month intervals. For applying the wettable powder surface treatment, use a coarse spray. Use spray equipment with sufficient capacity and power to treat large surfaces thoroughly.

HOUSEHOLD INSECTS

For the control of ants, clover mites, crickets, earwigs, pantry pests, roaches, scorpions, silverfish and spiders in and about house, dairy and food processing plants, use Malathion 57% Emulsifiable Emulsion at the rate of 6 1/2 fluid ounces per gallon of water. Apply by means of a coarse spray or paint brush to walls, window frames, baseboards, under sinks, drainboards, stoves, cracks and crevices and in other areas frequented by insects.

For Control of Bedbugs in dwellings, use Malathion 57% Emulsifiable Emulsion at a rate of 1 to 2 fluid ounces per gallon of deodorized kerosene. Apply lightly to all infested surfaces in sufficient quantity to moisten the fabric and generously to beds and woodwork with special care taken to wet all possible harboring places.

For Fleas and Brown Dog Ticks inside dwellings, thoroughly spray around the stoves, window frames, wall cracks, baseboards, closets, storage cabinets, openings around water pipes, or wherever there are likely to enter, congregate, feed or hide in the home or other buildings, plus the sleeping quarters of household pets (dogs and cats) and localized areas of floors and wall covering. Place fresh bedding on animal quarters following treatment. Avoid contamination of food, utensils, table, table equipment or linens. Repeat applications as necessary.

Pantry pests such as the exposed stages of saw toothed grain beetle, Indian meal moth, lesser grain borer, and Indian meal moth.

NOTE: This is the only use of this formulation as a pre-treatment for stored grain. It is not to be used on grain in other and asphal materials, such as tiles, carpet, and so on.

CONTINUED FROM SIDE PANEL

BEST DOCUMENT AVAILABLE

ACCEPTED
WITH CERTAIN
IN E.P.A.

10370-58

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH 567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: ☐ REGISTRATION ☐ REREGISTRATION
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

16L 93,306
 18 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been used by others.

☐ ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
--	-------------