

ford's

**SBAN* 1-E
INSECTICIDE**

sets of Households, Turf and Ornamental Plants, Mosquito Control and Area Control of Ticks and Chiggers, or under the supervision of pest control operators, and other trained personnel responsible for insect control programs.

(3,5,6-trichloro-2-pyridyl) phosphorothioate 12.6%
 (6-tert-butyl-4-methylpyridin-2-yl) phosphorothioate 21.5%
 Inert 65.9%

**OUT OF REACH OF CHILDREN
WARNING**

CAUTIONARY STATEMENTS

Hazards to Humans
 IRRITATION — HARMFUL IF SWALLOWED

Get in eyes, on skin or clothing • Wash thoroughly after handling
 before reuse • Avoid breathing vapors and spray mist • Avoid spraying
 on food, feedstuffs, and domestic water supplies • Keep container
 in a ventilated area.

METHOD OF PRACTICAL TREATMENT

Apply thoroughly. Contains aromatic petroleum solvent. Call a physician in
 event of contact, remove contaminated clothing and immediately wash
 with water. If in Eyes: Flush eyes with plenty of water for at least 15

minutes. It is a cholinesterase inhibitor. Treat symptomatically. Atropine only

PHYSICAL OR CHEMICAL HAZARDS

Do not store near heat or open flame • Do not cut or weld container.

PRECAUTIONS FOR ADDITIONAL PRECAUTIONS

AGRICULTURAL CHEMICAL

Do not store with food, feed, drugs or clothing.

NET CONTENTS: 1 GALLON

MANUFACTURED BY

Chemical & Service, Inc.

2730 PASADENA BLVD.
 PASADENA, TEXAS 77502

**PRECAUTIONARY STATEMENTS
ENVIRONMENTAL HAZARDS**

SBAN 1-E is toxic to fish, birds, and some animals. Do not apply directly to water
 streams or ponds. SBAN 1-E may be toxic to aquatic organisms. Do not apply directly to
 the soil. Do not apply where runoff is likely to occur. Do not apply where runoff
 contains toxic materials. Do not contaminate soil by cleaning of equipment
 used or disposal of wastes. Apply this product only as specified on this label.
 This product is highly toxic to bees exposed to direct treatment or residues on plants.
 Protective information may be obtained from your Cooperative Agricultural Extension
 Service.

PRECAUTIONS AT WORK: Do not use this product in a manner which would be
 likely to cause undue exposure to the user. Do not use this product where
 the label has been fully explained to you.

USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with the following

PESTS Indoors

General Information: Use SBAN 1-E insecticide to control the pests indicated in the
 table below by application as a 0.25% or 0.5% chlorpyrifos spray. Use the 0.25%
 spray to control light infestations and the 0.5% spray to control heavy infestations
 or as directed. To prepare any spray, dilute SBAN 1-E with water as shown
 in the following SPRAY SOLUTION CHART. Repeat the treatment as needed.

SPRAY SOLUTION CHART

| Gallons of Spray Solution Needed | Pound Amount* of SBAN 1-E TO USE | |
|-------------------------------------|----------------------------------|-------------|
| | 0.25% Spray | 0.50% Spray |
| 1 gallon | 0.25 | 0.50 |
| 2 gallons | 0.50 | 1.00 |
| 3 gallons | 0.75 | 1.50 |
| 4 gallons | 1.00 | 2.00 |
| 5 gallons | 1.25 | 2.50 |
| 10 gallons | 2.50 | 5.00 |

*Note: 1 Pound equals 16 ounces. 1 Gallon equals 128 ounces.
 NOTE: Where other insecticides and/or fumigants are desired, determine
 (DOSE) or percent may be added to the spray mixture containing SBAN 1-E in
 accordance with the label of the other product used to determine
 the amount needed to give the desired spray concentration. Do not mix and store in
 use and handling instructions for these products. Do not use SBAN 1-E in combina-
 tion with other insecticides (DOSE) or fumigants for treatment of food areas of food handling
 establishments.

Application within food handling establishments (except other than private residences)
 in which food is held, prepared or served (including, but not limited to, restaurants,
 grocery stores, bakeries, eating places, delicatessens and grain mills): Use as a dust treat-
 ment in and around food handling areas to control cockroaches (including the
 resistant to certain other insecticides) ants, crickets, beetles, silverfish weevils and
 stored product insects such as confused flour beetles, less than grain beetles, le-
 velled mill moths, rice weevils, lesser grain weevils and red flour beetles. Apply
 as a spray, use aerosol (20 gal or less) spray in isolated areas where the above pests
 have been seen or suspected of being or breeding. Apply only enough spray to
 thoroughly cover the surfaces treated using special care to avoid unnecessary runoff.
 Do not introduce the spray into the air or allow the spray to contact food or food-contacting
 surfaces. Avoid treated areas except for cleanup of spills and cleanup, along and behind
 bars, counters, storage and behind doors. Refrigeration areas and similar food preparation
 areas and equipment, floor drains, and around plumbing and other utility equipment.
 For ants apply to ant trails and wherever these pests may be present. Spot treatment
 may include cracks and crevices treatment by applying small amounts of material directly
 into openings leading to walls and below baseboards in walls and on and around equipment,
 legs and bases of equipment. Equipment capable of delivering a pressure of insecti-
 cide should be used. Repeat treatment as needed but not more than once every 7 days
 in restaurants and similar food service establishments. In case of emergency (call back),
 SBAN 1-E may be applied after 2 days from the treatment, but emergency use should
 be limited to once per year.

Applications of this product in food handling establishments other than as a spot and/or
 crack and crevice treatment are not permitted.
 Application within residential buildings including homes and apartment buildings and
 within industrial areas of industrial, institutional, and commercial buildings including
 hospitals, stores, manufacturing plants and warehouses. For spot treatment only — Apply
 as a contact spray of water suspension to localized areas where cockroaches (including
 strains resistant to certain other insecticides), ants, silverfish weevils, lesser grain
 weevils, and lesser grain beetles are normally occurring including dark corners of rooms and
 cracks, cracks and crevices in walls, along and behind baseboards, beneath and behind
 doors, around plumbing and other utility installations.
 For ants apply to ant trails, also around doors and treatment and wherever else these
 pests may be present. For cockroaches use a 0.5% spray only. Use a period of
 4 to 7 days if necessary required for maximum effect on cockroaches.

CONTINUED ON BACK PANEL

CONTINUED FROM SIDE PANEL

For control of carpet beetles: Thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting under carpeting edge
 and furniture in carpets and on lamping and wherever else these insects are seen or suspected.

For control of flies and house flies: Thoroughly apply a 0.5% spray as a spot treatment to unclean areas such as pet pens and resting quarters,
 manure pits and feed troughs, along and behind baseboards, window and door frames and to other areas of their and keep covering and other pests
 may be present. On bedding of pigs should be removed and replaced with clean, fresh bedding after treatment. DO NOT TREAT PETS WITH THIS PRO-
 DUCT. To control the source of fly infestation, spot treating the treated premises should be treated with a product registered for application to animals
 (poultry houses). Do not apply more than 17 days of this product in poultry houses, junction and service boxes or other electrical equip-
 ment because of possible insecticide residue. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and
 water supplies. Thoroughly wash dishes and feed handling equipment with soap and water if they become contaminated by application of this product.
 Do not allow chickens or pigs to contact treated surfaces used for feed, water and other feedstuffs (poultry). Before spraying.

PESTS OUT-OF-DOORS

General Information: Use SBAN 1-E insecticide for control of certain pests on outside surfaces of buildings and on trees and ornamental plants
 for area control of ticks and chiggers and for mosquito control. Consult the following directions for dosage and application details to control the pests listed
 in the table below. Use SBAN 1-E at the rate of 5% fluid ounces per gallon of water (0.5% chlorpyrifos emulsion) to control
 ants, silverfish weevils, crickets, stored product insects, lesser grain beetles, levelled mill moths, rice weevils, lesser grain weevils and other
 pests on outside surfaces of buildings including porches, screens, window frames, doors, garages, refuse dumps and other
 areas where these pests are likely to be present. Do not treat or allow the spray to contact plants, animals, humans, birds, dogs, cats, or
 other animals or people who may be present. Repeat treatment as needed to maintain effectiveness.

NOTE: To help prevent infestation of buildings by ticks, crickets, silverfish weevils, lesser grain weevils, levelled mill moths, rice weevils, lesser grain weevils and other
 pests on outside surfaces of buildings, use the building treatment to a height of 2 to 3 feet where pests are active and they find entrance. Use 2 gallons
 of SBAN 1-E per 100 gallons of water and spray as a contact spray at the rate of about 10 gallons spray mixture per 1000 sq. ft. to thoroughly
 and uniformly wet the walls.

Pests of trees and other ornamental and recreational turf grass areas: Use SBAN 1-E to control pests listed in the following table by applica-
 tion of the recommended dosage and in accordance with the directions given below. Dilute SBAN 1-E in enough water to obtain complete and uniform
 coverage of pest-infested areas and apply as a contact spray using suitable application equipment. Thoroughly wet the vegetation after
 treatment to reach the insecticide into the leaf canopy as needed. For best results, use the spray as soon as possible after treatment.

DOSAGES and DIRECTIONS for USE on TURF

| PEST | Dosage of SBAN 1-E per 100 gal. W | REMARKS |
|---|---|---|
| Ants (Carpenter, Formicid, etc.) | 3.0 oz | Spray every 7 to 10 days until control is obtained. For all outdoor areas, including in housing (porches, patios, etc.) and in areas where they are active. |
| Chiggers (Chiggers, etc.) | 3.0 oz | Spray every 7 to 10 days until control is obtained. |
| Crickets (House, etc.) | 3.0 oz | Spray as directed to control pests on the lawn and edges of the lawn as recommended by your local Agricultural Extension Service. |
| Lesser Grain Weevils (Lesser Grain Weevils, etc.) | 6.0 oz | For lawn grass, spray every 7 to 10 days and actively feeding near the soil surface. Usually active in July and August. If infestation is severe, repeat application as needed. Do not use if there is a possibility of contact with people or animals. |
| Levelled Mill Moths (Levelled Mill Moths, etc.) | 3.0 to 12.0 oz | Spray every 7 to 10 days until control is obtained. |

DOSAGES and DIRECTIONS for USE on ORNAMENTAL PLANTS

| Pests Controlled | Amount of SBAN 1-E to Water in Galons | | |
|---|---------------------------------------|---------------|-----------------|
| | 5 Gallons | 10 Gallons | 100 Gallons |
| Ants (Carpenter, Formicid, etc.) | 1.0 oz | 2.0 oz | 2.0 oz** |
| Chiggers (Chiggers, etc.) | 2.0 oz | 4.0 oz | 4.0 oz** |
| Crickets (House, etc.) | 2.0 oz | 4.0 oz | 4.0 oz** |
| Lesser Grain Weevils (Lesser Grain Weevils, etc.) | 4.0 oz | 8.0 oz | 8.0 oz** |
| Levelled Mill Moths (Levelled Mill Moths, etc.) | 1.0 to 4.0 oz | 2.0 to 8.0 oz | 2.0 to 8.0 oz** |

*Equivalent to 1 ounce of chlorpyrifos per 100 gallons
 **Equivalent to 3 ounces of chlorpyrifos per 100 gallons

CONDITIONS and RATES to USE for MOSQUITO CONTROL

| Condition of Area to be Treated | Amount to Use per Acre | | Number Acre of Water |
|--|-------------------------------|---------------|----------------------------|
| | of SBAN 1-E (fluid ounces) | (pounds) | |
| Light to medium vegetative cover | 1.5 to 3.0 | 0.075 to 0.15 | 150 to 300 |
| Medium to heavy vegetative cover | 3.0 to 6.0* | 0.15 to 0.3* | 30 to 60 |
| Adult Control — Light to medium vegetative cover | 3.0 | 0.15 | 30 |
| Medium to heavy vegetative cover | 6.0 | 0.3 | 60 |

*The higher dosage will give better knockdown and longer residual control where there is less or no vegetative cover.

10370-64

Ford's

CONTINUED FROM BACK PANEL

Apply DURSBAN 1-E to the trunk and lower ends of structural trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service for proper time to treat in your area. Mix in the ratio of one gallon of DURSBAN 1-E to 100 gallons of water and spray uniformly on trunk at a 200-300 ft. rate.

Apply DURSBAN 1-E to the trunk and lower ends of structural trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service for proper time to treat in your area. Mix in the ratio of one gallon of DURSBAN 1-E to 100 gallons of water and spray uniformly on trunk at a 200-300 ft. rate.

Apply DURSBAN 1-E to the trunk and lower ends of structural trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service for proper time to treat in your area. Mix in the ratio of one gallon of DURSBAN 1-E to 100 gallons of water and spray uniformly on trunk at a 200-300 ft. rate.

Apply DURSBAN 1-E to the trunk and lower ends of structural trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service for proper time to treat in your area. Mix in the ratio of one gallon of DURSBAN 1-E to 100 gallons of water and spray uniformly on trunk at a 200-300 ft. rate.

Apply DURSBAN 1-E to the trunk and lower ends of structural trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service for proper time to treat in your area. Mix in the ratio of one gallon of DURSBAN 1-E to 100 gallons of water and spray uniformly on trunk at a 200-300 ft. rate.

DURSBAN* 1-E INSECTICIDE

Controls Numerous Pests of Households, Turf and Ornamental Plants. Also Recommended for Mosquito Control and Area Control of Ticks and Chiggers. To be applied only by or under the supervision of pest control operators, public health organizations, mosquito abatement districts, and other trained personnel responsible for insect control programs.

Table with 2 columns: Ingredient Name and Percentage. Includes Chlorpyrifos (12.6%), Aromatic Petroleum Distillate (21.5%), and Inert Ingredients (65.9%).

KEEP OUT OF REACH OF CHILDREN WARNING PRECAUTIONARY STATEMENTS Hazards to Humans CAUSES EYE IRRITATION - HARMFUL IF SWALLOWED

Do not take internally. Do not get in eyes, on skin or clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs, and domestic water supplies. Keep container closed. Handle concentrate in a ventilated area.

STATEMENT OF PRACTICAL TREATMENT If Swallowed: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. If On Skin: In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water. If In Eyes: Flush eyes with plenty of water for at least 15 minutes. Call a physician. Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

PHYSICAL OR CHEMICAL HAZARDS COMBUSTIBLE - Do not use or store near heat or open flame. Do not cut or weld container. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS AGRICULTURAL CHEMICAL Do not ship or store with food, feed, drugs or clothing. EPA Reg No. 10370-64 NET CONTENTS: 1 GALLON

MANUFACTURED BY Ford's Chemical & Service, Inc. 2739 PASADENA BLVD. PASADENA, TEXAS 77502

PRECAUTIONARY STATEMENTS ENVIRONMENTAL

DURSBAN 1-E is toxic to fish, aquatic invertebrates, and birds. Do not apply where it will contact water. Do not apply where it will contact birds. Do not apply where it will contact bees. Do not apply where it will contact honey bees. Do not apply where it will contact beneficial insects.

USE: This is a statement of Federal law to the effect that any person who manufactures, distributes, or sells this pesticide in violation of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) is guilty of a criminal offense.

Application within 100 feet of water bodies is prohibited. Do not apply to water bodies. Do not apply to lawns, lawns, or other areas where it will contact water. Do not apply to lawns, lawns, or other areas where it will contact water.

Application within 100 feet of water bodies is prohibited. Do not apply to water bodies. Do not apply to lawns, lawns, or other areas where it will contact water. Do not apply to lawns, lawns, or other areas where it will contact water.