

57% MALATHION EMULSIFIABLE CONCENTRATE**ACTIVE INGREDIENT:**

Malathion: 0,0-dimethyl phosphorodithioate of
diethyl mercaptosuccinate.....57.00%

INERT INGREDIENTS**.....43.00%

TOTAL.....100.00%

**This product contains xylene range aromatics

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with soap and water.

IF IN EYES: Flush with water for 15 minutes. Contact a physician.

IF INHALED: Remove victim to fresh air and apply artificial respiration if indicated.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing vapors or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat and open flame.

EPA REG. NO. 10370-68
EPA EST. NO. 10370-TX-1

PAGE 1

Not Reviewed. Registrant claims to be in compliance with
Registration or Pesticide Product Guidelines (21)
in force. Page 2 12. Active to Inert Change in

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store product in original labeled container in a cool, dry, locked place out of the reach of children.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

PESTICIDE DISPOSAL: Securely wrap container in several layers of newspaper and discard in trash.

NOTE: Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces. Do not allow contact with treated surfaces until spray has dried.

HOUSE FLIES

For reduction of fly population use 2 gallons per 100 gallons of water, or 1 pint in 6 gallons of water for control in dairy barns, horse and sheep barns, poultry houses, corrals, manure piles, garbage heaps, city dumps, etc. - apply spray thoroughly to wet ceilings, walls and other places where flies congregate.

FOR A BAIT SPRAY - use 2 gallons with 2 gallons unsulfurized molasses or corn syrup, or 20 pounds of sugar per 100 gallons of water. Repeat application as necessary.

NOTE: Do not use where milk is processed or stored or in human dwellings. Avoid contamination of feed and feed products. Remove animals and poultry before treating. **DO NOT USE ON ANIMALS.**

FLEAS

For control of fleas in dog houses, garages and other buildings, except dwelling houses, use a mixture of 5 ounces product per gallon of water - apply at a rate of 1 gallon per 1,000 sq. ft. Remove animals before spraying. Thoroughly treat infested areas. Fresh bedding should be placed in animal quarters following treatment. Repeat application if necessary. Do not use on pets and other animals. To control fleas on pets and other animals, use an insecticide recommended for this purpose.

ORNAMENTALS

For the control of aphids, spider mites, whiteflies, mealybugs, thrips, Japanese beetle adults, four-lined leaf bug, tarnished plant bug, rose leafhopper and potato leaf hopper, use 1-1/2 pints per 100 gallons of water, making one or more applications as needed.

Manufactured by
FORD'S CHEMICAL & SERVICE, INC.
2739 Pasadena Blvd.
Pasadena, TX 77502