

Reg # 10370--71

PM-10

191

4/4/91

Not Reviewed. Registrant claims to be in compliance with...

WARNING!

Before using this compound, exposed food products and packaging materials must be removed from the room or protected. After using this compound, equipment and utensils must be thoroughly rinsed with potable water before operations are resumed. Application confined to time periods when plant is not in operation. CONTACT SPRAY: Dilute with Deodorized Kerosene on a 1 to 6 basis. To be used as a surface spray for control of roaches, waterbugs and silverfish in the edible products areas of Federally Inspected Meat and Poultry Processing Plants. For control of roaches, waterbugs and silverfish, thorough and repeated spraying of the infested area, working the spray into cracks and other hiding places is recommended. Emphasis should be placed on hitting the insects directly and on repeated applications. Direct the spray to insect hiding places.

PROFESSIONAL Spray Concentrate

FOR USE IN FEDERALLY INSPECTED MEAT & POULTRY ESTABLISHMENTS

For control of Roaches, Waterbugs & Silverfish
For use by Pest Control Operators Only

ACTIVE INGREDIENTS:

Pyrethrins	0.62%
Piperonyl Butoxide, Technical*	5.03%
Piperonyl Butoxide	94.35%
TOTAL:	100.00%

*Equivalent to 4.02% (Butylcarbityl) (6-propylpiperonyl) Ether and 1.01% related compounds.

WARNING

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

CONTAINER DISPOSAL-Do not reuse container. Crush used container and bury in non-cropland soil away from water.

FOGGING

Close windows and doors and shut off ventilating system. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 oz. per 1000 cu. ft. of space. Direct the spray toward the ceiling and upper corners of area and behind obstructions. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment as necessary. Contact spray treatment for crawling insects should be used in combination with the fogging treatment.

The above described application will kill accessible, exposed stages of flies, fruit flies, mosquitoes, silverfish, small flying moths, crickets, gnats, roaches, spiders, scorpions, wasps, hornet, clerid mites, cheese mites, cheese skippers and exposed or accessible stages of stored product pests: granary weevils, rice weevils, confused flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, meal worms and grain mites.

E.P.A. Reg. No. ~~10370~~ 10370-071
NET CONTENTS _____ GALLONS