

Reg # 10376-83

PM-10

192

Ford's 4/4/91

MULTIPURPOSE AEROSOL

CONTAINS PYRENONE

PROFESSIONAL STRENGTH — LOW DOSAGE REQUIRED

For use in Schools, Hospitals, Institutions, Supermarkets, Dairies, Hotels, Restaurants, Drive-ins, Motels, Drug Stores, Food Plants, Milk Rooms and Animal Quarters.

Provides rapid control of Flies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Roaches, Ants, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders.

KILLS AND REPELS Stable Flies, House Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats. Controls Garden insects on Ornamental and Vegetable Plants. Acceptable for use to day of harvest on edible plants.

ACTIVE INGREDIENTS 100%

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Hydrocarbons	4.3%

** INERT INGREDIENTS: 94.5%

TOTAL: 100.0%

** CONTAINS PETROLEUM DISTILLATE

*Equivalent to 4% (butylcarbityl) (6-propylpiperonyl) ether and 1% related compounds.

PYRENONE—Registered trademark of Fairfield American Corporation.

CAUTION: Keep out of Reach of Children
See Additional Precautionary Statements

PRECAUTIONARY STATEMENTS—HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After spraying in processing plants, wash with portable water all equipment, benches, shelving, etc., where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL & PHYSICAL OR CHEMICAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE FOR USE

STORAGE: Contents under pressure. Do not store near heat or open flame. Store only in locked storage areas inaccessible to children.

DISPOSAL: Replace cap and discard containers in trash. Do not incinerate or puncture.

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE. Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

STORAGE AND DISPOSAL:

STORAGE: Contents under pressure. Do not store near heat or open flame. Store only in locked storage areas or areas inaccessible to children.

DISPOSAL: Replace cap and discard containers in trash. Do not incinerate or puncture.

Not Reviewed. Registrant claims to be in accordance with:
Reregistration of Pesticide Product Guidance for Phase 2
estimate, Para 2.12, Article to Inert Chemicals.

(Page 1 of 2)

Roaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Sow Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary. **Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Exposed States Indian Moths and Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flour Beetles, Weevils and Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

To Treat Animals: To protect beef and dairy cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters and Milk Rooms: For rapid control house flies, mosquitoes, gnats, wasps and small flying moths, close windows and doors and spray at the rate of 1 second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

For Use in Gardens: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

USE ON	TO CONTROL
Roses	Aphids, Rose Chafers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scales.
Asters	Leafhoppers, Tarnish Plant Bugs, Lace
Dahlias	Aphids, Red Spider Mites, Plant Bugs
African Violets	Broad Mites, Citrus Mealybugs
Azaleas	Azalea Leaf Bugs, White Flies, Leaf Miners
Euonymus	Euonymus Scales, Aphids, Thrips (exposed)
Hydrangeas	Aphids, Red Spider Mites, Hydrangea Leafhoppers
Junipers	Juniper Scales, Red Spider Mites, Bagworms
Lilacs	Leaf Miners, Leaf Rollers, Scales
Tomatoes	Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Green Peach Aphids
Pepper, Eggplants, Potatoes	Colorado Potato Beetles, Leafhoppers, Blister Beetles, Flea Beetles, Green Peach Aphids
Cucumbers, Squash, Melons	Striped & Spotted Cucumber Beetles
Beans	Mexican Bean Beetles, White Flies, 12-Spotted Beetles
Sweet Corn	Japanese Beetles

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 10370-83
EPA Est. No. 10370-TX-1
Net Contents:

Manufactured By
Ford's Chemical & Service, Inc.
2739 Pasadena Blvd., Pasadena, Texas 77502