

Reg # 10370-92

PM-10

4/4/91

191

ford's

PYRE-DUST ROACH POWDER

**PYRENONE® PROVIDES QUICK INSECT CONTROL
CONTROL COCKROACHES, CRICKETS, FRUIT FLIES AND SIXTEEN COMMON GARDEN PESTS.
LEAVES NO OBJECTIONABLE RESIDUES**

For use in Residences, Institutions, Food Plants, Canneries, On Stored Seed and Grain and On Vegetable and Ornamental Plants
Protects Stored Grain and Seed from Insect Infestations.

ACTIVE INGREDIENTS.

Pyrethrins
*Piperonyl Butoxide, Technical
Petroleum Distillate
** INERT INGREDIENTS

*Equivalent to 0.8% (butylcarbamyl) (6-propylpiperonyl) ether and 0.2% related compounds. TOTAL
PYRENONE® Registered trademark of Fairfield American Corporation

0.10%
1.00%
~~1.10%~~
~~2.10%~~ 98.92
100.00%

* Contains Petroleum Distillate

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures under the Resource Conservation and Recovery Act. CONTAINER DISPOSAL: Dispose of bags according to approved Federal state or local procedures under the Resource Conservation and Recovery Act.

FOR CONTROL OF COCKROACHES AND CRICKETS: Using conventional dusting equipment, apply this product at 2 ounces per 100 sq. ft. to all indoor areas where these pests may crawl or hide. Apply in cracks and crevices and hidden surfaces around, under and behind baseboards, cabinets, refrigerators, stoves, sinks, doors, pipes, windows, electrical fittings, electrical equipment and other possible harborages. Repeat application as required.

ON VEGETABLE AND ORNAMENTAL PLANTS: Pyre-Dust Roach Powder may be used throughout the growing season, or as a preharvest treatment. Acceptable for use up to the day of harvest on edible plants when other insecticides are prohibited due to residue restrictions and use limitations. Apply dust uniformly to upper and lower leaf surfaces at the rate of 30 pounds of dust per acre. Treat only when the insects are present and repeat as required to maintain control.

CROP	TO CONTROL
ORNAMENTALS	Aphids, Flea Beetles, Leafhoppers, White flies on African Violets, asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwoods, Geraniums, Gladioli, Marigolds, Roses, etc.
ASPARAGUS	Asparagus Beetles
BEANS	Aphids, Leafhoppers, White flies, Mexican Beetles
BROCCOLI	Aphids, Cabbage Loopers
CABBAGE	Cross striped Cabbageworms
BRUSSEL SPROUTS	Diamondback Moth Larvae
CAULIFLOWER	Flea Beetles, Harlequin Bugs, Imported Cabbageworms
CELERY	Green Peach Aphids, Leaf Tiers, Cabbage Loopers

CROP	TO CONTROL
CRANBERRIES	Fireworms, Leafhoppers
EGGPLANTS	Colorado Potato Beetles, Flea Beetles, Green Peach Aphids
LETTUCE	Cabbage Loopers, Green Peach Aphids, Diamondback Moth Larvae, Imported Cabbageworms
PEPPERS	Green Peach Aphids
POTATOES	Aphids, Colorado Beetles, Flea Beetles, Leafhoppers
RADISH	Aphids, Flea Beetles
SPINACH	Aphids, Cabbage Loopers, Webworms
TOMATOES	Green Peach Aphids, Flea Beetles, Colorado Potato Beetles

Fruit Fly Control of Fruits and Vegetables: To control Fruit flies on apples, blackberries, boysenberries, currants, dewberries, grapes, loganberries, peaches, pears, plums, prunes, melons and tomatoes, apply as a post-harvest application. Dust fruits and vegetables immediately after picking while in containers in the field. Remove crops from field as soon as they are picked and have been dusted. Do not hold picked crops in the field overnight. If the crops are held at the processing plants more than 20 hours, a pre-harvest application should be made. Dust lightly but thoroughly.

Stored Grain Protectant. For use on corn, barley, rice, rye, buckwheat, birdseed, and other seeds in storage. This product kills accessible stages of many grain insects, particularly rice weevils and granary weevils, and protects against many insects in the bran bug group, including flour beetles and grain beetles. One treatment is usually adequate for a full storage season. Treated grains require only normal cleaning before processing for human food and may be fed to livestock without hazard. Shelled corn may require cleaning, subject to inspection for grading purposes. Whole grains and seeds to be treated should be in condition for normal storage. Treatment should be applied when grain goes into storage and before serious infestation has developed.

How much to apply: Application on grains and seeds should be made at the rate of 80 to 100 pounds per 100 bushels (13 ounces to 1 pound per 10 bushels). The dust should be sufficiently well mixed with the whole grain seeds to assure even distribution. It can be applied to a moving stream of grain in the usual or typical conveyor. Small lots of grain or seeds can be treated by hand, with a shovel or other suitable equipment. On ear corn this insecticide should be applied at the same rate specified above and should be sufficiently well mixed with the corn to assure even distribution. One method is by dusting the protectant over alternate shallow layers of the corn.

EPA Reg. No. 10370-92
EPA Est. No. 10370-TX 1

ford's

Manufactured By
Chemical & Service, Inc.
2739 Pasadena Blvd.
Pasadena, Texas 77502



4/86

CPS 7004

Not Reviewed. Registrant claims to be in accordance with