

12  
VOT I

**FOR THE EFFECTIVE CONTROL OF APHIDS, SPIDER MITES, FLEAS, CHICKEN RED MITES AND OTHER INSECTS**

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS & DOMESTIC ANIMALS**  
**CAUTION** Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin or clothing. Wash skin thoroughly after handling. Remove and wash contaminated clothing before reuse.  
**ENVIRONMENTAL HAZARDS**  
 This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.  
**BEE CAUTION** This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treated area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**DIRECTIONS FOR USE**  
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
 Do not treat areas where unprotected humans or domestic animals are present. Do not allow entry into treated areas until sprays have dried or dusts have settled, unless full protective clothing is worn.

**STORAGE AND DISPOSAL**  
 Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Apply this product only as specified on this label. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

PLANT	PEST	AMOUNT		Days Before Harvest	
		Dust Per Acre	Number of Applic. Applications	Last Applic. Application	Days
<b>VEGETABLES</b> (Thorough full coverage applications should be made)					
Tomatoes	Spider mites	35-45 lbs	1-2	3	
	Aphids	35-45 lbs	1-2	3	
Peas	Trioxys, lesser mites	35-45 lbs	1-2	3	
	Leafhopper	25 lbs	2-3	3	
Beans	Meadow sparrow beetle	30-35 lbs	2-3	3	
	Leafhopper	30-35 lbs	2-3	3	
Broccoli	Aphids	30 lbs	1-2	7	
Cabbage	Imported cabbageworm	30 lbs	1-2	7	
Kale	Cabbage looper				
Mustard	Cabbage looper				
Peanut	Aphids	30 lbs	1-2	7	
	Aphids	50 lbs	1-2	7	
Soybean	Aphids	30-35 lbs	1-2	3	
	Spider mites	30-35 lbs	1-2	3	
Watermelon	Leafhopper	30-35 lbs	1-2	3	
Do not apply to other crops or plants unless permitted by label.					
Potatoes	Leafhopper	25-30 lbs	3-4	3	
	Thrips	30-40 lbs	3-4	3	
Spinach	Aphids	30-35 lbs	1-2	7	
	Aphids	30-40 lbs	1-2	7	
Lettuce	Mites	30-40 lbs	1-2	7	
	Cabbage looper	30-40 lbs	1-2	7	

**ACCEPTED**  
 SEP R 1977  
 UNDER THE FEDERAL PESTICIDE  
 FURNISHING AND RECORDING ACT  
 FOR BOOKING OF THIS PESTICIDE  
 ED WITH NO. 3 28 64 4



**ACTIVE INGREDIENT:** Malathion\* ..... 5.0%  
**INERT INGREDIENTS:** ..... 95.0%  
 \*O-O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**CAUTION**  
 Keep Out Of Reach Of Children.  
 See back panel for additional cautions.  
 Read All Directions Before Using.  
 IN CASE OF CONTACT: Wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes. Get medical attention if irritation persists.  
 NOTE TO PHYSICIAN: Malathion is an acetyl cholinesterase inhibitor. Atropine is antidotal.

E.P.A. Reg No 3286-44      E.P.A. Est. No. 3286-TX-1  
 Net Weight 1 lb.  
 FERD STAFFEL COMPANY, San Antonio, TX 78298