Please read instructions on reverse before	completing form.	Form Appr	roved. OMB No. 2070	0-0060. Approval expires 2-28-9!
United States Environmental Protection Age Washington, DC 20460		incy	Registratio Amendment X Other	
Application for Pesticide - Section I				
1. Company/Product Number 10370-129		2. EPA Product Mana D. Edwards	ger	3. Proposed Classification
4. Company/Product (Name) Ford's Sevin 10 Dust		РМ# 19		X None Restricted
5. Name and Address of Applicant (Include ZIP Code) AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale NJ 07645 Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name		
		tion - II		
Amendment - Explain below. Resubmission in response to Agen Notification - Explain below.	cy letter dated	Final printed Agency letter "Me Too" Ap Other - Explain	pplication.	
Explanation: Use additional page(s) if necessary. (For section and Section .) This notification is consistant with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistant with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under 12 and 14 of FIFRA. Alternate name: Ford's Sevin Dust 10%. Additional bullet point; Use Indoors and Outdoors. Amended Environmental Hazards section as per PR Notice 93-8.				
Section - III				
1. Material This Product Will Be Packaged In:				
Yes' No If 'Yes'	Yes No		Pla Gla Pag	ainer stal stic ass per her (Specify)
be submitted 3. Location of Net Contents Information	4. Size(s) Retail Contail	t	. Location of Label Di	
Label Container			On Labeling a	ccompanying product
6. Manner in Which Label is Affixed to Product Lithograph Other Stencifed				
		ion - IV		
1. Contact Point	below for identification of indivi	dual to be contected, if	necessary, to process	this application.)
Name Christina M. Lawes	Titte Manag	er, Federal F		phone No. (Include Area Code) 5 (201) 307-3355
I certify that the statements I have n I acknowledge that any knowingly for both under applicable law.			•	ಕೆ. Date Application Beckined ^Stamped)
2. Signature With Maan	3. Title Manag	er, Federal I	Registration	is
Christina M. Lawes	5. Date	January 21	1, 1996	



SEVIN® DUST 10%

- FOR INSECT CONTROL ON VEGETABLE CROPS
- ORNAMENTALS DOGS AND CATS
- APPLY AS A CRACK AND CREVICE TREATMENT FOR INDOOR **PEST CONTROL**
- Use Indoors and Outdoors

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate) 10.0% INERT INGREDIENTS:.....90.0%

Contains Sevin® brand Carbaryl Insecticide Sevin® is a Registered Trade Mark of Rhone-Poulenc, Nederland B.V. for Carbaryl Insecticide.

EPA REG. NO. 10370-129

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN

CAUTION

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado amplia-

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back or throat with linger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Apply Artificial Respiration if needed.

IF IN EYE5: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, Opiates, or cholinesterase inhibiting drugs.

> See Side Panel For Additional Precautionary Statements

> > **NET WEIGHT:**

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. For indoors when treating attics, storage rooms or other enclosed areas such as crawl spaces wear a pesticide respirator jointly approved by the Mine Safety and Health Administration and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed.

RE-ENTRY & WORKER SAFETY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dust has settled. Because certain States may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Environmental Hazards

This products is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is extremely toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the proposed treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other Carbaryl labels.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store product in original container in a cool, dry, locked place out of reach of children.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a senitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Household Disposal: Do not re-use empty container. Rinse burning. If burned stay out of smoke.

thoroughly before discarding in trash.

VEGETABLES AND ORNAMENTALS Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes, ornamentall. gardens or parks.

Apply when air is calm, using a hand duster. Unless otherwise specified, apply at the first sign of infestation and repeat at 7 to 10 day interval when needed to maintain course. Observe use limits given for specific crops.

MAY BE APPLIED UP TO DAY OF HARVEST ON FOLLOWING CROPS.

CORN: For control of *Corn Earworm* apply 5-1/2 to 7-1/2 oz, per 1,000 sq. ft. starting when the first silks appear. Repeat at 2 to 3 day intervals throughout the silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. May be applied up to harvest on Sweet Corn ears for human consumption. To avoid excessive residues on fodder or forage, do not apply within 7 days of harvest. Reapply immediately after rain. Applications of Sevin to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

CUCUMBERS, SUMMER SQUASH: For Flea Beetles, Melonworms, Pickleworms, Spotted Cucumber Beetle, and Squash Bugs apply 3-1/2 oz. per 1,000 sq. it. Apply when insect damage first appears and repeat at 10 to 14 day intervals.

EGGPLANT, PEPPER, TOMATO: For Fall Armyworm, Flea Beetle, and Tomato Fruitworm apply 5-1/2 to 7-1/2 oz. per 1,000 sq. ft. For European Corn Borer, Stink Bugs, and Tomato Hornworm, apply 5-1/2 to 7-1/2 oz. per 1,000 sq. ft. Apply when insect damage first appears and repeat as needed for control.

GRAPES (Western U.S.): For *Grape Leaf Folder, and Grape Leafhopper* early season control use 5-1/2 oz, per 1,000 sq. it, and 5-1/2 to 7-1/2 oz, for summer control. For the control of first brood Grape Leaf Folder apply when maximum number of eggs have hatched. For Leathopper control apply at first sign of infestation.

POTATOES: For *Colorado Potato Beetle, Flea Beetle, and Leaf-hoppers* apply 5-1/2 to 7-1/2 oz. per 1,000 sq. ft. Apply when insect damage first appears and repeat at 10 to 14 day intervals or as needed for control.

STRAWBERRIES: For *Meadow Spittle Bug and Strawberry Leaf Roller* apply 7-1/2 oz. per 1,000 sq. it. Apply when "Spittle" and/or leaf "Rolling" first appears. Repeat at 7 to 10 day intervals as needed for control. May be applied up to 3 days before harvest.

ORNAMENTALS. For control of the following apply when insects or their damage first appears and repeat as needed. Apply 5-1/2 to 7-1/2 oz. per 1,000 sq. it. or thoroughly apply to place a thin layer of dust on both upper and lower sides of leaves, stems and flowers. Controls: Apple Aphids, Bagworms, Blister Beetle, Boxelder Bug, certain Leaf Miners (Boxwood, Holly, Oak, and Spruce Needle), Elm Leaf Beetle, Exposed Thrips, Flea Beetles, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plants Bugs, Scale Insects, Tent Caterpillars.

In tests Sevin[®] Brand Carbaryl Insecticide has been used at recommended rates without damage to Andromeda, Arbor Vitae, Ash, Aster, Bamboo, Beech, Birch, Boxwood, Camellia, Carnation, Ceanothus, Cedar, Chinese Elm, Chrysanthemum, Cypress, Dahlia, Delphinium, Dogwood, Elm, Euonymous, Fir, Fuschia, Hemlock, Hickory, Holly, Hornbeam, Hydrangea, Inkberry, Iris, Juniper, Laurel, Lilac, Locust, Magnolia, Maidenhair Fern, Maple, Nasturtium, Oak, Orchid, Ornamental Mulberry, Petunia, Pine, Poplar, Privet, Rhododendron, Rose of Sharon, Sassaíras, Spirea, Verbena, Viburnum, Wigelia, Wisteria, Yew, Zinnia.

FOR FLEAS, TICKS, MITES, LICE FOR FLEAS AND BROWN DOG TICK CONTROL ON DOGS AND CATS*: Using rubber or neoprene gloves thoroughly rub 1 to 2 level tablespoonsfuls into the DOG'S coat, back line, neck, belly and legs, avoiding the eyes. Thoroughly clean the area and dust pet sleeping quarters (lightly), then outdoor runways, and pens using 1 lb. per 800 sq. ft. Replace with fresh pet bedding.

DO NOT USE ON PUPPIES OR KITTENS UNDER 4 WEEKS OF AGE. DO NOT USE ON PREGNANT DOGS. Make second application for newly hatched Fleas in 7 to 10 days or weekly as needed for continued control. *FOR CATS USE DUST SPARINGLY LIMITING THE TREATMENT TO 1 LEVEL TEASPOONSFUL PER CAT. While wearing gloves rub dust into fur. Make sure to treat back of neck and belly.

FOR MITES, LICE, FLEAS, AND TICKS ON CHICKENS: One pound is enough dust for 100 chickens. Can also be used on Pheasants. Pigeons, and Partridge. Do not repeat application to Poultry within four weeks. Dust chicken house using one pound per 40 sq. ft. Do not house birds in treated poultry house within seven days of slaughter. Do not dust birds within 7 days of slaughter. Do not dust nest or eggs.

INDOOR PEST CONTROL

INDOOR PEST CONTROL USE IS LIMITED TO CRACK AND CREVICES TREATMENT ONLY IN THE FOLLOWING DESIGNATED AREAS. THIS PRODUCT CANNOT BE USED INDOORS AS AN OVERALL BROADCAST TREATMENT.

WHERE TO USE: For use in and around buildings such as Homes, Apartments, Warehouses, Barns, and Municipal and Recreational Areas. DO NOT USE IN FOOD AREAS of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in service areas while food is exposed. Do not treat surfaces that leave visible dust residues. Remove any excess surface dust by brushing or wiping into the intended cracks or crevices. Apply with a Hand Dispenser.

HOW MUCH TO USE: Apply dust only into cracks and crevices. Any visible dust must be brushed into cracks or crevices.

HOW TO APPLY AS A CRACK AND CREVICE TREATMENT For control of Ants, Cockroaches, Crickets, Millipedes, Silverfish & Fleas which may hide, live and/or breed in cracks and crevices use a "Squeeze" or "Puri" Duster to apply thoroughly into all cracks and crevices adjacent to baseboards, corners, walls, door sills, and window frames, quarters of household pets, ant trails, outside perimeters of dwelling and other crack and crevices where insects may rest or hide such as in building foundations and entrances, in walls or floors near refrigerators, stoves, sinks, and around bookcases, or in cracks or crevices in attic areas (see Precautionary Statement) or other storage areas where insects may be present. For Fleas and Brown Dog Ticks thoroughly dust all cracks and crevices around and near pet living quarters, pet's bedding and resting areas. (See Directions for Use on Dogs and Cats). For Ants treat cracks and crevices where Ant trails may be present and where ants may gain entrance to buildings. Repeat as needed for continued control.

IMPORTANT NOTICE

When treating cracks and crevices in attics, storage rooms or other enclosed areas such as crawl spaces, for protection against carbaryl dust wear a pesticide respirator or dust mask jointly approved by the Mine Safety and Health Administration and by the National Institute for Occupational Safety and Health under provisions of 30 CFR Part 11.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645

BEST AVAILABLE COPY

IMPORTANT