

Reg # 10 70-157

PM 16

lot 2

ORIGINAL

Page 1 of 2

TREATS 5,000 SQ. FT.

Ford's

TURF-G **GRANULAR INSECTICIDE** Contains © **OFTANOL** Insecticide

ACCEPTED
SEP 10 1985
U.S. Federal Insecticide, Fungicide and Rodenticide Act
as amended for the pesticide registered under EPA Reg. No. 10370-157

FOR INSECT CONTROL ON TURF GRASSES

WHITE GRUBS - FIRE ANTS - SOD WEBWORMS - CHINCH BUGS

ACTIVE INGREDIENT:

1-Methylethyl 2-[[ethoxy((1-methylethyl)amino)phosphinothioyl]oxy]benzoate* 1.5%

INERT INGREDIENTS 98.5%

Total 100.0%

*U.S. Patent No. 3,621,082

Canadian Patent No. 825,198

STOP — READ THE LABEL BEFORE USE

CAUTION: KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Give a glass or two of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If in eyes, flush with plenty of water. If on skin, wash with soap and hot water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty container. Wrap and put in trash.

EPA REG. NO. 10370-RLO

EPA Est. No. 44616-MO-1

10370-TX-1

NET WEIGHT 15 POUNDS

MANUFACTURED BY

Ford's

Chemical & Service, Inc.

2738 PASADENA BLVD.

It is violation of Federal law to use this product in a manner inconsistent with its labeling.
IMPORTANT: Read these entire Directions for Use before using Turf-G Insecticide. Turf-G Insecticide is recommended for use on lawns, and other ornamental and recreational turf grass areas.

APPLICATION EQUIPMENT

Apply Turf-G uniformly over the turf area with suitable granular turf application equipment. Both drop-type and rotary-type spreaders may be used to apply Turf-G. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

APPLICATION TECHNIQUES

Turf-G applications using a pull-type drop spreader can be aided by a splash board mounted under the hopper for more uniform granule distribution. A more uniform application can be made by spreading half the required amount of Turf-G Granular over the area, and then immediately apply the remaining half at right angles or crosswise. The full recommended rate of Turf-G Granular must be used for maximum benefit.

When applying Turf-G, use appropriate marking systems and make application when the wind is calm to prevent missed area.

SPREADER SETTINGS

Approximate Spreader Settings for applying Turf-G at the rate of 3 pounds per 1,000 sq. ft.

Scotts PF-1/011B #6.0 : Cyclone B-1 #3.5. Consult Spreader Manual for 1/2 rat

RECOMMENDED APPLICATIONS

CROP	PEST	DOSAGE TURF-G	REMARKS
Turf Grasses	SOIL INSECTS White grub larvae (such as Japanese beetle, Black turf grass ataenius Chafers, <i>Phyllophaga</i> spp., Green June beetle, Oriental beetle, Asiatic garden beetle), Billbug (larval stage), <i>Hyperodes</i> weevil (larval stage) and Mole crickets.	3 lbs./1,000 sq. ft.	Be Sure to Read "APPLICATION EQUIPMENT" Section of this label. Do not apply more than 6 lbs. of Turf-G per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. ft. regardless of formulation used. Minimum interval between applications must be 30 days. For White Grub and Billbug Larval Control: Water turf area thoroughly immediately after application, using sufficient water to wet soil 1 to 1-1/2 inches deep. For <i>Hyperodes</i> Weevil Control: Apply between full bloom periods of forsythia and flowering dogwood, usually April 15 to May 15, and water-in as for white grub larval control. For Mole Cricket Control: Water immediately after application using 1/2 to 1 inch of water. Consult your local turf state Agricultural Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.
	FIRE ANTS	3 lbs./1000 sq. ft.	Area Fire Ant Control: Apply specified dosage uniformly over area to be treated for residual control. Water immediately after application using 1/2 to 1 inch of water. If established fire ant mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.
		0.5 oz. per 9 square feet	Established Fire Ant Mounds - Drench Treatments: Wet Conditions: Distribute dosage evenly over 9 sq. ft. and immediately sprinkle gently with 3 gallons of water. Dry Conditions: Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment.
	SURFACE INSECTS Sod webworm larvae, Chinch Bugs	1.5 or 3 lbs. /1,000 sq. ft.	Apply 1/2 to 1 inch of water immediately after treatment. Reapply as necessary for pest re-infestations. Do not apply more than 6 lbs. per 1,000 sq. ft. per year or an equivalent amount of 1.45 ounces active ingredient per 1,000 sq. feet regardless of formulation used. Minimum interval between applications is 15 days for 1.5 lb. rate and 30 days for the 3 lb. rate. Consult your local turf, state Agricultural Experiment Station, or state Extension Service Specialists for more specific information regarding timing of application.

GENERAL APPLICATION RESTRICTIONS

For effective insect control and as a general practice or as indicated above make sure to "water-in" the insecticide after each application. Do not mow treated turf area before rainfall or irrigation has been applied as per recommended applications. Do not graze treated areas or use clippings from treated areas for feed or forage. Keep children and pets off treated area until insecticide has been watered in and the grass is dry.

OFTANOL is a Reg. T.M. of the parent company of Farbenfabriken Bayer GmbH Leverkusen.

NOTICE: Buyer assumes all risk and liability of use, and/or handling of this material not in accordance with the terms of this label.