



P 2/2

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Thoroughly wash after handling and before eating or smoking. Keep pets and children off sprayed areas until dry.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water. Do not treat under the drip line or to the root systems of sensitive ornamental shrubs and plants. Clover, Dichondra and certain grasses such as Bermudagrass, St. Augustine or Carpetgrass are sensitive to herbicides and may be injured by the use this product. Do not contaminate waters used for irrigation or domestic purposes. Be careful when treating weeds not to spray surrounding desirable grasses unless damage can be tolerated. Avoid spraying and spattering of nearby flowers and other desirable plants.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not re-use empty container. Wrap and put in trash.

**PARTIAL LIST OF WEEDS CONTROLLED** - BLACK MEDIC, CANADA THISTLE, CHICKWEED, CLOVER, CURLY DOCK, DANDELION, ENGLISH DAISY, FLEABAN, FLORIDA PUSLEY, GROUND IVY, HEAL-ALL, HENBIT, KNOTWEED, LAMBSQUARTER, MUSTARD, PENNYWORT, MORNING GLORY, PIGWEED, PLANTAIN, POISON IVY, POISON OAK, PRICKLY LETTUCE, PURSLANE, SHEEP SORREL, SPOTTED SPURGE, THISTLE, WILD ONION. (This product will not kill "Grassy Weeds". Use Liquid Edger EPA Reg. No. 10370-146.)

**WHERE TO USE**- Lawns, Picnic Areas, around Patios and Driveways, Cemeteries, along and under Fencing, and Vacant Lots, Filling Stations, Parking Lots, around Schools, Playgrounds, Parks, Commercial Warehouses and Offices.

**WHEN TO APPLY** - Spot treat weeds when air temperature is above 70 degrees and the weeds are actively growing. Lower temperatures will reduce herbicidal effectiveness. Do not over dose. Spray when air is calm in morning or late afternoon to avoid wind drift of herbicide. Treat weeds at any time they become unsightly or when they present a hazard or problem.

**HOW TO APPLY** - This product is ready to use. Do not dilute. Hold sprayer as close as possible ( 1 to 2 feet ) to the target weed in order to wet the foliage. Direct the sprayer nozzle at each weed to point of run-off. Hit the growing point. Re-treat hard-to-kill weeds within 10 to 14 days.

**RESEEDING THE AREA** - After weeds have died remove the dead vegetation. Prepare the seed bed by additions of topsoil, peat or other amendments. Work into the soil. Wait 2 weeks then seed the area at 1 1/2 times the recommended seeding rate. Keep soil moist through the first cutting.

**NOTICE** - Buyer assumes all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.