

**ford's**

In Accordance with EPA Reg. 103.70-2-2,  
Based on Draft Labeling of 1971

## NUTGRASS & CHICKWEED KILLER

### A MSMA LIQUID CONCENTRATE FOR SELECTIVE POST-EMERGENT WEED CONTROL

One Pint Treats 4,000 Sq.Ft. of Turf

*Controls Crabgrass, Foxtail, Dallis Grass, Sand-Bur and For Grassy Weeds such as Johnson Grass and Certain Broadleaf Weeds such as Cocklebur, Chickweed and other weeds as listed.*

#### ACTIVE INGREDIENT.

Monosodium Methanearsonate

16.0%

#### INERT INGREDIENTS

84.0%

#### TOTAL

100.0%

Total Arsenic (as elemental) all in Water Soluble form 7.40%

Contains 1.5 lb. MSMA Per Gallon

### CAUTION: KEEP OUT OF REACH OF CHILDREN

#### STATEMENTS OF PRACTICAL TREATMENTS

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing. Wash affected area with soap and warm water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air. Get medical attention if indicated.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Avoid contact with skin. Do not breathe spray mist. Wear protective clothing when handling this product including long pants, long sleeved shirt and impermeable gloves and boots. Wash hands after using and before eating, drinking. Avoid contamination of feed and foodstuffs. Keep children and animals off treated areas until spray has dried or has been washed into the soil. Do not apply this product with a Hose End Applicator.

##### ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not feed treated forage to livestock or graze treated areas. Do not spray or allow to drift onto vegetables, ornamentals or other desirable plants. Do not apply when weather conditions favor drift from target areas. Do not apply directly to water.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. Do not store near fertilizer, seeds, insecticide or fungicides. **DISPOSAL:** Do not reuse empty container. Wrap and put in trash.

#### DIRECTIONS FOR USE (Cont.)

**General Information** — This product is recommended for selective, Post Emergent weed control in lawns and Ornamental Turf and Non-Crop areas as listed. Apply when temperatures are warm (70°F +) and weeds are actively growing. **Mixing** — Fill the spray tank 1/2 full and then thoroughly dissolve this product into the spray solution and fill the tank to proper level. Do not store solution in tank overnight. Clean application equipment after using by flushing with water. (See Storage & Disposal Statement). **How To Apply** — Use a hand pump sprayer, a garden type pressure sprayer, or sprinkler can. Do not use a sprayer that is used for the application of insecticides or fungicides due to possible contamination. Thoroughly wet all target weeds. Avoid wetting desirable turf grass where possible. A fan type nozzle will give best results. Apply on a calm day to avoid spray drift to susceptible plants.

**Weeds Controlled** — **Grassy Weeds:** Crabgrass, Dallisgrass, Sandbur, Goose Grass, Nutgrass (nut and type), Foxtail, Barnyard Grass, Bahiagrass, Johnson Grass. **Broadleaf Weeds:** Cocklebur, Puncture Vine, Chickweed, Wood Sorrel, and Pigweed.

#### ORNAMENTAL TURF GRASS

Apply as a post-emergent spray to control weeds listed herein. Do not use on new lawns until after the second cutting. Desirable turf grass may be temporarily discolored without lasting effects. Zoysia, Bluegrass, and Bermuda grass are quite tolerant. Do not use on St. Augustine or Centipede grass. Bent grass and Fescues may be damaged.

Mow the turf to 1 1/2 inches before treatment. **MIXING RATE** — Add 4 fluid ounces of this product to 3 gallons water to treat 1,000 sq. ft. (This is equal to 1 1/2 gallons of this product per acre.) Thoroughly wet all weeds. Plan on an additional application within 10 to 14 days to give desired control.

This spray is particularly effective for control of Crabgrass, Foxtail and Chickweed in the lawn. If possible treat weeds when in the seedling stage and before they go to seed. Best results will be obtained if the soil is moist. Small areas may be watered before application to increase the spray effectiveness.

#### NON CROP AREAS

This product is useful in the maintenance of drainage ditches, rights of way, fence rows, storage yards and similar areas to control weeds as listed herein. **MIXING RATE** — Add 4 fluid ounces in a minimum of 3 gallons of water and apply to 1,000 sq. ft. This is equal to 1 1/2 gallons of this product per acre. Make sure to wet plants thoroughly. Plan on a second treatment for best control.

Buyer assumes all risk and liability for use, storage and/or handling of this product not in accordance with the terms of this label.

PAR 130

**PRECAUTIONARY STATEMENTS**

**Hazards To Humans & Domestic Animals**

CAUTION: Do not eat, drink, or smoke while using this spray. Wash hands thoroughly after use. Do not get on clothing. Wear protective clothing, including pants, long sleeves, hat and gloves. Do not eat, drink, or smoke until after using and before eating or smoking. Avoid contact with skin until after spraying. Do not spray on treated areas until spray has dried. Do not wash treated areas until after spraying. Do not use on lawns or lawns treated with a Mistle-Seed Application.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water by leaking, spillage, or disposal of waste. Do not use this spray on lawns or lawns treated with a Mistle-Seed Application. Do not spray on lawns or lawns treated with a Mistle-Seed Application. Do not spray on lawns or lawns treated with a Mistle-Seed Application. Do not spray on lawns or lawns treated with a Mistle-Seed Application.

**STATEMENTS OF PRACTICAL TREATMENT**

If Swallowed: Drink plenty of water. Do not induce vomiting. If on skin: Wash with plenty of water. If in eyes: Flush with plenty of water. If inhaled: Move to fresh air. If on skin: Wash with plenty of water. If in eyes: Flush with plenty of water. If inhaled: Move to fresh air.

**DIRECTIONS FOR USE**

Use as directed on lawns and lawns treated with a Mistle-Seed Application.

**STORAGE AND DISPOSAL**

Store in original container. Do not use if container is damaged. Do not use if container is damaged. Do not use if container is damaged.

**DIRECTIONS FOR USE (cont.)**

General information: Do not use on lawns or lawns treated with a Mistle-Seed Application. Do not use on lawns or lawns treated with a Mistle-Seed Application.

KILLS CRABGRASS



WEED

**CZAR 529**

Herbicide — Contains MSMA

MSMA Liquid Plus Surfactant For Selective Post Emergent Weed Control  
 Contains 1.5 lb. MSMA Per Gallon  
**CONTROLS CRABGRASS, FOXTAIL, DALLIS GRASS, SAND BUR** and for Grassy Weeds such as Johnson Grass and Certain Broadleaf Weeds such as Cocklebur, Chickweed and Other Weeds as Listed.  
 One Pint Treats 4,000 Sq. Ft. of Turf

ACTIVE INGREDIENT  
 MSMA Liquid Plus Surfactant 1.5 lb. MSMA Per Gallon  
 Inert Ingredients 8.5 lb. MSMA Per Gallon  
 Total Weight 10.0 lb. MSMA Per Gallon

**CAUTION:**  
 KEEP OUT OF REACH OF  
 CHILDREN

See Side Panel For Additional Precautionary Statements

NOT REVIEWED  
 In Accordance with FR Notice 92-2,  
 Based on Draft Label: 92-2-1