



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

APR 13 1994

Christina M. Lawes  
ROUSSEL UCLAF CORP.  
95 Chestnut Ridge Road  
Montvale, NJ 07645416

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

SUBJECT: Label Amendment Submissions of 11/08/93 in Response to PR Notice 93-7  
EPA Reg. No.: SEE ATTACHED 10370-169


Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully **REMOVES THE PRODUCT FROM THE SCOPE** of PR Notice 93-7 once the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements), the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

Sincerely,

  
James Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Enclosures



Recycled/Recyclable  
Printed with Soy/Canola Ink on paper that  
contains at least 50% recycled fiber

194

ROUSSEL UCLAF CORP.  
10370 11/08/93

Original Submissions

- FORD'S LIQUID SEVIN 2L BRAND CARBARYL INSECTICIDE - 10370-32
- FORD'S DURSBAN 2.5% G - 10370-54
- FORD'S 50% MALATHION EC - 10370-147
- FORD'S CRABGRASS AND FOXTAIL KILLER - 10370-169
- LAWN FUNGICIDE GRANULES - 10370-202
- ORGANICIDE DIP - 10370-207
- CLEAN CROP SEVIN BRAND CARBARYL INSECTICIDE LIQUID - 10370-285

**BEST COPY AVAILABLE**

# ford's

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

JUN 30 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

10370-169

## C R A B G R A S S & F O X T A I L K I L L E R

### A MSMA LIQUID CONCENTRATE FOR SELECTIVE POST-EMERGENT WEED CONTROL

One Pint Treats 2,600 Sq.Ft. of Turf

\*\*\*\*\* Controls Crabgrass, Foxtail, Dallis Grass, Sand-Bur \*\*\*\*\*  
and For Grassy Weeds such as Johnson Grass and Certain  
Broadleaf Weeds such as Cocklebur, Chickweed and other  
weeds as listed.

ACTIVE INGREDIENT:	
Monosodium Methanearsonate . . . . .	11.50%
INERT INGREDIENTS . . . . .	88.50%
TOTAL	100.00%

Total Arsenic (As Elemental) all in Water Soluble form 5.32%  
Contains 1.0 lb. MSMA Per Gallon

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

#### STATEMENTS OF PRACTICAL TREATMENTS


- IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF IN EYES: Flush eyes with plenty of water. Get medical attention.
- IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and warm water. Get medical attention if irritation persists.
- IF INHALED: Remove victim to fresh air. Get medical attention if indicated.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 10370-169

EPA EST. NO.

NET CONTENTS

**ROUSSEL UCLAF**   
 Rousset Uclaf Corporation  
 95 Chestnut Ridge Road  
 Montvale, New Jersey 07645

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid contact with skin. Do not breathe spray mist. Wear protective clothing when handling this product including long pants, long sleeved shirt and impermeable gloves and boots. Wash hands after using and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep children and animals off treated areas until spray has dried or has been washed into the soil. Do not apply this product with a Hose-End Applicator.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not feed treated foliage to livestock or graze treated areas. Do not spray or allow to drift onto vegetables, ornamentals or other desirable plants. Do not apply when weather conditions favor drift from target areas. Do not apply directly to water.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original container in a cool, dry, locked place out of reach of children. Do not store near fertilizer, seeds, insecticide or fungicides. **DISPOSAL:** Do not re-use empty container. Wrap and put in trash.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**DIRECTIONS FOR USE(Cont.)**

**General Information** — This product is recommended for selective "Post-Emergent" weed control in Lawns and Ornamental Turf and Non-Crop areas as listed. Apply when temperatures are warm (70° F+) and weeds are actively growing.

**Mixing** — Fill the spray tank 1/2 full and then thoroughly dissolve this product into the spray solution, then fill tank to proper level. Do not store solution in tank over night. Clean application equipment after using by flushing with water. (See "Storage & Disposal" Statement).

**How To Apply** — Use a hand pump sprayer, a garden type pressure sprayer, or sprinkler can. Do not use a sprayer that is used for the application of insecticides or fungicides due to possible contamination. Thoroughly wet all target weeds. Avoid wetting desirable turf grass where possible. A fan type nozzle will give best results. Apply on a calm day to avoid spray drift to susceptible plants.

**Weeds Controlled** — Grassy Weeds; Crabgrass, Dallisgrass, Sandbur, Goose grass, Nutgrass (Nutsedge), Foxtail, Barnyard Grass, Bahiagrass, Johnson grass, Broadleaf Weeds; Cocklebur, Puncture Vine, Chickweed, Wood Sorrel, and Pigweed.

**ORNAMENTAL TURF GRASS**

Apply as a post-emergent spray to control weeds listed hereon. Do not use on new lawns until after the second cutting. Desirable turf grass may be temporarily discolored without lasting effects. Zoysia, Bluegrass, and Bermuda grass are quite tolerant. Do not use on St. Augustine or Centipede grass. Bent grass and Fescues may be damaged.

Mow the turf to 1 1/2 inches before treatment. **MIXING RATE** — Add 8 fluid ounces of this product to 3 gallons water to treat 1,000 sq. ft. (This is equal to 2 gallons of this product per acre.) Thoroughly wet all weeds. Plan on an additional application within 10 to 14 days to give desired control.

This spray is particularly effective for control of Crabgrass, Foxtail and Chickweed in the lawn. If possible treat weeds when in the seedling stage and before they go to seed. Best results will be obtained if the soil is moist. Small areas may be watered before application to increase the spray effectiveness.

**NON-CROP AREAS**

This product is useful in the maintenance of drainage ditches, rights-of-way, fence rows, storage yards and similar areas to control weeds as listed hereon. **MIXING RATE** — Add 8 fluid ounces in a minimum of 3 gallons of water and apply to 1,000 sq. ft. This is equal to 2 gallons of this product per acre. Make sure to wet plants thoroughly. Plan on a second treatment for best control.

Buyer assumes all risk and liability for use, storage and/or handling of this product not in accordance with the terms of this label.