

5418846
18
300
CD

Richard B. Boyd, Jr.
Ford's Chemical & Service, Inc.
3741 Red Bluff Rd., Suite 200
Pasadena, TX 77503

MAR 30 1993

Subject: Revised Labeling Submitted on 12/20/91
EPA File Number: 10370-07

Dear Mr. Boyd:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments below.

1. Within the Environmental Hazards Statements, change "Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes)." to "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark." In addition, change the statement "Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources." to "Do not contaminate water when disposing of equipment washwaters."

2. In the Storage and Disposal section, within a Container Disposal subheading change the statement to, "Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash."

Five copies of the finished labeling must be submitted prior to releasing the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Refer to the "A-79 Enclosure" regarding finished labeling. A stamped copy is enclosed for your records.

Sincerely,



Phil Hutton
Product Manager (18)

CONSUMER Pesticide-Rodenticide Branch

| | | | | | | | |
|---------|--------|--|--|--|--|--|--|
| SYMBOL | H7505C | | | | | | |
| SURNAME | CDP | | | | | | |
| DATE | 3/2/93 | | | | | | |

Office of Pesticide Programs



295

ORGANICIDE DIP

Contains Rotenone & Pyrethrins
For Dilution with Water

FOR FLEAS & TICKS ON PETS

8 Ounces Can Make up to 16 Gallons of Dip, Wash or Spray

Can Be Used on Your Pet as a Spray, Wash or Dip
Kills Fleas, Ticks and Lice on Dogs or Cats

May be Used as a Spray in Pet Living and Sleeping Quarters

For Insect Control on Ornamentals — Fruit — Vegetables
One 8 Ounce Bottle Makes up to 48 Gallons of Finished Spray


This product Will Kill and Control 43 Common Insect Pests That Attack
Ornamentals, Shrubs & Flowers, Vegetables, Fruits & Berries

ACTIVE INGREDIENTS:

| | |
|-------------------------------|------|
| Rotenone | 1.1% |
| Other Associated Resins | 2.2% |
| Pyrethrins | 0.8% |

| | |
|--------------------------|--------|
| INERT INGREDIENTS* | 95.9% |
| Total | 100.0% |

*This product contains an aromatic petroleum solvent.



[PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.]

Keep Out of Reach of Children

WARNING


See side [back] panel for additional precautionary statements.

EPA Reg. No. 10370-207

EPA Est. 10370-TX-1

RECEIVED
with COMMENTS
in EPA Letter Dated

Net Contents
(8 ounces, 1 Pint, 1 Quart or 1 Gallon)



March 30, 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
registered for the pesticide
under EPA Reg. No.
10370-207

FORD'S CHEMICAL & SERVICE, INC.
3741 Red Bluff, Suite 200
Pasadena, TX 77503

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: May be fatal if swallowed. Harmful if absorbed through the skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. provide adequate ventilation in treatment area. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse. Avoid contamination of food, utensils and food preparation areas. **NOTE TO VETERINARIANS:** If animal swallows significant amount of dip, administer an emetic, or gastric lavage if not convulsive; assist respiration as necessary. Deep IM or slow IV drip of diazepam to control convulsions. Calcium gluconate IV and Vitamin B should be administered to preserve liver functions.

Statement of Practical Treatment

If Swallowed: Call a physician or poison control center immediately. Promptly drink large quantities of milk, egg whites, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting. Aromatic solvent content may present aspiration hazard. **If Inhaled:** Remove victim to fresh air. Apply artificial respiration if needed. **If in Eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists. **If On Skin:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store product in its original labeled container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not reuse empty container. Wrap container and place in trash collection.

KILLS FLEAS, TICKS AND LICE ON DOGS AND CATS AND IN THEIR LIVING QUARTERS

IMPORTANT

Avoid spray or dip contact with pet's eyes, nose or mouth. Treat pet's ears for ticks. To prevent reinfestation, thoroughly spray the animal's living quarters and bedding with 2 fluid ounces per 1 gallon of water. Do not use 2 ounce dilution rate directly on cats. After treatment place freshly laundered bedding in sleeping quarters. Repeat as needed for continued control through "flea" season.

HOW TO APPLY TO YOUR PET FOR USE ON DOGS

While wearing neoprene or rubber household gloves spray or sponge in against the lay of pet hair until liquid penetrates to skin; or, dip the entire pet's body making sure not to get head or face into dipping solution. Make sure to treat animal's back, head to tail, around the neck and ears, the underbelly, the legs, the base of the tail and where Fleas, Ticks or Lice are seen. Do not get solution into pet's eyes, nose or mouth. Afterwards, do not rinse pet since treatment will be ineffective.

FOR PUPPIES

Dilute 1/2 fluid ounce (1 tbsp.) of concentrate in 1 gallon of water for moderate Flea or Tick infestations; or, 1 fluid ounce (2 tbsp.) in 1 gallon of water for heavy infestations. Do not exceed the 1 ounce per gallon rate on puppies. See "HOW TO APPLY TO YOUR PET." Do not use on puppies less than 4 weeks old.

FOR ADULT DOGS

FLEAS AND LICE: Dilute 1 fluid ounce (2 tbsp.) per 1 gallon of water. For young adults use 1/2 fluid ounce per 1 gallon of water. Apply directly to animals as a spray, or as a sponge-on wash; or, dip animals until animal's skin is wet. See "HOW TO APPLY TO YOUR PET."

TICKS: Dilute at the rate of 2 fluid ounces (4 tbsp.) per 1 gallon of water. For young animals use 1 fluid ounce per 1 gallon of water. Apply directly to the animals as a spray or as a sponge-on wash; or dip until animal's skin is wet. For Ticks, examine pet's ears and treat as needed. See "HOW TO APPLY TO YOUR PET."

FOR USE ON CATS

HOW TO APPLY TO YOUR CAT

While wearing neoprene or rubber household gloves spray or sponge in against the lay of pet fur until liquid penetrates to skin; or dip the entire pet's body making sure not to get head or face into dipping solution. Make sure to treat animal's back, head to tail, around the neck and ears, the underbelly, the legs, the base of the tail and where Fleas, Ticks or Lice are seen. Do not get solution into pet's eyes, nose or mouth. Afterward, do not rinse pet since treatment will be ineffective.

FOR ADULT CATS AND KITTENS

Do not use on kittens less than four weeks old. **FLEAS, TICKS and LICE:** dilute at the rate of 1/2 ounce (1 tbsp.) per 1 gallon of water. Apply directly to animal as a spray or as a sponge-on wash. Dipping may be difficult.

Liquid Measuring Conversion:

1. 1/2 oz = 1 tablespoonful per gallon = 15 milliliter per 3.79 Liters
2. 1 oz = 2 tablespoonful per gallon = 30 milliliter per 3.79 Liters
3. 2 oz = 4 tablespoonful per gallon = 60 milliliter per 3.79 Liters

FOR USE ON ORNAMENTAL TREES, SHRUBS AND FLOWERING PLANTS; FOR VEGETABLES; FOR FRUITS AND BERRIES

Mix this product with water as described in the following charts. Use a hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Repeat every 5 to 10 days or as needed. Mix only the amount of spray needed for one complete spray coverage. Do not feed treated foliage to livestock. Do not apply rotenone in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried or dusts have settled.

ORNAMENTALS

For use on Ornamental Trees, Shrubs and Flowering plants such as Poplar, Linden, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Soirea, Rose, Lilac, Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation and Chrysanthemum. Do not apply this product within one day of collecting flowers or cuttings.

PESTS OF ORNAMENTALS

AMOUNT TO USE PER 1 GALLON WATER

Aphids, Leafhoppers, Exposed Thrips

1 teaspoonful

Oblique-Banded Leaf Rollers, Sawfly Larvae, Boxelder Bugs, Japanese Beetles, Elm Leaf Beetle Larvae, Mites, Lygus Bugs, Lace Bugs, Greenhouse Whiteflies, Stink Bugs, Garden Fleahoppers, Greenhouse Leafhoppers

2 teaspoonfuls

Cucumber Beetles, Rose Chafer, Ants, Aster Beetles, Flea Beetles

4 teaspoonfuls

VEGETABLES

For use on Vegetable Plants such as Peas, Beans, Beets, Cucumbers, Squash, Muskmelons, Potato, Cabbage, Spinach, Lettuce, Celery, Asparagus, Pepper, Eggplant, Onions, Carrots and Corn. Do not apply within one day of harvest.

| VEGETABLE PESTS | AMOUNT TO USE PER 1 GALLON WATER |
|--|-------------------------------------|
| Aphids, Leafhoppers, Exposed Thrips | 1 teaspoonful |
| Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetle, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leaf-tiers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites | 2 teaspoonfuls |
| Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (Nymphs), Squash Vine Borer, [European Corn Borers; 3 to 4 complete stalk and leaf applications or more; as eggs start to hatch. Repeat at 5 to 7 day intervals.] Consult your Agricultural Extension Agent for Details. | 4 teaspoonfuls |

FRUITS & BERRIES

For use on Apples, Pears, Peaches, Plums, Cherries, Grapes, Currants, Gooseberry, Raspberry, Blackberry, Strawberry and Blueberry, consult the following chart. Do not apply within one day of harvest.

| PESTS ON FRUITS & BERRIES | AMOUNT TO USE PER 1 GALLON WATER |
|--|-------------------------------------|
| Aphids, Leafhoppers, Exposed Thrips | 1 teaspoonful |
| Imported Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms - adult, Red-necked Borers, Japanese Beetles and Mites | 2 teaspoonfuls |
| Strawberry Leafrollers and Rose Chafers | 4 teaspoonfuls |

Liquid Measuring Conversion:

1. 1/2 oz = 1 tablespoonful per gallon = 15 milliliter per 3.79 Liters
2. 1 oz = 2 tablespoonful per gallon = 30 milliliter per 3.79 Liters
3. 2 oz = 4 tablespoonful per gallon = 60 milliliter per 3.79 Liters

NOTICE: Buyer and user assume all risk of liability of use, storage and/or handling of this product not in accordance with the terms of this label.

