



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 13 1994

Christina M. Lawes
ROUSSEL UCLAF CORP.
95 Chestnut Ridge Road
Montvale, NJ 07645416

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SUBJECT: Label Amendment Submissions of 11/08/93 in Response to PR Notice 93-7
EPA Reg. No.: SEE ATTACHED 10370-207


Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully **REMOVES THE PRODUCT FROM THE SCOPE** of PR Notice 93-7 once the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements), the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

Sincerely,


James Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Enclosures



Recycled/Recyclable
Printed with SoyCandle Ink on paper that
contains at least 50% recycled fiber

194

ROUSSEL UCLAF CORP.
10370 11/08/93

Original Submissions

FORD'S LIQUID SEVIN 2L BRAND CARBARYL INSECTICIDE - 10370-32
FORD'S DURSBAN 2.5% G - 10370-54
FORD'S 50% MALATHION EC - 10370-147
FORD'S CRABGRASS AND FOXTAIL KILLER - 10370-169
LAWN FUNGICIDE GRANULES - 10370-202
ORGANICIDE DIP - 10370-207
CLEAN CROP SEVIN BRAND CARBARYL INSECTICIDE LIQUID - 10370-285

Ford's

ORGANICIDE™ DIP

- * CONTAINS ROTENONE & PYRETHRINS
- * FOR DILUTION WITH WATER
- * FOR FLEAS & TICKS ON PETS
- * 8 OUNCES CAN MAKE UP TO 16 GALLONS OF DIP, WASH OR SPRAY
- * CAN BE USED ON YOUR PET AS A SPRAY, WASH OR DIP
- * KILLS FLEAS, TICKS AND LICE ON DOGS
- * MAY BE USED AS A SPRAY IN PET LIVING AND SLEEPING QUARTERS
- * FOR INSECT CONTROL ON ORNAMENTALS - FRUIT - VEGETABLES
- * ONE 8 OUNCE BOTTLE MAKES UP TO 48 GALLONS OF FINISHED SPRAY
- * THIS PRODUCT WILL KILL AND CONTROL 43 COMMON INSECT PESTS THAT ATTACK ORNAMENTALS, SHRUBS & FLOWERS, VEGETABLES, FRUITS & BERRIES

ACTIVE INGREDIENTS:

Rotenone	1.1%
Other Associated Resins	2.2%
Pyrethrins	0.8%
INERT INGREDIENTS*	95.9%
	100.0%

*This product contains an aromatic petroleum solvent.

™Organicide is a trademark of Roussel Uclaf Corporation.

[PRECAUCION AL USUARIO: Si usted no lee ingles, no use producto hasta que la etiqueta haya sido explicado ampliamente.]

EPA REG. NO.10370-207

EPA EST.NO.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center immediately. Promptly drink large quantities of milk, egg whites, gelatin solution, or, if these are not available, large quantities of water. Avoid Alcohol. Do not induce vomiting. Aromatic solvent content may present aspiration hazard.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if needed.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

See Side Panel for Additional

Precautionary Statements

NET CONTENTS

ROUSSEL UCLAF

Roussel Uclaf Corporation

95 Chestnut Ridge Road

Montvale, New Jersey 07645

(207) 081993tk

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUN 30 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10370-267

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

WARNING

May be fatal if swallowed. Harmful if absorbed through the skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Provide adequate ventilation in treatment area. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse. Avoid contamination of food, utensils and food preparation areas. NOTE TO VETERINARIANS: If animal swallows significant amount of dip, administer an emetic, or gastric lavage if not convulsive; assist respiration as necessary. Deep IM or slow IV drip of diazepam to control convulsions. Calcium gluconate IV and Vitamin B should be administered to preserve liver functions.

Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwater.

Physical and Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store product in its original container in a cool, dry, locked place out of reach of children.

Disposal: Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes, or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

KILLS FLEAS, TICKS, AND LICE, ON DOGS AND IN THEIR LIVING QUARTERS.

IMPORTANT: Avoid spray or dip contact with pet's eyes, nose or mouth. Treat pet's ears for ticks. To prevent reinfestation, thoroughly spray the animal's living quarters and bedding with 2 fluid ounces per 1 gallon of water. After treatment, place freshly laundered bedding in sleeping quarters. Repeat as needed for continued control through "flea" season.

HOW TO APPLY TO YOUR PET

FOR USE ON DOGS: While wearing neoprene or rubber household gloves spray or sponge in against the lay of pet hair until liquid penetrates to skin; or dip the entire pet's body making sure not to get head or face into dipping solution. Make sure to treat animal's back, head to tail, around neck and ears, the underbelly, the legs, the base of the tail and where Fleas, Ticks or Lice are seen. Do not get solution into pet's eyes, nose or mouth. Afterwards, do not rinse pet since treatment will be ineffective.

FOR PUPPIES: Dilute 1/2 fluid ounce (1 tbsp.) of concentrate in 1 gallon of water for moderate Flea or Tick infestations; or, 1 fluid ounce (2 tbsp.) in 1 gallon of water for heavy infestations. Do not exceed the 1 ounce per gallon rate on puppies. See "How

To Apply To Your Pet." Do not use on puppies less than 4 weeks old.

FOR ADULT DOGS: *Fleas and Lice*: Dilute 1 fluid ounce (2 tbsp.) per 1 gallon of water. For young adults use 1/2 fluid ounce per 1 gallon of water. Apply directly to animals as a spray, or as a sponge-on wash; or, dip animals until animal's skin is wet. See "How To Apply To Your Pet."

TICKS: Dilute at the rate of 2 fluid ounces (4 tbsp.) per 1 gallon of water. For young animals use 1 fluid ounce per 1 gallon of water. Apply directly to the animals as a spray or as a sponge-on wash; or dip until animal's skin is wet. For ticks, examine pet's ears and treat as needed. See "HOW TO APPLY TO YOUR PET."

Liquid measure conversion:

1. 1/2 oz = 1 tablespoonful per gallon = 15 milliliter per 3.79 Liters
2. 1 oz = 2 tablespoonful per gallon = 30 milliliter per 3.79 Liters
3. 2 oz = 4 tablespoonful per gallon = 60 milliliter per 3.79 Liters

FOR USE ON ORNAMENTAL TREES, SHRUBS AND FLOWERING PLANTS; FOR VEGETABLES; FOR FRUITS AND BERRIES Mix this product with water as described in the following charts. Use a hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Repeat every 5 to 10 days as needed. Mix only the amount of spray needed for one complete spray coverage. Do not feed treated foliage to livestock. Do not apply rotenone in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried or dusts have settled.

ORNAMENTALS

For use on ornamental trees, shrubs and flowering plants such as Poplar, Linden, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Spirea, Rose, Lilac, Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation and Chrysanthemum. Do not apply this product within one day of collecting flowers or cuttings.

Pests of Ornamentals	Amount to Use Per 1 Gallon Water
Aphids, Leafhoppers, Exposed Thrips	1 teaspoon
Oblique-Banded Leaf Rollers, Sawfly Larvae, Boxelder Bugs, Japanese Beetles, Elm leaf Beetle Larvae, Mites, Lygus Bugs, Lace Bugs, Greenhouse Whiteflies, Stink Bugs, Garden Fleahoppers, Greenhouse Leafhoppers	2 teaspoons
Cucumber Beetles, Rose Chafer, Ants, Aster Beetles, Flea Beetles	4 teaspoons

VEGETABLES

For use on Vegetable Plants such as Peas, Beans, Beets, Cucumbers, Squash, Muskmelons, Potato, Cabbage, Spinach, Lettuce, Celery, Asparagus, Pepper, Eggplant, Onions, Carrots and Corn. Do not apply within one day of harvest.

Vegetable Pests

Amount to Use Per
1 Gallon Water

Aphids, Leafhoppers, Exposed Thrips	1 teaspoon
Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetle, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leafhoppers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites	2 teaspoons
Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (Nymphs), Squash Vine Borer, [European Corn Borers; 3 to 4 complete stalk and leaf applications or more; as eggs start to hatch. Repeat at 5 to 7 day intervals.] Consult your Agricultural Extension Agent for details.	4 teaspoons

FRUITS AND BERRIES

For use on Apples, Pears, Peaches, Plums, Cherries, Grapes, Currants, Gooseberry, Raspberry, Blackberry, Strawberry and Blueberry, consult the following chart. Do not apply within one day of harvest.

Pests on Fruits and Berries	Amount of use per 1 Gallon Water.
Aphids, Leafhoppers, Exposed Thrips	1 teaspoon
Imported Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms - adult, Red-necked Borers, Japanese Beetles and Mites	2 teaspoons
Strawberry Leafrollers and Rose Chafers	4 teaspoons

Liquid measuring Conversion:

1. 1/2 oz=1 tablespoonful per gallon=15 milliliter per 3.79 Liters
2. 1 oz=2 tablespoonful per gallon=30 milliliter per 3.79 Liters
3. 2 oz=4 tablespoonful per gallon=60 milliliter per 3.79 Liters

NOTICE: Buyer assumes all risk of liability of use, storage and/or handling of this product not in accordance with the terms of this label.