

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Ford's Chemical & Service, Inc.
2739 Pasadena Boulevard
Pasadena, CA 77502

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 10370-210."
 - b. Change the statement "Do not contaminate water by cleaning of equipment or disposal of wastes" to read "Do not contaminate water when disposing of equipment washwaters."
3. A 1-year storage stability study must be submitted. The formulated product must be analyzed for its active ingredient at time zero and during a year of storage. These data must be submitted within 15 months from the date of this Registration Notice.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

8/23/88

95

10/10

Compound Evaluation Laboratory
Scientific Services, FSQS
Food Ingredient Assessment Division
U.S. Department of Agriculture
Building 306, BARC-East
Beltsville, MD 20705

5. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Phil Hutton
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

26/10

ACCEPTED
MAY 19 1988
EPA REGION 1
BOSTON, MASS.

3



**FORD'S PPT NO. 1515
INSECT SPRAY**

"PYRETHRINS / PIPERONYL BUTOXIDE TRANSPARENT EMULSION" 10370-210

An Insecticide for Control of House Flies, Mosquitoes, Gnats(Midges), Wasps, Small Flying Moths(Millers), Cockroaches(Waterbugs), Ants, Silverfish, Crickets, Fleas, Bedbugs, Lice, Earwigs, Centipedes, Sowbugs, Boxelder Bugs, Horn Flies, Face Flies, Deer Flies, Stable Flies, Sucking and Biting Lice, Ticks, Sheep Keds, and Poultry Mites.

Product May Be Used Full Strength Or Diluted To Make Up To 16 Gallons of Finished Spray

ACTIVE INGREDIENTS:

Pyrethrins	0.15%
*Piperonyl Butoxide Technical	1.50%
INERT INGREDIENTS:	98.35%
Total	100.00%

*Equivalent to 1.2% (Butylcarbityl)(6-propylpiperonyl) Ether and 0.3% related compounds.
This Product is a Water Base Insecticide Concentrate.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Have victim drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. Thoroughly rinse. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 10370-___

EPA EST NO. 10370-TX-1

NET CONTENTS __1__ GALLON

FORDS CHEMICAL & SERVICE, INC.
2739 PASADENA BLVD.
PASADENA TEXAS 77502

3c/16

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating or smoking.

Do not apply directly to food or feedstuffs. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply when food processing is underway. After space spraying in bakeries and other food processing plants wash thoroughly all equipment, benches, shelving, etc, where exposed food will be handled, with an effective cleaning compound and then rinse with potable water. In the home all food processing surfaces and utensil, should be covered during treatment, or thoroughly washed before use. Remove pets and cover fish aquariums before space spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS

Do not expose this water base product to freezing temperatures. Do not apply in electrical conduits, motor housings, junctions, and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be stained or harmed by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. Do not store in temperatures below 35°F.
PESTICIDE DISPOSAL: For Containers Over 1 Gallon Size - Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility.
CONTAINER DISPOSAL: For Containers 1 Gallon And Under; Do not reuse empty container. Wrap and put in trash. For Containers over 1 Gallon; Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture & dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities.

HOW TO MIX

Use this product undiluted or mixed with water. See SPRAY DILUTION CHART.

- | | |
|----------------------------------|---|
| Use Full Strength Product As Is | * For reduction of heavy infestations of flying and crawling insects as listed indoors and outdoors. |
| Use Dilution No.1 (0.1% + 1.0%) | * For Light Infestations of Flying and Crawling Insects.
* For an Indoor Maintenance Insect Control Program.
* To Kill and Repel flying insects as listed on Animals. |
| Use Dilution No.2 (0.01% + 0.1%) | * For Control of Lice, Fleas, Ticks, Sheep Ked on Animals.
* For Use on Poultry for Bed Bugs & Mites. |

4/6/16

NOTE TO APPLICATORS: ONE GALLON (3.79 L) OF FINISHED SPRAY IS SUFFICIENT TO TREAT 1500 LINEAR FEET (457 METERS) WITH A 6 INCH (15 CM) SPRAY BAND.

SPRAY DILUTION CHART

AMOUNT OF FINISHED SPRAY MIXTURE DESIRED		ADD THE FOLLOWING AMOUNTS OF THIS PRODUCT TO THE GALLONS OR LITERS SHOWN IN COLUMN NO. 1 OR 2	
1) GALLONS	2) LITERS	3) FOR DILUTION "NO. 1" (0.1% + 1.0%)	4) FOR DILUTION "NO. 2" (0.01% + 0.1%)
1	3.79	2-3/4 qt = 2.60 L	9 fl oz = 266 ml
3	11.37	2 gallon = 7.57 L	26 fl oz = 769 ml
5	18.95	3-1/3 g = 12.60 L	1-2/5 qt = 1.32 L
10	37.90	6-3/4 g = 25.55 L	2-3/4 qt = 2.60 L

Note: 1 Fluid Ounce = 2 Tablespoonfuls = 6 Teaspoonfuls = 29.574 ml
L = Liter; ml = Milliliter; qt. = quart; g = gallon; oz = ounce

INDOOR PEST CONTROL

WHERE TO USE

For use in Hotels, Motels, Homes, Pre-schools, Apartment Buildings, Commercial Offices, Institutional and Storage Buildings, Recreational Buildings, Packaging and Processing Plants, Milk Handling Areas of Milk Plants, Milk Storage Rooms, Egg Processing Plants, Supermarkets, Restaurants, Bottling Plants, Canneries, Flour and Feed Mills, Bakeries, Boats, Ship Cabins, and Hospitals, In Hospitals, patient should be removed prior to treatment. Also for use in Campers, Mobile Homes, Buses, Automobiles, Trucks, Boxcars, and Ship Cargo Holds, Also for use Barns, Stables, Kennels, Poultry Houses, and Other Animal Quarters. When Spraying in Animal Quarters.

DO NOT SPRAY FOOD OR FEED ITEMS.

DO NOT APPLY OR ALLOW TO CONTACT PLANT FOLIAGE BECAUSE OF POSSIBLE INJURY.

HOW TO USE AS A SPACE SPRAY

To control House Flies, Mosquitoes, Gnats, Wasps, & Small Flying Moths close all doors, windows, and ventilators. Use Full Strength or Dilution No. 1. Adjust spray equipment to deliver a fine mist and direct the nozzle upward and outward in all directions. Fill the room with mist at the rate of 0.5 to 1 ounce of finished spray for each 1,000 Cu. Ft. (28.3 Cu M) of space. Keep the area closed for 15 minutes, then thoroughly ventilate before reentering the treated area. Sweep up and dispose of dead insects.

HOW TO USE AS A CONTACT SPRAY

Use Full Strength to kill Cockroaches (Waterbugs), Fleas, Earwigs, Centipedes, Sowbugs, Boxelder Bugs, Ants, Silverfish, and crickets; adjust spray nozzle to product large wetting type spray particles. Direct spray to contact insects in cracks and crevices in floors, baseboards, around pipes, behind cabinets, and other hiding places. Spray as many insects directly as possible. Apply the spray to the point of run-off. Repeat as necessary.

TO KILL BEDBUGS AND LICE

Apply Full Strength as a spot treatment to cracks and crevices, around baseboards, floorboards, bedboards, and walls. First, thoroughly clean and air mattresses, and springs. Then spot treat only areas that may harbor pests, concentrating on tufts, folds, and seams, Repeat as necessary. For persistent infestations, consult a Professional Applicator.

50/10

OUTDOOR PEST CONTROL

DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (LAWNS, FLOWERING PLANTS, SHRUBS, TREES, EVERGREENS ETC.) BECAUSE OF POSSIBLE INJURY.

HOUSEFLIES, MOSQUITOES, GNATS, FLYING MOTHS, AND WASPS OUTDOORS

For outdoor use as an aid in reducing annoyance from these insect in Municipalities, Zoos, Parks, Playgrounds, Stadiums, Open-air Theaters, Recreational Areas, Dump Sites, Manure & Compost Piles. Use Full Strength and treat the area with 1.5 gallons per acre (1.3 L/ha). Spray Garbage Cans, Cuthouses, Barbecue Pits, Compost and Manure Piles and other breeding areas until surfaces are thoroughly wet. Spray outside surfaces of Tents, Lean-tos, and Shelters in Camping Areas, the outside of Barns, Dairies, Kennels, Creameries, Food Processing Plants, and Institutions until surfaces are thoroughly wet, paying particular attention to screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around outdoor light fixtures on porches, garages, and other places where these insect alight or congregate. Repeat as necessary.

INSECT CONTROL ON DOMESTIC ANIMALS

DAIRY & BEEF CATTLE - HORSES - HOGS - SHEEP - GOATS - DOGS

To kill and repel House Flies, Horn Flies, Face Flies, Deer Flies, Stable Flies and Horse Flies, on above listed animals Use Dilution No. 1 and apply as a coarse particle spray at the rate of 2 ounces per large animal on a daily basis. Apply sufficiently to wet the hair but not to soak the hide.

For longer insect kill and repellent action use up to 1 to 2 quarts (0.95 to 1.89 L) thoroughly applied to an animal the size of a mature cow. This latter treatment will protect large animals up to 3 to 8 days from Stable Flies, Horn Flies, Face Flies and House Flies, and offers up to 3 days protection from Horse Flies and Deer Flies. Avoid spraying in eyes. Repeat as necessary.

Any animals nervous from the sound or feel of the spray can be treated by applying the material with a sponge. Applicator should wear protective clothing and gloves for sponging.

For control of Sucking and Biting Lice, Fleas, Ticks and Sheep Ked on the above listed animals use Dilution No. 2 as a dip or spray. If used as a spray, make sure the animal hair is thoroughly wet including the head and tail. Repeat the treatment in ten days to kill newly hatched insect.

FOR POULTRY

For Biting Lice, Fleas, Ticks, Bed Bugs, and Mites on Poultry: Use Dilution No. 2 to treat birds by dipping them in the solution while holding the wings away from the body to insure thorough coverage. To avoid chilling of the poultry, dip on a warm day, or in a warm building. Applicator should wear protective clothing and gloves.

FOR POULTRY HOUSES

Apply Full Strength as a crack and crevice/spot treatment with a coarse spray droplet to areas where the Bed Bugs, and mites hide, such as cracks in walls and floor, crevices, roost poles and nests.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

6/16