ريم

ca the s	<u> </u>	
US ENVIRONMENTAL PROTECTION SENCY OFFICE OF PESTICIDES PROGRAMS	10370-226	JUL 12 1900
REGISTRATION DIVISION (75-767) WASHINGTON, DC 20480	TERM OF ISSUANCE	1000
MATION OF DESTICINE THE REGISTRATION	NAME OF PESTICIDE PRO	DUCT
NOTICE OF PESTICIDE: REGISTRATION	figet Garain	foray
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	<u> </u>	
NAME AND ADDRESS OF REGISTRANT (Include 21P code)		1/ - 1.
_	٦	160 207017
Fig. Commission Courts , Inc.		
2759 Fri (d) 38 Fr (4, 45 a)		18
Papadonu, TX 77502		. 0
ı	1	•
_	-	
NOTE: Changes in labeling formula differing in substance fr submitted to and accepted by the Registration Division prior	to use of the label in comm	erce. In any correspondence on this
product always refer to the above U.S. EPA registration num	ber.	
On the besis of information furnished by the registrant, the substitute formation for the federal insecticide, Fungicide, and Rodenticide Act.	bove named pesticide is he	reby Registered/Reregistered under
A copy of the labeling accepted in connection with this Reg		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
Incomposed to condity maily reging recondition (c) (d) (A) provided that you:	stareu en accesdanc	c with FTFRA
1. Submit/cit/ all data required	for remains take on the	eredis cration of
your product under FIFRA section 3(c)(5		
registrance of similarepreducts similar	-products to submit	to such adatabasement asset (*)
2. Make the labeling changes list	ad hellow to form you	u velosou the
product for shipment:	bu butow herore you	d release the
	1	
a. Add the phrase "EPA Regist	ration No. 1370-220	6. "
n. Delete the statements "To rate of 7.75 gallons per a feet. Apply as a coarse w shallow waters of wetlands larvae are known to breed.	ore or .19 gallons etting spray to the , swamps and march	per 1000 square border of, or ch
c. Place the signal word "Cau Children" statement.	tion" <u>under</u> the "Ke	nep Out of Re ach (f
ATTACHMENT IS APPLICABLE	BEST AVA	MLABLE COPY

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED

- d. Add the statement "If irritation persists, get medical attention" after the statement "If On Skin: Wash with plenty of soap and water."
- 3. Submit a 1-year storage stability study. The formulated product must be analyzed for its active ingredient at time zero and during a year of storage. The storage should be in warehouse conditions of temperature and humidity and stored in similar containers you will be using the trade. Note: For the storage stability study you cannot reference the concentrate you are using to formulate your product.
- 4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the λ -79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFEA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.


Phil Mutton

Product Manager (17)

Insecticide-Rodenticide Branch

Registration Division (H7505C)

Enclosures



FORD'S GARDEN SPRAY

Kills Gypsy Moth Caterpillars and Japanese Beetles
An Effective Insecticide for Use on
Vegetable and Ornamental Pests
A Concentrated Pyrenones Micro-Emulsion, Clear, Stable

Apply With Conventional Mechanical Compressed Air Equipment or Non-Aerosol Pump

ACTIVE INGREDIENTS:

Pyrethrins	
	rechnical
•	
	Total100.00%

*Equivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

DO NOT TRANSPORT OR STORE UNDER 32°
If Freezing Occurs, Thaw and Agitate Vigorously

KEEP OUT OF REACH OF CHILDREN - C A U T I O N

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Calling physician or Poison Control Center: immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person. IF IN EYES: Flush eyes with plenty of water. If irritation persists, get medical attention. IF ON SKIN: Wash with plenty of soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estauries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

FORD'S CHEMICAL & SERVICE, INC.

EPA REG. NO. 10370 - XXX EPA EST. NO. 10370-TX-1

DIRECTIONS FOR USE

(

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach, and Tomatoes. To control: asparagus beetle, aphids, leafhoppers, whiteflies, Mexican bean beetles, 12-spotted cucumber beetles, cabbage loopers, cross-striped cabbageworms, diamondback moth larvae, flea beetles, harlequis, bugs, imported cabbageworms, stink bugs, green peach aphids, leaftiers, fireworms, blister beetles, Colorado potato beetles and webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jews, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolis and Marigolds - To control: aphids, flea beetles, leafhoppers and whiteflies.

To kill gypsy moth caterpillars and adults, adult Japanese

beetles and Eastern and forest tent caterpillars on contact:
On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats, and small flying moths: Apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or .18 gallons per 1000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

with COMMENTS in EPA Unter Detect

JUL 12 1989

Under the residence de la Función de la Companya de