


FORD'S GARDEN SPRAY

Kills Gypsy Moth Caterpillars and Japanese Beetles
 An Effective Insecticide for Use on
 Vegetable and Ornamental Pests
 A Concentrated Pyrenone^a Micro-Emulsion, Clear,



Apply With Conventional Mechanical Compressed Air
 Equipment or Non-Aerosol Pump

ACTIVE INGREDIENTS:

Pyrethrins.....	0.02%
*Piperonyl Butoxide, Technical.....	0.20%
INERT INGREDIENTS:.....	99.78%
Total.....	100.00%

^aEquivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds.

DO NOT TRANSPORT OR STORE UNDER 32°
 If Freezing Occurs, Thaw and Agitate Vigorously

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person. IF IN EYES: Flush eyes with plenty of water. If irritation persists, get medical attention. IF ON SKIN: Wash with plenty of soap and water. If irritation persists, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing.

DISPOSAL: Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

Or triple rinse dispose of container with recyclables where local law permits.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct the spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach, and Tomatoes. To control: asparagus beetles, aphids, leafhoppers, whiteflies, Mexican bean beetles, 12-spotted cucumber beetles, cabbage loopers, cross-striped cabbageworms, diamondback moth larvae, flea beetles, harlequin bugs, imported cabbageworms, stink bugs, green peach aphids, leaf tiers, fireworms, blister beetles, Colorado potato beetles and webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jews, Asters, Carnations, Chrysanthemums, Geraniums, Gladioli and Marigolds - To control: aphids, flea beetles, leafhoppers and whiteflies.

To kill gypsy moth caterpillars and adults, adult Japanese

BEST AVAILABLE COPY

393
beetles and Eastern tent caterpillars on contact:
On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats, and small flying moths: Apply using a good space sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

ROUSSEL UCLAF

Roussel Uclaf Corporation
95 Chestnut Ridge Road
Montvale, New Jersey 07645