



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

Oct 26 1994

CHRISTINA M. LAWES
ROUSSEL UCLAF CORP.
95 CHESTNUT RIDGE ROAD
MONTVALE, NJ. 07645

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 03/11/94 in Response to PR Notice 93-7
EPA Reg. No. 10370-236
FORD'S PERMETHRIN 10%

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

Ford's

PERMETHRIN CONCENTRATE

- * *RECOMMENDED FOR COMMERCIAL/AGRICULTURAL USE AND MOSQUITO ABATEMENT DISTRICT OFFICIALS.*
- * *FOR USE OUTDOORS AND IN HOMES AND NON-FOOD AREAS OF COMMERCIAL BUILDINGS, INSTITUTIONS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS, HOTELS, RESTAURANTS, KENNELS AND FOOD/FEEED PROCESSING ESTABLISHMENTS.*
- * *NOT FOR USE IN USDA MEAT AND POULTRY PLANTS.*
- * *KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES*

ACTIVE INGREDIENT:

*Permethrin: (3-phenoxyphenyl) methyl (±)
cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethyl-
cyclopropanecarboxylate. 10.0%

**INERT INGREDIENTS: 90.0%

100.0%

*Cis/trans ratio: Max. 65% (±) trans and min. 35% (±) cis.
**Contains aromatic petroleum distillate.

EPA REG. NO. 10370-236

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Due to the presence of solvent, gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove person to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with plenty of soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

See Side Panel For Additional
ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 26 1994

NET CONTENTS:

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
10370-236

ROUSSEL UCLAF



Roussel Uclaf Corporation
95 Chestnut Ridge Road
Montvale, New Jersey 07645
2365 110493TK

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Remove and wash clothing before reuse. Avoid contamination of food and feed.

Wear full-faced gas mask with canister type recommended for general insecticide protection for applying indoors as a space spray or fog.

Environmental Hazards

This pesticide is highly toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical Hazards

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

Do not use this products in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of the Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage And Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. **CONTAINERS ONE GALLON AND SMALLER:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

CONTAINERS LARGER THAN ONE GALLON: Metal Containers - Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic Containers - Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Glass Containers - Triple rinse (or equivalent) then dispose of in a sanitary landfill or by other approved state and local procedures.

Do not allow spray treatment to drift on pastureland, cropland, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture.

To prepare dilutions, the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. Do not hold dilutions for more than 24 hours.

For maximum effectiveness, a combination of localized application and space treatment is recommended. Remove pets, birds and cover fish aquaria before spraying.

Do not use in food areas of food-handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

NON-FOOD/FEED AREAS including (but not limited to) garbage rooms, lavatories, floor drains (to sewers) entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

INDOOR USE AREAS:

| | |
|--------------------------|---------------------|
| Bakeries | Kennels |
| Beverage Plants | Meat Packing Plants |
| Canneries | Motels |
| Flour Mills | Office Buildings |
| Food Processing Plants | Railroad Cars |
| Grain Elevators | Restaurants |
| Granaries | Schools |
| Homes | Ships Holds |
| Hospitals | Supermarkets |
| Hotels | Truck Trailers |
| Industrial Installations | Warehouses |

AS A SURFACE SPRAY: For initial cleanup of severe insect infestation, dilute at a rate of 1 part concentrate in 19 parts water (6.7 ounces per gallon).

For normal infestations, dilute 1 part concentrate in 100 parts water (1.28 ounces per gallon).

To control *Cockroaches, Waterbugs, Palmetto Bugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Millipedes, Centipedes, Sow Bugs, Pill Bugs, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Saw-toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Mealworms, Grain Mites And Cadellles:* Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. Spray bookcases for silverfish. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

TREATMENT OF FLYING INSECTS RESTING PLACES

TO AID IN CONTROL OF: Mosquitos, Wasps, Gnats, Skippers: Treat walls, ceilings, moldings, screens, door frames, window frames, beams, light cords and similar resting places using the above dilution of 1-19.

TO CONTROL: Mosquitos, Gnats, Small Flying Moths. For best results, use truck mounted equipment and apply when there is a light breeze of approximately 5 miles per hour. Spray from the windward side so the fog will be carried across the area to

be treated. Treat shrubbery and vegetation where mosquitoes may rest.

Dilute 1 part concentrate in 19 parts water and apply at a rate of 2.5 pints per acre (0.9 ounce per 1,000 sq. ft.), using truck mounted conventional misting equipment such as Rotomist models 51C, 91 or 100HT traveling at 5 m.p.h. To treat a 300 foot swath, calibrate equipment to deliver 0.95 gallon per minute. Under no conditions should open water be treated with this product. Not to be used within 100 feet (30 meters) of lakes and streams.

OUTDOOR USE AREAS:

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes, or climate modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

| | | |
|----------------------|--------------|--------------------|
| Corrals | Gardens | Recreational Areas |
| Drive-In Restaurants | Golf Courses | Urban Areas |
| Drive-In Theaters | Parks | Zoos |
| Feedlots | Playgrounds | |

FOR TEMPORARY REDUCTION OF ANNOYANCE FROM: Flies, Gnats, Mosquitos. In drive-in restaurants, drive-in theaters, golf courses, urban areas, parks, playgrounds and other recreational areas: Dilute 1 part concentrate in 40 parts water (3.2 fluid ounces per gallon of water). Direct the spray into tall grass, shrubbery and around lawns, where these pests may hover or rest. Apply while the air is still and avoid wetting foliage. Application should be made prior to use. Repeat as necessary.

In the treatment of corrals, feedlots, swing lots and zoos: Cover any exposed water, drinking fountains and animal feed before applications. Fill the area with mist and direct the spray into tall grass, shrubbery and around lawns where the above pests may hover or rest. Apply while the air is still and avoid wetting foliage. In zoos, prevent exposure of reptiles to the product. Repeat as necessary.

To kill *Gypsy Moth Caterpillars And Adults, Adult Japanese Beetles, and Eastern and Forest Tent Caterpillars* on contact: Combine 3.2 fluid ounces of concentrate per 1 gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Allow all treated articles to thoroughly dry before use. Repeat treatment as needed.

RESIDUALLY CONTROLS FLEAS AND TICKS ON DOGS FOR 35 DAYS. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor coverings. Fresh bedding should be placed in animal quarters following treatment. To kill *Fleas and Ticks* on dogs, and to obtain protection against reinfestation, dilute 1 part

concentrate to 100 parts water (1.28 ounces per gallon). Using protective gloves (mitts), wet the animal by dipping or spraying. **DO NOT TREAT PUPPIES LESS THAN FOUR WEEKS OLD. DO NOT USE THIS OR ANY OTHER PESTICIDE ON SICK, OLD, OR DEBILITATED PETS.**

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 19 parts water (6.7 fluid ounces per gallon). Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of *House Flies, Fruit Flies, Gnats, Mosquitos, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths and Tobacco Moths:* Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1,000 cu. ft. of space. Keep area closed for at least 10 minutes. Vacate areas after treatment and ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label: Dilute 1 part concentrate in 19 parts water (6.7 ounces per gallon). Apply using conventional, mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at a rate of 1 ounce per 1,000 cu. ft. of space.

TREATMENT ON ANIMALS AND ANIMAL QUARTERS: To aid in control of: *Horn Flies, Face Flies, Houseflies, Mosquitoes, Gnats, Stable Flies, Horse Flies, Deer Flies, Blood Sucking Lice, Poultry Lice, Bedbugs, Mites, Fleas and Ticks.* Dilute this product at a rate of 1 part concentrate in 19 parts water (6.7 ounces per gallon).

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs, and horses from horn flies, face flies, houseflies, mosquitoes, and gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To control *Fleas, Ticks, and Lice:* Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on puppies less than 4 weeks old.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

OUTDOOR APPLICATION

Dilute at a rate of 1 part concentrate in 19 parts water (6.7 ounces per gallon).

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamental to kill Japanese beetles hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leaf Miners.*

May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against *Whiteflies, Aphids, Spider And Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.*

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.