

**CLEAN
CROP**

ACCEPTED

JAN 23 1989

Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for this
pesticide registered under
EPA Reg. No. ~~34704-104~~

Transferred to
10370-278

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

CAUTION

Harmful if Swallowed. Avoid contact with skin.
Wash thoroughly after using. Avoid breathing of
spray mist. Keep out of reach of children. Keep
container closed and away from heat and open
flame. Do not combine emulsifiable liquids with
wettable powders in the same spray tank unless
previous use of the materials being combined has
proven them to be physically compatible.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic
invertebrates, and aquatic life stages of amphibians. Do not
apply directly to water except as specified on this label.
Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed
to direct treatment on blooming crops or weeds. Do not apply
this product or allow it to drift to blooming crops or weeds
if bees are visiting the treatment area.

**PHYSICAL AND
CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**STATEMENT OF
PRACTICAL TREATMENT**

If Swallowed—Call a physician or Poison Control
Center immediately. Drink one or two glasses of
water and induce vomiting by touching the back
of throat with finger. Repeat until vomit fluid is clear.
Do not induce vomiting or give anything by mouth
to an unconscious person.
If Inhaled—Remove victim to fresh air. Apply ar-
tificial respiration if indicated.
If On Skin—Remove contaminated clothing and
wash affected areas with soap and water.
If In Eyes—Flush eyes with plenty of water. Call
a physician if irritation persists.
Note to Physician: This product may cause
cholinesterase inhibition. Atropine is antidotal.
2-PAM may be effective as an adjunct to atropine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.
Apply specified dosage (tablespoons) of this pro-
duct per 2 gallons of water unless specified other-
wise. Begin applications when insects first appear
and repeat as needed.

STORAGE AND DIS-

STORAGE: Store in original co
preferably in a locked area. Keep c
ed when not in use. Store in a co
Do not use food or drink contain
or storage.

PESTICIDE DISPOSAL: Securely
container in several layers of ne
discard in trash.

CONTAINER DISPOSAL: Do nc
lainer. Rinse thoroughly before
trash.

ALMONDS: For control of Aphids a
Borer, use 6 to 12 tbs. per 11½ ga
No time interval between last ap
harvest.

**APPLES, APRICOTS, AVOCADOS,
FIGS, NECTARINES, PEACH
PECANS, PLUMS, PRUNES, QUIN
7 tbs. per 10 gallons of water to c
Thrips, Spider Mites, Mealybugs,
Eye-Spotted Bud Moth, and Orange
Willamette Mite and Two-Spotted M
make at least 2 applications 10 to
in summer months. For Scale on p
to completion of hatching, about la
Cherry Fruit Fly repeat applications
injury may occur to McIntosh a
varieties of apples in summer sprai
tain varieties of sweet cherries part
Northwest, and fruit spotting may
tainies. Do not apply within 1 day
pears; within 3 days on apples, i
plums, prunes, and quinces; with
apricots, avocados, nectarines, &
limitation on Pecans.**

**ASPARAGUS, BEETS, CARROTS,
DANDELIONS, EGGPLANT, END
HORSERADISH, LEEKS, PARSLEY
PEAS, PEPPERS, SALISFY,
SPINACH, SWISS CHARD, AND W
Use 1½ to 3 tbs. per 2 gals. of wa
Aphids, Spider Mites, Thrips,
Asparagus Beetle, Pea Weevil, and
Beetle. Do not apply within 1 day
Asparagus; within 3 days on Garl
Leeks, Peas, Peppers, and Shallots,
on Beets, Carrots, Dandelions, End
Salisfy, Swiss Chard, Spinach, W
within 21 days on Parsley.**

BEANS: For the control of Aphids,
Thrips, Leafhoppers, Asparagus
Weevil, and Mexican Bean Beetle use
per 2 gallons of water. Do not apply
of harvest.

MALATHION 2 HOME LAWN AND GARDEN SPRAY

ACTIVE INGREDIENTS:

Malathion*	25.0%
INERT INGREDIENTS**	75.0%
TOTAL	100.0%

*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.
(contains 2 lbs. malathion per gallon)

**This product contains xylene range aromatics.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panels for Additional Precautionary Statements

EPA REG. NO. 34704-104

EPA EST. NO. 34704-CO-2

NET CONTENTS _____ GALLONS

Malathion Solution
RC MA2783

ORM-A

BEST AVAILABLE COPY

BERRIES: CANEBERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, and Raspberries), **CRANBERRIES, & STRAWBERRIES:** Use 4½ to 6 tbs per 2 gallons of water to control Aphids, Thrips, Spider Mites, Whiteflies, Leafhoppers, Rose Scale, Lygus Bugs, Spittlebugs, Leafrollers, Backheaded Fireworms, and Strawberry Root Weevil. For Strawberry Root Weevil apply to soil surface before planting and work into top 6 to 8 inches. Do not apply within 3 days of harvest on Cranberries and Strawberries and within 1 day on Caneberries.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, KALE, MUSTARD GREENS, TURNIPS: Use 3 to 6 tbs. per 2 gallons of water to control Aphids, Imported Cabbage-worm, & Cabbage Looper. Do not apply within 7 days of harvest on Brussels Sprouts, Cabbage, Kale, Mustard Greens, and within 3 days on Broccoli and Turnips.

CANTALOUPE, PUMPKINS, SQUASH, MELONS: For control of Aphids, Leafhoppers, Pickworm, Spider Mites, Cucumber Beetles. Use 2¼ to 2½ tbs per 1 gallon of water. Do not apply within 3 days of harvest on cantaloupes and melons, 1 day on squash.

CELERY & CUCUMBERS: Use 4 tbs. per 2 gallons of water to control Aphids, Spider Mites, and Cucumber Beetle. Do not apply within 7 days of harvest on Celery or within 1 day on Cucumbers.

CITRUS: (GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGERINES, TANGELOS, AND KUMQUATS): Use 3 to 6 tbs per 2¼ gals. of water to control Aphids and Scale Crawlers. Use 3¼ tbs per 4½ gals. of water to control Thrips. Do not apply when trees are in bloom. Do not apply within 7 days of harvest.

CORN: Aphids, Corn Rootworm Adults, Sap Beetles, Thrips, Young Grasshoppers. Use 3 to 4 tbs per 2 gallons of water. Do not apply within 5 days of harvest or forage use. Caution: injury may occur in the whorl and silk stages with emulsifiable liquids.

GRAPES: Use 4 to 7 tbs per 2 gals. of water to control Grape Leafhoppers, Mealybugs, and Spider Mites. Injury may occur to Ribier, Italia, Cardinal, and Almeria grape varieties when Malathion is applied after the clusters appear. Do not apply within 3 days of harvest.

LETTUCE: Use 3 to 6 tbs per 2 gals. of water to control Aphids and Leafhoppers. Do not apply within 7 days of harvest on Head-Lettuce and within 14 days on Leaf-Lettuce.

ADULT MOSQUITO & FLY CONTROL: Apply 1½ tbs per 2 gallons of water for control of Adult Flies and Mosquitoes. Repeat as necessary. Treat shrubbery and vegetation where mosquitoes may result. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shorelines may be treated. See environmental hazards. Repeat ap-

plications should be made as necessary.

ONIONS: Use 4½ tbs per 2 gallons of water to control Thrips. Do not apply within 3 days of harvest.

PEPPERS: Aphids, Pepper Maggot. Use 4 tbs per 2 gallons of water. Do not apply within 3 days of harvest.

POTATOES: Aphids, Blister Beetles, False Chinch Bugs, Leafhoppers, Mealybugs. Use 3 to 9 tbs per 2 gallons of water.

TOMATOES: Aphids, Spider Mites: Use 3 to 4½ tbs per 2 gallons of water. Drosophila Flies: Use 6 tbs per 2 gallons of water in full coverage application to fruit and foliage. Do not apply within 1 day of harvest.

FLOWERS, ORNAMENTALS AND SHADE TREES: Follow general directions using 3 tbs per 2 gallons of water. Aphids on Elms, Linden Maple; European Pine Shoot Moth on Pines; Japanese Beetle (adults) on Roses, Flowers, Flowering ornamentals; Leaf Hoppers on Hawthorn, Poplar, Willow; Mealybugs & Whiteflies on Azalea, Rhododendron, Laurel, Andromeda, Taxus; Thrips (exposed) on Roses and Flowers; Tarnished Plant Bug on ornamental fruit trees and flowers; Sawflies on Ash, Larch, Pines, Spruce; Lacebug on Rhododendron, Azalea, Sycamore. Use 2-4 ounces per 10 gallons for the following insects: Leaf Miner on Birch, Boxwood (Apply early to mid May. Repeat in 10 days); Tent Caterpillars on Flowering ornamentals. (Apply when scale crawlers appear on the following insects: Oak Kermes on Oak, Fletcher Scale on Taxus, Juniper, Arborvitae, Florida Juniper Scale on ornamental Evergreens, Euonymus Scale on Pachysandra; Pine Needle Scale on Pines.)

NOTICE

Platte warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte. In no case shall Platte be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Platte makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
180 SO. MAIN STREET
FREMONT, NEBRASKA 68025

BEST AVAILABLE COPY