

**AgrEvo**

A company of Hoechst and Schering, Berlin

PM 14

10370-291

1.43

50% Malathion Insect Spray

- * Homeowner Lawn and Garden Pest Control. For Homeowner Use Only
- * Controls Many Insect Pests Including: Aphid, Red Spider Mites, Thrips, Scales, Leafhoppers, Whitefly, Mealybugs and Tent Caterpillars.

ACTIVE INGREDIENT:

Malathion: O,O-dimethyl phosphorodithioate of diethyl
mercaptosuccinate.....

50.00%

INERT INGREDIENTS*

50.00%

100.00%

ACCEPTED

OCT 26 1995

Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended for the pesticide
registration system
EPA Reg. No. 10370-291

*This product contains xylene range aromatics

EPA Reg. No. 10370-291

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN**WARNING****STATEMENT OF PRACTICAL TREATMENT**

If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected area with soap and water.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: This product may cause cholinesterase inhibition. Atropine by injection is antidotal. 2-Pam may be effective as an adjunct to atropine.

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals****Warning**

May be fatal if swallowed. May be absorbed through the skin. Avoid contact with skin, eyes or clothing. Do not breath spray mist. Wash thoroughly after handling. Avoid contamination of food or feed stuff. Do not permit children or pets to go onto sprayed grass until sprayed grass until spray has completely dried. Keep out of reach of domestic animals.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in areas near application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical & Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water food or feed by storage or disposal.

Storage: Keep container tightly sealed during storage. Do not use, pour or store near heat or open flame.

Disposal: Do not reuse empty container. Wrap container in several layers of newspaper and put in trash container.

50% Malathion Insect Spray controls many common garden insects. Use any type of garden sprayer to apply. Make new dilution for every use. Spray thoroughly to cover both upper and lower leaf surfaces and all infested portions of plants.

Do not use on ferns. May injure foliage of certain varieties of apples and cherries. Apply this product only as specified on this label.

Ornamentals (Roses, Azaleas, Camellias, Evergreens and many other flowers and shrubs): Aphids, tent caterpillars: 2 teaspoons per gallon of water. Red spider mites, mealybugs, woolly aphids, whitefly, thrips, rose leafhoppers: 1 table spoon per gallon of water. Black scale, soft brown scale, Monterey pine scale: 1 1/2 to 2 tablespoons per gallon of water.

Fruits and Nuts: Number between parentheses () indicates days required between last spray and harvest. Apples (3), Pears (1), Apricots (7), Cherries (3), Peaches (7), Pecans (1), Walnuts (1), Aphids: 2 teaspoons per gallon of water.

Apples(3), Pears (1), Red spider mites, woolly aphids, whiteflies, thrips, pear psylla: 1 tablespoon per gallon of water.

Apples (3), tent caterpillars: 2 teaspoons per gallon of water.

Citrus Fruits (grapefruit, lemons, limes, oranges, kumquats): Do not apply within 7 days of harvest. Blackscale, California red scale, purple scale, thrips: 2 teaspoons per gallon of water. Do not apply during full bloom.

Vegetables: Numbers between parentheses () indicate days required between last spray and harvest.

Cole crops, including broccoli, Brussels sprouts, cabbage, and cauliflower (7), peas (3), potatoes (1), squash (1), melons (1), tomatoes (1), aphids: 2 teaspoons per gallon of water. Beans (1), aphids: 1 tablespoon per gallon of water.

Premise Treatment: Animals become re-infested unless the pet's yard and lawns are treated. Fleas, ticks; Remove manure or debris from these areas and spray using 10 tablespoons (5 oz.) in 1 gallon of water and spray at a rate of 1 gallon of mixture per 1,000 square feet. Repeat treatment in 3 to 4 weeks if necessary.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health
95 Chestnut Ridge Road
Montvale NJ 07645