



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
10370-310

Date of Issuance:
APRIL 18, 1997

NOTICE OF PESTICIDE:
 x Registration
 Reregistration

Term of Issuance:
Conditional

(under FIFRA, as amended)

Name of Pesticide Product:
INTERCEPT H&G READY-TO-USE INSECT CONTROL

Name and Address of Registrant (include ZIP Code):

AGROEVO ENVIRONMENTAL HEALTH
95 Chestnut Ridge Road
Montvale, NJ 07645

160/58701
17/1

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c) (7) (A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c) (5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 10370-310."
- b. Delete the claim "Can be used where kids and pets play." Claims as to the safety of the pesticide in any form are not acceptable. Delete the claim "Less is more, etc...." Vague or non-specific terms implying unstated benefits related to the composition of a product are not acceptable.
- c. Revise the environmental hazards statement to read as follows:

"This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water."

Signature of Approving Official:

George T. LaRocca
George T. LaRocca
Product Manager (13)

Date:

APR 18 1997

3. Submit three (3) copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure



AgrEvo™

A company of Hoechst and Schering Berlin

3/7

InterCept H&G Ready-to-Use Insect Control, Flower, Vegetable, Lawn and Garden Spray

- * Ready-to-Use.
- * It's the Least You Can Do For Your Garden. ?
- * Easy to Use
- * For Use on Plants.
- * Gentle to Most Plants.
- * Less is More, Low Dose with Maximum Effectiveness ✓
- * Safe to Use on Over 60 Varieties of Ornamentals*
- * Can be Used Where Kids and Pets Play* ✓
- * For Controls of Vegetable Pests
- * Controls Most Ornamental Pests
- * Controls Most Garden Insect Pests
- * Use Indoors and Outdoors.
- * For Use In and Around the Home.
- * For Control of Various Plant Pests.
- * For Plant Pest Control
- * Water-based. No Unpleasant Odor.
- * Use Around Buildings, Homes, Flower and Vegetable Gardens, and on Fruit, Nut and Ornamental Trees.

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 18 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

10370-310

* When Used According to label instructions.

ACTIVE INGREDIENT:

*Permethrin.....0.02%

INERT INGREDIENTS:.....99.98%

Total 100.00%

*(3-phenoxyphenyl)methyl (±)cis/trans 3-(2,2-dichloroethenyl)2,2-dimethylcyclopropane
carboxylate. Cis/trans ratio: min. 35% (±)cis and max. 65% (±)trans.
InterCept H&G is a trademark of AgrEvo Environmental Health.

EPA Reg. No. 10370-

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

See side panel for additional Precautionary Statements.

Net Contents:

4/7

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

For use in and around the home. Turn nozzle to spray position, point away from face and squeeze trigger to spray.

INDOOR USES

INDOOR PLANTS: Use sufficient spray to obtain full coverage of all foliage. Adjust the amount of spray volume used to the size of the plant(s) being treated.

To protect plants in indoor landscaping - Spray directly on insects to kill;
chrysanthemums - *leafrollers, armyworms, cutworms, cabbage worms*
velvet plant - *fungus gnats*

Spray as directed above to control *aphids, armyworms, corn earworms, fallcankerworms, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale crawlers, spring cankerworms, and exposed thrips* on the following plants:

- | | | |
|-----------------|------------------|-------------------|
| Ageratum | English Ivy | Poinsettia |
| Ardisia | Fuchsia | Portulaca |
| Aster | Gladiolus | Pothos |
| Azalea | Grape ivy | Prayer Plant |
| Baby's breath | Hypoestes | Purple Passion |
| Begonia | Ivy | Rabbits foot fern |
| Birds-nest fern | Juniper | Rose |
| Bleeding Heart | Lily | Schefflera |
| Carnation | Marigold | Snapdragon |
| Chrysanthemum | Orchid | Snake Plant |
| Coleus | Palm | Statice |
| Crown of thorns | Pansy | Velvet Plant |
| Cockscomb | Peperomia | Verbena |
| Cyclamen | Petunia | Zinnia |
| Dracaena | Philodendron | |
| Dumbcane | Piggy-back Plant | |

If application is made indoors, remove infested plants from living area prior to application. Spray plants in an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exist a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.

Repeat applications may be necessary to control citrus blackflies on non-bearing citrus and adult Japanese beetles on roses

OUTDOOR USES

PESTS ON THE OUTDOOR SURFACE OF BUILDINGS: To temporarily control *house flies, mosquitoes, cluster flies, gnats, flying moths*. For outdoor use only as an aid in reducing annoyance from these insects. Spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects alight or congregate. Repeat as necessary.

OUTDOOR ORNAMENTALS: Spray upper and lower leaf surfaces. Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. Adjust the amount of spray volume used to the size of the plant(s) being treated. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as needed.

Controls *aphids, cicadas, corn earworms, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, bagworms, spider mites, exposed thrips* and *whitefly* on the following plants:

Ageratum	Gladiolus	Pea Shrub
Arizona Cypress	Gold Bells	Petunia
Aster	Hypoestes	Philodendron
Azalea	Ivy	Poinsettia
Begonia	Lilac	Rhododendron
Coleus	Marigold	Rose
Common Ninebark	Mock Orange	Snapdragon
Common Snowberry	Nannyberry	Taxus
Exacum	Orshid	Wax Plant
Euonymous	Palm	Weeping Fig
English Ivy	Pansy	Zinnia

May also be used on such plants as African violets, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, fuschia, geraniums, crassula, grape vines (ornamental), laurels, stocks and wandering Jews.

ORNAMENTAL TREES AROUND THE HOME: Use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: *pine moths*, *pine needle miners*, *pine beetles* and *needle scales* on pines; *budworms* on conifers; *oakworms* on oak; *fall webworms* on cherry, elm and honeysuckle; *aphids* on tulip poplar; *citrus black flies* on non-bearing citrus; *tussock moths* on Douglas and grand firs; *Leafminers* on birch; *elm leaf beetle* and *elm spanworms* on elms.

FOR HOME VEGETABLE GARDENS: Spray upper and lower leaf surface. Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. Apply no more than 13 ounces over 100 linear feet of row, (13 ounces per 100 square feet).

Vegetable	Target Pests	Use Directions
Asparagus (pre-harvest)	cutworms asparagus beetle	Spray plants as needed to ensure uniform coverage. Can be applied up to 3 days prior to harvest.
(post-harvest)	larval stages of asparagus beetle; tarnished plant bug; lygus bugs and adult Japanese beetle.	Treat ferns or bush growth after spear harvest when insects are present.
Celery - see Leafy Vegetables		
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	beet armyworm; diamond back moth; imported cabbageworm; cabbage- aphid, cabbage looper.	Apply ever 5 days or as needed. Can be applied up to one day prior to harvest.
Corn - see Sweet Corn		
Cucurbits: Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon	aphids, cabbage looper, cucumber beetle (adults), cutworms, leafhoppers, leafminers, melonworm, pickleworm, plant bugs, squash bugs,	Spray plants as needed to ensure uniform coverage. Can be applied up to the day of harvest.
Eggplant	Colorado potato beetle, vegetable leaf miner	Spray plants as needed to ensure uniform coverage. Can be applied up to 3 days prior to harvest.
Horseradish	imported crucifer weevil	Apply as needed. Can be applied up to 22 days prior to harvest.
Leafy Vegetables: amaranth, arrugula, celery, celtuce, chervil, corn salad, chrysanthemum (edible-leaved and garland), cress (garden and upland), dandelion, dock, endive, fennel, lettuce (head and leaf), orach, parsley, purslane (garden and winter), rhubarb, spinach, Swiss chard	alfalfa looper, aphids, beet armyworm, cabbage looper, corn ear worm, cutworms, european corn borer, fall armyworm, green cloverworm, southern armyworm, tobacco budworm, vegetable leaf miner.	Spray every 5 to 10 days as needed. Can be applied up to 1 day prior to harvest.
Melons - see Cucurbits		

Vegetable	Target Pests	Use Directions
Potatoes	beet armyworm; colorado potato beetle; potato leafhopper; potato aphid; potato tuberworm; cabbage looper; cutworms; European corn borer; potato psyllid; tarnished plant bug.	Spray plants as needed. Can be applied up to 7 days prior to harvest.
Peppers	vegetable leafminer; cabbage looper; flea beetle; corn earworm; pepper weevil; European corn borer.	Spray plants as needed. Can be applied up to 3 days prior to harvest.
Spinach	beet armyworm; corn earworm; cabbage looper; cutworms; European corn borer; green cloverworm and leafhoppers.	Spray plants as needed for uniform coverage. Can be applied up to 7 days prior to harvest.
Sweet Corn	corn earworm; European corn borer; fall armyworm; Southern armyworm.	Spray plants every 5 days and when needed. Can be applied up to 1 day prior to harvest.
Tomatoes	beet and Southern armyworm; tomato pinworm; tomato fruitworm; horn worms; granulate cutworms; colorado potato beetle; cabbage looper	Spray plants when needed for uniform coverage. Can be applied up to day of harvest.

HOME OWNER GROWN FRUIT AND NUT TREES

Apply InterCept H&G Ready-to-Use Insect Control, Flower, Vegetable, Lawn and Garden Spray to trees until foliage is dripping wet. Make sure to spray the undersides of leaves and into dense foliage.

Trees (fruit or nut)	Target Pests	Use Directions
Almonds	navel orange worm; peach twig borer.	Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	plum curculio; redbanded leafroller; rosyapple aphid; spotted tentiform leafminer; tarnished plant bug; white apple leafhopper; olique-banded leafroller; green fruitworm.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
Peaches	lesser peachtree borer; peach twig borer; plum curculio; tarnished plant bug; green fruit worm; oriental fruit moth.	Apply when insects first appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears	pear psylla; green fruitworm and codling moth.	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health
95 Chestnut Ridge Road
Montvale, NJ 07645