

SANITROL MICROBIOCIDE

FOR CONTROL OF BACTERIA AND FUNGI IN CANE SUGAR MILLS

ACCEPTED
AUG 31 1971

UNITED STATES DEPARTMENT OF AGRICULTURE
FUNGICIDE AND ANTIMICROBIAL DIVISION
FOR ECONOMIC POISON REGISTRATION
REGISTRATION NO. 10378-1

Comments

**NET CONTENTS
APPROX.
55 GALLONS
NET WEIGHT
500 POUNDS**

ACTIVE INGREDIENTS . . . 35%
**Nabam (Disodium Ethylene
 Bisdithiocarbamate) . . 13%**
**Sodium Dimethyl Dithio-
 carbamate 12%**
Ethylenediamine 10%
INERT INGREDIENTS 65%
TOTAL 100%

USDA REG. NO 10378-1

WARNING: KEEP OUT OF REACH OF CHILDREN

May cause eye injury and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling the concentrate. Avoid breathing of vapor. In case of contact, immediately flush with large amount of water; for eyes wash with for at least 15 minutes and get medical attention. Harmful if swallowed. Use with adequate ventilation. Keep container closed when not in use. Remove contaminated clothing and wash before reuse.

In case of swallowing, induce vomiting with mustard and warm water and immediately call a physician for further attention.

DIRECTIONS FOR USE:

HODAG SANITROL MICROBIOCIDE is a liquid product which can be metered into the cane juice by means of a small pump. It is important to feed the product in such a manner as to insure complete circulation of treated juice throughout the mill tandem. The product should be added continuously to the juice at a minimum rate of 10 ppm up to but not exceeding a maximum of 20 ppm based on the solution as received in the drum. The feed rate must not exceed 43 pounds (4.5 gallons) per 1000 tons of sugar cane.

When used regularly in this manner, inversion losses, bad odors and gummy deposits resulting from the growth of micro-organisms are reduced to a minimum.

See Bulletin No. D11-568 For Additional Use-Directions

FOR INDUSTRIAL USE ONLY

HODAG CHEMICAL CORPORATION - SKOKIE, ILLINOIS U.S.A.

ACCEPTED WITH COMMENTS