DIRECTIONS

Designed for use as a periodic shock treatment. Use at the rate of one quart per 5000 Gallons of pool water. This will add 5 p.p.m. of available chlorine to the water, which will help prevent algae from growing and kill many bacteria that may be present. Pools should be shocked once every week. Add directly to pool water near inlet to aid dispersion. Do not enter pool until chlorine residuals drop between 1.0 p.p.m. and 1.5 p.p.m. as determined by test kit.


This product degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.

E. P. A. Reg. No. 10380 - 9

NET CONTENTS: 1 GALLON

Q-PAK CORPORATION

2145 Mc CARTER HIGHWAY, NEWARK, N. J. 07104



Sodium Hypochlorite

10%

INERT INGREDIENTS

90%

CAUTION

KEEP OUT OF REACH OF CHILDREN.

SEE ANTIDOTE STATEMENT AND OTHER PRECAUTIONS ON BACK / SIDE PANEL.



ACTIVE INGREDIENT:

Sodium Hypochlorite

10%

INERT INGREDIENTS

90%

CAUTION
KEEP OUT OF REACH OF CHILDREN.

SEE ANTIDOTE STATEMENT AND OTHER PRECAUTIONS ON BACK / SIDE PANEL.

CAUTION HARMFUL IF SWALLOWED

Keep out of reach of children.

his product will severely irritate skin and byes.

Avoid transfer to food or beverage containers.

Avoid spillage on clothing as deterioration and discoloration may occur.

Keep container tightly capped in an upright position. Store in a cool place.

Rinse empty container thoroughly with water and discard.

Do not mix with any other chemicals. Violent chemical reactions may occur. Add directly to pool water as per instructions.

FIRST AID

EXTERNAL: In eyes, flush with water, im-

mediately, at least 5 minutes.

Call a physician.

On skin, flood and rinse immediately with copious amounts of water. If irritation persists, con-

tact a physician.

INTERNAL: Drink milk, white of egg or feed

cooked cereal. Call a physician.

MOYCO GERMICIDAL SOLUTION

is a most effective chemical solution for the cold disinfection of dental and medical instruments, prosthetic and orthodontic appliances, plastic and rubber apparatus, thermometers, porcelain bowls and non-porous surfaces such as stainless steel table tops, etc.


DIRECTIONS FOR USE

Wash instruments to be disinfected with warm water and soap to remove all pus, blood or exudates. Thoroughly rinse away all soap with clear water, immerse in properly diluted Germicidal Solution (1 ounce to 3 quarts of water) for 10 to 15 minutes. Leave instruments in solution until ready to be used. Replace solution frequently.

DANGER: This liquid is highly concentrated, therefore it can cause severe eye irritation. Do not get in the eyes or on the skin. Avoid contamination of food. Harmful, or fatal, if swallowed. First Aid: Immediately flood eyes or skin with plenty of water for 15 minutes. For eyes, call a physician. Remove or wash contaminated clothing before re-use. If swallowed, do not induce vomiting. Drink large quantities of fluid.

IMPORTANT

See literature for additional information before using.



H.I. 4624-1835

CONCENTRATED GERMICIDAL SOLUTION

FOR THE COLD DISINFECTION OF DENTAL AND MEDICAL INSTRUMENTS

Active Ingredients:

Diisobutylcresoxyethoxyethyl dimethyl benzyl ammonium. chloride: o-phenylphenol ethoxynonylphenol complex..10% lnert Ingredients90% (Sodium nitrite...1%)

DANGER

Keep Out of Reach of Children

(See reverse panel for further cautions.)

1 PINT

Manufactured for:
THE J. BIRD MOYER COMPANY, INC.
Philadelphia, Pa. 19132

U.S.D.A. Reg. No. 10438-1. made in U.S.A.

SEP 181970

UNDER THE FENERAL INSFOTICIDE
FUNGICIET OF TWEETHINGLE ACT
FOR LOCIETARY POURTER NOTE TO ATTACHED COMMENTS.

1-88401