

PRECAUTIONARY STATEMENTS

WARNING

Hazards to Humans and Domestic Animals

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food. Do not use on humans or permit children to go onto sprayed grass until spray has completely dried.

Statement of Practical Treatment

If swallowed - Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Note to Physician: Gastric lavage is indicated. Contains a petroleum-based solvent which presents aspiration hazard. Atropine sulfate and 2-PAM are specific antidotes.

Environmental Hazards

This product is toxic to birds, fish, and wildlife including waterfowl. Birds and waterfowl feeding or drinking on treated areas may be killed. Apply this product only as specified on label. Do not exceed maximum permitted label rates, as rates above the those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended apply a minimum of 0.25" of water immediately after application. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries (except for effluence from treated watercross beds). Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency for information pertaining to areas having threatened endangered species. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

General Information

Lesco Diazinon 500 is an emulsifiable solution which when diluted with water according to directions, is intended for the control of insect pests on turf and ornamentals. Do not use in barns or dwellings. Do not use on livestock. In the directions for use which follow, best control is obtained when application is made at the first sign of insects. Application should be repeated as necessary to maintain control.

Directions for Use and Conditions of Sale

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Failure to follow the Directions for Use and Precautions on this label may result in poor insect control or crop injury. **CHEMIGATION:** Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

A. STORAGE INSTRUCTIONS:

Store in original container only, separate from other pesticides, fertilizer, food and feed. Do not store near open flame or heat sources.

B. PESTICIDE DISPOSAL INSTRUCTIONS:

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or residue is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

C. CONTAINER DISPOSAL INSTRUCTIONS:

METAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
PLASTIC - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Range, Pasture, and Grassland Insects

Unless otherwise stated, Lesco Diazinon 500 should always be applied as a spray in a minimum of 1 gal. of water per acre for aerial applications and a minimum of 5 gals. of water per acre for ground applications.

Bermudagrass and Forage Grasses (other than Range and Pasture Grasses)

Thrips, Whiteflies, Grasshoppers: Apply 1 pt. per acre. Dairy and beef cattle and sheep (1) may be grazed immediately following application; (2) may be fed green forage immediately following cutting. Forage grasses may be cut for hay 7 days following treatment. Do not repeat application for 30 days.

BEST AVAILABLE COPY

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



DIAZINON® 500 INSECTICIDE

This product must not be used on golf courses and sod farms FOR CONTROL OF CERTAIN INSECTS ON ORNAMENTALS AND LAWNS as listed on label.

LESCO DIAZINON 500 CONTAINS 4 LBS DIAZINON PER GALLO Diazinon® Registered Trademark Ciba-Geigy

ACTIVE INGREDIENTS:

O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	48.00%
Xylene-Range Aromatic Solvent	38.00%
INERT INGREDIENTS:	16.00%
TOTAL	100.00%

E.P.A. Reg. No. 10404-11

E.P.A. Est. No. 10404-OH-02

See Directions for Use on Panel at Right.

KEEP OUT OF REACH OF CHILDREN WARNING
See additional warning statements on side panel.

Manufactured by:

LESCO, INC.

20085 Lake Road, P.O. Box 16915
Rocky River, Ohio 44116

Ornamental Insect Control

To control certain insects on Arborvitae, Azalea, Birch, Borwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladiol, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Plum, Poplar, Rhododendron, Rose, Spruce, and Willow; apply the recommended rates indicated as follows:

Insects	Rate Per 3 gals. water	Rate Per 120 gals. water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Diglossid Leaf-miners, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leafhoppers, Privet Mites, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Mites, Webworms, Whiteflies	1/2 fl. oz.*	1 pt.**
Apple-and-Thorn Skeletonizers, Cottonweaver Webworms, Fall Webworms, Hamlock Chermes, Oak Looper, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars	1 1/2 fl. oz.	3 pts.**

* 1 fl. oz. = 2 tablespoons

** 1 fl. oz. = 1 pint

Do not use on Ferns, Forsythia, Hibiscus, Papaya, Pines, and Gardens since injury to the plants may occur. Mimosa Webworms: Mix 1/2 fl. oz. in 3 gals. water (or 1 pt. in 100 gals. water) and apply as a thorough spray when adults first appear. Make a second application at 4-5 week intervals.

NET CONTENTS: 1 GAL. 8897349 5 GAL. 8897350 30 GAL. 8897358 55 GAL. 8897348

NOTIFICATION TO PRINTER

Please Reference the Front Panel of This EPA Labeling EPA no longer classifies the Solvent in this product as an Active Ingredient therefore entirely remove the Solvent Name from the Ingredient Statement. Adjust the figure to add percentage which is now shown for the solvent to 100% total percentage of the INERT INGREDIENT. Proof ASA

Lawn Insect Control

To control insects listed below, apply recommended rates. For control of Ant Eaters, Fleas, Millipedes, Sowbugs, and Springtails (Collembola), which enter the house, spray a 5 ft. band of soil around the house, as well as the h Repeat application if necessary.

Insects	Rate/1000 sq. ft.	
	Lesco	Water
Lawn Chinch Bugs	2-3 fl. oz. or 3-6 fl. oz.	25 gal.*

*Or, make application in 3 gals. of water per 1000 sq. ft. and thoroughly water

Insects	Rate/1000 sq. ft.	
	Lesco	Water
Ants, Armyworms, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Fire Flies, Lawn Billbugs, Sod Webworms (Lawn Moths), Sowbugs	2-4 fl. oz.	3 gals.

For lawn thoro treat dry. C

Brown Dog Ticks, Bermuda Mites, Chiggers, Fleas, Leafhoppers	1 1/2 fl. oz.	3 gals. shrub
--	---------------	---------------

Insects	Rate/1000 sq. ft.	
	Lesco	Water
Millipedes	8 fl. oz.	3 gals.
Rhodesgrass Scales	5 fl. oz.	25 gals.*

* Or make application in 3 gals. water per 1000 sq. ft. and thoroughly water

Range, Pasture, Ditch Banks, Roadsides, Wasteland, Non-crop Areas, Grasshoppers: Apply 1/2-1 pint per acre in water or oil (minimum of 1 gal. preferably in the nymphal stage. Thorough coverage of the foliage is essential. Dairy and beef cattle and sheep (1) may be grazed immediately following immediately following cutting, and (2) may be fed hay provided grass is not cut solution or for 30 days if the spray is an oil solution. Do not spray livestock direct nor repeat application for 30 days.

Boxelder Bugs: Spray bugs outside the home with 1 1/2-2 1/2 fl. oz. (2 1/2-5 table; thoroughly spray around foundation, windows, doors, tree trunks, and in insects hide.

Billbugs, White Grub (such as Japanese Beetle Larvae, European Chafer, Sout of lawn. Water grass thoroughly after application. For grub control make applic October. For billbug control apply when activity is first observed or when the damage from this insect.

Hypoxandrus Weevil (New York only): Apply 3 fl. oz. per 1000 sq. ft. of grass. Water hyperandrus in mid-April and again in mid-May.

Atlantic Spherotheca (Atlantic Beetle) (Ohio, West Virginia only): Apply 4 oz. summer when damage is first noticed. Water grass thoroughly with 1/2-1/4 inch of summer or early fall if a second generation occurs.

WARRANTY

LESCO warrants that this insecticide conforms to the chemical described dance with label directions under normal conditions, this insecticide is not Since timing, method of application, weather, plant and soil conditions, m factors affecting the use of this product are beyond our control, no warrant product contrary to label directions or under conditions which are strictl user assumes all risks of any such use.