LESCO DIAZINON 14G

FOR CONTROL OF CERTAIN SOIL INSECTS

56 Pounds **Net Weight**

ACTIVE INGREDIENT: O, Odiethyi O-(2-isopropyl-6methyl-4-pyrimidinyl) Phosphorothicate... INERT INGREDIENTS:....

LESCO Diazinon 14G is a granular insecticide Diazinon Registered Trademark Ciba-Geigy

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

It is a violation of Federal law to use this product in a manner inconsistent with its label.

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller.

Setter makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

LAWN INSECTS

Lawn Chinch Bugs: Apply 12-1 lb. per 1,000 sq. ft. (21/2-5 lbs. per 5,000 sq. ft). Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf such as St. Augustine grass, use 1-112 lbs per 1,000 sq. ft. (5-71/2 lbs. per 5,000 sq. ft.)

Billbugs; White Grubs such as Japanese Beetle Larva, European Chater, and Southern Chater: Apply 1 lb. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.

Hyperodes Weevil (New York only): Apply 3/4 lb. per 1.000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Cyclone Spreader Settings	Once Over	Twice Over	Spread Width
1/2 pound per 1,000 sq. ft.	2.00	1.75	' 6 feet
1 pound per 1,000 sq. ft.	2.50	2.00	6 teet
1-1/2 pounds per 1,000 sq. ft.	2.75	2.25	6 feet

KEEP OUT OF REACH OF CHILDREN. **CAUTION**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through skin. Avoid contact with skin, syes, or clothing. Avoid contamination of feed. Wash thoroughly after handling product and before eating or smoking. Keep children and pets off treated area until lawn has

Practical Treatment

If swallowed, drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin - Remove contamined clothing and wash affected areas with soap and water.

Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

Note to Physician: If symptoms of chotinergic poisoning occur, administer atropine therapeutically. If swallowed, lavage stomach or induce vomiting

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Birds feeding on treated areas may be killed. this product only as specified on this label

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal. Open diimping is prohibited

2. PESTICIDE DISPOSAL

Pesticide, so(ay mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies

PATE RESPOSAL

Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place

Consult federal, state of local disposal authorities for approved alternative procedures such as limited open to

Distributed by: LAKESHORE EQUIF NT & SUPPLY CO. ELYRIA, OHIO 44035 ()

E.P.A. Reg. No. 10104-12 E.P.A. Est. No. 10404-OH-01