

LESCO 2.32 GRANULAR IN

ACCEPTED
MAY 31 1981
Dursban is a registered trademark of the Dow Chemical Co.
Fungicide, Insecticide, and Fertilizer
an amended and registered pesticide
registered under EPA Reg. No. 10404-15

E

FOR PROFESSIONAL TURF MAINTENANCE OPERATORS ONLY

A Dursban[®]-coated corn cob granular for control of Ants, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Earwigs, Hyperodes (Turfgrass) Weevils, Sod Webworms (Lawn Moths), Sowbugs (Billbugs) and White Grubs on lawns and around buildings. Manufactured specifically for use in cyclone-type spreaders.

ACTIVE INGREDIENTS:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate) 2.32%

INERT INGREDIENTS: 97.68%

TOTAL 100.00%

*Dursban - Trademark of the Dow Chemical Co.
U.S. Patent No. 3,244,586

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label instructions.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Read evenly at a rate of 1 pound per 1,000 square feet. After application, water lawn using 30 gallons per 1,000 square feet. To help obtain uniform distribution, apply one half the required amount in one direction; then apply the balance at a right angle.

To control **ANTS, BROWN DOG TICKS, CRICKETS, CUTWORMS, EARWIGS, SOD WEBWORMS, CHINCHBUGS and SOWBUGS**—spread uniformly at a rate of 1 pound per 1,000 sq. ft. in northeastern states; make first application in May or June or when insects first appear. Re-treat as needed, usually in July or August.

NOTE: To reduce initial infestation or reinfestation of buildings by Earwigs, treat a band of ground 6 to 10 feet wide around and adjacent to the buildings where these pests are active.

To control **HYPERODES WEEVILS**—apply as above except at twice the rate of two pounds per 1,000 square feet. Apply over suspected problem area in mid-April and again in mid-May.

To control **WHITE GRUBS** (larvae of Japanese Beetle and European Chafer), use 4 pounds per 1,000 square feet. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local agricultural extension service specialist for proper time to treat in your area.

Cyclone Spreader Setting	Once Over	Twice Over	Spread Width
1 pound per 1,000 sq. ft.	2.75	2.50	6 feet
2 pounds per 1,000 sq. ft.	3.00	2.75	6 feet
4 pounds per 1,000 sq. ft.	—	3.00	6 feet

APPROXIMATE SPREADER SETTINGS

IMPORTANT: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed. Do not take internally. Avoid contact with skin, eyes, or clothing. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Avoid breathing of dust. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Keep away from food, foodstuffs and domestic water supplies.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: remove victim to fresh air. Apply respiration if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water.

If in eyes: Flush out eyes for at least 15 minutes with water. Call a physician if irritation persists.

NOTE TO PHYSICIAN

Active ingredient is cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleansing of equipment or disposal of wastes. Do not apply when run off is likely to occur. Use this product only as specified on this label.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures under the Resource, Conservation and Recovery Act.

3. CONTAINER DISPOSAL

Dispose of bags according to approved Federal, state or local procedures under the Resource, Conservation and Recovery Act.

4. GENERAL

Consult Federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

Distributed by: Lakeshore Equipment & Supply Co.
300 South Abbe Rd., P.O. Box 4015
Elyria, Ohio 44036

EPA Reg. No. 10404-15
EPA Est. 10404-OH-01

NET WEIGHT:
50 LBS./22.7 KILOGRAMS

26