LESCO 36-0-0 Fertilizer with Dursban*

CONTAINS DURSBAN*

For the control of soil insects such as ants, armyworms, cutworms, sod webworms, chinche bugs, earwigs, sowbugs, crickets and certain white grubs.

ACTIVE INGREDIENT:

Chlorpyrifos (0, 0-diethyl-0-(3,5,6-trichloro-2-	
pyridyl) phosphorothioate	2.32%
INERT INGREDIENTS:	97.68%
TOTAL	100.00%

DURSBAN* - Registered Trademark of Dow Chemical Company U.S. Patent No. 3,244,586

KEEP OUT OF REACH OF CHILDREN CAUTION

Read all directions before using.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Ma ba harmful if swallowed. Avoid eye contact.

Wr thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep away from food, foodstuffs and domestic water supplies. Keep children and pets off of treated area until watered into soil and soil is dry.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician.

If on skin: Wash thoroughly with soap and water.

If in eyes: Flush eyes with plenty of water. If irritation persists, call a physician.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Fish and crustaceans may be

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE

To control chinch bugs, sod webworms, ants, armyworms, crickets, cut worms, earwigs, and sowbugs, uniformly apply one (1) pound of this product to cover 1,000 square feet of non-crop turfgrase area. Treat when insects first appear; re-treat as needed. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply balance at a right angle. Immediately after application, thoroughly water the treated turf to wash the insecticide into the zone of insect infestation.

To control certain white grubs (istvae of European chafer and Japanese beetle), interchiptor and on the product per 1,000 square feet. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash insectivities and only in the time of the time of the grubs are young and actively feeding near soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

36-0-0 with DURSBAN* SPREADER SETTINGS

Cyclone	Once over T	wice over	Spread Width
1 3b./1.000 sq. ft.	3.75	3.25	7 feet
4 ibs/1,000 sq. ft.	5.50	4.50	7 feet

APPROXIMATE SPREADER SETTINGS.

IMPORTANT: These settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application.

DISPOSAL DIRECTIONS

Do not reuse empty container. Wrap container and put in trash collection

Lakeshore makes no warranty expressed or implied concerning the use of this product other than indicated on the

- killed if their waters are contaminated with this product. Do not contaminate water by cleaning equipment or disposal of wastes.
- naterial when such use and handling are contrary to label instructions.

EPA Reg. No. 10404-17 EPA Est. No. 10404-0H-01 Net Weight 50 Lbs. (22.7 kg.)

Manufactured by: Lakeshore Equipment & Supply Co. 300 South Abbe Road, P.O. Box 4015 Elyria, Ohio 44036

