

JAN - 4 1994

Mr. Julius C. Prinzo
LESCO, Inc.
20005 Lake Road
Rocky River, Ohio 44116

Dear Mr. Prinzo:

Subject: Revised Directions and Precautions
LESCO 32-5-7 Fertilizer with Dursban
EPA Registration No. 10404-29
Your Application Dated October 29, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely,

DHE

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

LESCO

BEST AVAILABLE COPY

JAN - 4 1984
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registration number 10904-29

32-5-7 Fertilizer with Dursban

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Controls Ants, Armyworms (Beet, Fall, Yellowstriped), Brown Dog Ticks*, Centipedes, Chinch Bugs*, Clover Mites, Cutworms, Crickets, Earwigs, European Crane Fly larvae*, Flary Skipper, Fleas*, Grasshoppers, Leafhoppers, Millipedes, Sod Webworms* (Lawn Moth), Sow Bugs, Springtails, Turfgrass Weevil* (Hyperodes), White Grubs* (Species of Black Turfgrass Attenius, European Chafer, Southern Chafer, Japanese Beetle and June Beetle larvae), Bluegrass Billbugs*, Fire Ants*.

*See Specific Directions

ound Treats up to 333 sq.ft.

Residual insect control for parks - golf courses - athletic fields - home lawns - for outside of homes - apartments - condominiums - warehouses - hospitals - schools - food handling establishments - commercial and office buildings

Active Ingredient:
Chlorpyrifos: [D,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 0.74%
Inert Ingredients 99.26%
TOTAL 100.00%

GUARANTEED ANALYSIS

Total Nitrogen (N) 32.00%
Available Phosphoric Acid (P₂O₅) 5.00%
Soluble Potash (K₂O) 7.00%
Dursban® is a Registered Trademark of DowElanco
EPA Reg. No. 10404-29 EPA Est. No. 1C404-OH-04

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENTS OF PRACTICAL TREATMENT

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. **DO NOT** induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention.
If in Eyes: Flush eyes with plenty of water. **If on Skin:** In case of contact, remove contaminated clothing and wash affected area with soap and water.
Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and water supplies. As recommended, water lawn after application and allow to dry before permitting children and pets on treated area.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Use only as directed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal
Storage: Store product in its original labeled container in a cool, dry, locked place out of reach of children.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

*COMMERCIAL CONTROL

The following paragraphs correspond to insects followed by asterisk (*) on the above front panel. Insects not followed by an asterisk (*) may be controlled by following the instructions FOR USE ON LAWNS, PARKS, etc. Repeat as needed for control.

TICKS AND FLEAS

Use 3.0 lb./1,000 sq. ft. for area control of these insects infesting non-crop land areas such as roadsides, footpaths, and trails, picnic and camping sites, parks, recreational areas and around dwellings where these pests are present and are creating a nuisance. **DO NOT** allow use of treated area during applications. For Fleas (out-of-doors, only) also treat sleeping quarters and other areas frequented by pets such as beneath porches and along fencing.

CHINCH BUGS

Apply 3.0 to 4.8 lbs./1,000 sq.ft. when insects appear or their damage is imminent. To prevent damage apply as soon as daily temperatures reach 80° and repeat every 2 months. Use low rate for minimum infestations; higher rate for heavy infestations. For best results, product should be watered in after application.

SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30-day intervals may be needed for new infestations. Apply at rate of 3.0 lbs./1,000 sq.ft.

TURFGRASS WEEVIL

Use 6.1 lbs./1,000 sq.ft. Treat suspected problem areas in mid-April and again in mid-May or as recommended by your local agricultural extension service.

WHITE GRUB SPECIES - CRANE FLY LARVAE

To control White Grubs (Black Turfgrass Attenius, European Chafer, Southern Masked Chafer, Japanese Beetle, and June Beetle Larvae), use 6.1 lbs. to treat 1,000 sq.ft. After application, apply enough water (1/2") to thoroughly wash the insecticide into the turf. For best results, remove thatch first and treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local agricultural extension service specialist for best time to treat. European Crane Fly larvae will be controlled using 4.6 lbs./1,000 sq.ft. or will be controlled simultaneously with the same application being applied for control of "White Grubs" i.e., (6.1 lbs./1,000 sq.ft.). Consult your agricultural agent for proper time to apply. For best results, first remove thatch. Water the granules into the soil.

BLUEGRASS BILLBUGS

Use 6.1 lbs./1,000 sq.ft. for Billbug grubs. Remove thatch before treatment. Wash granules into soil as for White Grubs. Consult your local turf specialist for timing of applications.

IMPORTED FIRE ANT CONTROL - MOUND TREATMENT

To control Fire Ants in lawns, turf and premise areas, apply 1/2 cup (4 ounces) evenly over and around each mound. For best results, do not disturb the mound before or during the application. Apply in cool weather (65-80°F) or in early morning or late evening. Treat new mounds as they appear.

FOR USE ON LAWNS, PARKS, GOLF COURSES, PLAYGROUNDS, AND ALL OTHER ORNAMENTAL AND RECREATIONAL TURF AREAS: Apply uniformly to home lawns, golf courses, parks, playgrounds, and any other ornamental and recreational turf and non-crop land areas using any hand or power operated gravity or rotary type spreader designed for application of granules.

SURFACE INSECTS

For Surface Insects: Ants, Armyworms (Beet, Fall, Yellowstriped) Ticks, Centipedes, Cutworms, Crickets, Earwigs, Flary Skipper, Grasshoppers, Leafhoppers, Millipedes, clover Mites, Sowbugs, and springtails, use 2.4 lbs./1,000 sq.ft. (105 lbs./acre). Treat when insects first appear. For best results, product should be watered in after application. Observe delayed watering for Sod Webworm control.

SOIL INSECTS

For Soil Insects as listed on this label, increase the rate to 6.1 lbs./1,000 sq.ft. immediately after application, sprinkle the turf with 30 gallons of water/1,000 sq.ft. (1,300 gallons/acre) to move the granules down into the zone of insect infestation. Repeat treatment as necessary.

SPREADER SETTINGS*

Lbs. Product /1,000 sq.ft.	Lbs. A.I. /Acre	EXHAUSTIVE Spreader Width	Tire Ho/e	Approximate Setting
3.0 lbs	1.0 lbs.	7 feet	Open	E 1/4
6.1 lbs	2.0 lbs	7 feet	Open	G 1/2

* Above spreader settings are approximate. Rate of walk, moisture, and spreader condition can vary the setting necessary to apply the correct amount.

INSECT CONTROL USING BAND AND PERIMETER TREATMENT

Use for Crawling Insect Control Outside of Homes, Apartments, Condominiums, Warehouses, Hospitals, Schools, Food Handling Establishments, and Commercial Buildings: To protect against insects entering the indoors, a band of granules may be placed around the outside of buildings to kill Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails, and Spiders before they attempt to enter. Use 4.0 lbs. to 6.1 lbs. to treat 500 sq.ft. (as a band 5 ft.x100 ft. long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days.

WARRANTY

LESCO, Inc. warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purposes since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Net Weight 50 Pounds

#009681

LESCO, Inc.
2005 Lake Road, P.O. Box 16915
Rocky River, Ohio 44116

Rev. 10/83WP