

Controls Ants, Armyworms (Beet, Fall, Yellowstriped), Brown Dog Ticks*, Centipedes, Chinch Bugs*, Clover Mites, Cutworms, Crickets, Earwigs, Fiery Skipper, Fleas*, Grasshoppers, Leafhoppers, Milipedes, Sod Webworms* (Lawn Moth), Sowbugs, Springtails, Fire Ants*. *See Specific Directions

1 Pound Treats 333 Square Feet

Residual insect control for golf course turf, road medians and outside of industrial sites such as warehouses and food processing establishments.

ACTIVE INGREDIENT:

Chlorpyrifos [0,0-diethyl-0-(3,5,6-trichloro-2-pyridyl)
phosphorothioate]0.74%
INERT INGREDIENTS: 99.26%
Total

KEEP OUT OF REACH OF CHILDREN CAUTION

	FIRST AID
If inhaled	 FIRST AID Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth- to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-
	 Call a poison control center or doctor for further treatment advice.
lf in eyes	 Hold eye open and rinse slowly and gently with water for 15-10 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
	HOT LINE NUMBER
f	NOTE TO PHYSICIAN
Chloropyrifos is a cholir antidote by injection.	nesterase inhibitor, treat symptomatically. Atropine is

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS


CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and water supplies. Water lawn after application and keep the public, including children and pets, off treated area until the lawn is dried.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of equipment water by cleaning of equipment or disposal of equipment washwaters. Use only as directed

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal Storage: Store product in its original labeled container in a cool dry, locked place out of reach of children. Pesticide Disposal: Wastes resulting from the use of this product may be

disposed of on site or at an approved waste disposal facility. <u>Container Disposal:</u> Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay our of smoke.

*COMMERCIAL CONTROL + - - -

The following paragraphs correspond to insects followed by asterisk (*) on the above front panel. Insects not followed by an asterisk (*) may be controlled by following the instructions FOR USE ON GOLF COURSE. Repeat as needed for control.

TICKS AND FLEAS

Use 3.0 lb/1,000 sq ft for area control of these insects, infesting non-crop land areas; golf course turf only, road medians and industrial sites where these pests

are present and are creating a nuisance. DO NOT allow use of treated area during applications.

CHINCH BUGS

Apply 3.0 lb/1,000 sq ft when insects appear or their damage is imminent. To prevent damage, apply as soon as daily temperatures reach 80°F and repeat every 2 months. For best results, product should be watered-in after application. SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30-day intervals may be needed for new IMPORTED FIRE ANT CONTROL - MOUND TREATMENT infestations.

FOR PROFESSIONAL USE ONLY

To control Fire Ants in lawns, turf, and premise areas, apply 1/2 cup (4 ounces) evenly over and around each mound. For best results, do not disturb the mound before or during the application. Apply in cool weather (65-80°F) or in early morning or late evening. Treat new mounds as they appear.

FOR USE ON GOLF COURSE TURF ONLY, ROAD MEDIANS AND INDUSTRIAL SITE TURF AREAS: Apply uniformly to golf course turf, road medians and industrial site turf and non-crop land areas using any hand- or power-operated gravity- or rotary-type spreader designed for application of granules.

SURFACE INSECTS

For Surface Insects: Ants, Armyworms (Beet, Fall, Yellowstriped), Ticks, Centipedes, Cutworms, Crickets, Earwigs, Fiery Skipper, Grasshoppers, Leafhoppers, Millipedes, Clover Mites, Sowbugs, and Springtails, use 2.4 lb (1.000 sq ft (105 lb/acre). Treat when insects first appear. For best results, Chesen delayed watering for Sod <u>/1.000 sq ft</u> (105 lb/acre). Treat when insects first appear. For best results, product should be watered in after application. Observe delayed watering for Sod Webworm control.

Э:

SUGGESTED SPREADER SETTINGS* Recommended Rates in lb per 1,000 sq ft

242

Spreader	Ground Speed	XX lb	XX lb	XX lb	XX lb
888	XX mph	XX	XX	XX	XX
bbb	XX mph	XX	XX	XX	XX
CCC	XX mph	XX	XX	XX	XX
ddd	XX mph	XX	XX	XX	XX
fff	XX mph	XX	XX	XX	XX
<u>9</u> 99	XX mph	XX	XX	XX	XX
hhh	XX mph	XX	XX	XX	XX
iii	XX mph	XX	XX	XX	XX

*IMPORTANT: Above spreader settings are approximate. Rate of walk, moisture, and spreader condition can vary the setting necessary to apply the correct amount.

INSECT CONTROL USING BAND AND PERIMETER TREATMENT

Use of CONTROL USING BAND AND PERIMETER TREATMENT Use for Crawing Insect Control Outside of Industrial Plant Sites such as Warehouses and Food Processing Establishments. To protect against insects entering the indoors, a band of granules may be placed around the outside of buildings to kill Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails, and Spiders before they attempt to enter. Use 3.0 Ib to treat 1,000 sq ft (as a band 5 ft x 200 ft long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days.

WARRANTY

LESCO, Inc. warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normat conditions, this product is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 15885 Sprague Rd., Strongsville, OH 44136-1799, referring to the item number found on this bag.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-a1.com

Dursban is a registered trademark of Dow AgroSciences LLC

LESCO and Poly Plus are registered trademarks of LESCO Technologies, LLC o:\requi\product labels\masters\epa documents\dursban074.doc

Created: XX/XX/XX (XXXXXX) Rev. XX/XX/XX XXXX

Net Weight: XX lb (XX kg) PA REG. NO. 10404-29 EPA EST. NO. 10404-OH-4 (M), 10404-FL-1 (S) First letter of lot code indicates manufacturing site.



FXXXX Manufactured by: LESCO, Inc. 15885 Sprague Road Strongsville, Ohio 44136-1799

758073xxxxx

 Poly Plus Coating More Cost Effective performance than Standard Sulfur Coated Urea (SCU) Proven Technology 50% Poly Plus SCU for controlled growth and extended color. Controls surface insects such as chinch bugs, sod webworms and bluegrass billbugs.
 More Cost Effective performance than Standard Sulfur Coated Urea (SCU) Proven Technology 50% Poly Plus SCU for controlled growth and extended color.
50% Poly Plus SCU for controlled growth and extended color.
 Controls surface insects such as chinch bugs, sod webworms and bluegrass billbugs.
Contains micronutrients for improved plant vigor.
Greater Performance and Savings
Free flowing product makes X-X-X easy to handle and apply
• Fewer applications reduce mowing frequency